

35

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
17 May 2001 (17.05.2001)

PCT

(10) International Publication Number
WO 01/34627 A1

(51) International Patent Classification⁷: C07H 21/04,
21/02, C07K 5/00, 14/00, C12Q 1/68, C12N 1/21, 15/63,
15/85, 15/86

(74) Agents: HOOVER, Kenley, K. et al.; c/o Human Genome
Sciences, Inc., 9410 Key West Avenue, Rockville, MD
20850 (US).

(21) International Application Number: PCT/US00/30628

(22) International Filing Date:
8 November 2000 (08.11.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/164,744 12 November 1999 (12.11.1999) US
60/215,140 30 June 2000 (30.06.2000) US

(71) Applicant (for all designated States except US): HUMAN
GENOME SCIENCES, INC. [US/US]; 9410 Key West
Avenue, Rockville, MD 20850 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): RUBEN, Steven, M.
[US/US]; 18528 Heritage Hills Drive, Olney, MD 20832
(US). KOMATSOULIS, George, A. [US/US]; 9518
Garwood Steet, Silver Spring, MD 20901 (US). BAKER,
Kevin, P. [GB/US]; 14006 Indian Run Drive, Darnestown,
MD 20878 (US). YOUNG, Paul, E. [US/US]; 122 Beck-
with Street, Gaithersburg, MD 20878 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- With international search report.
- Before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments.

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.



WO 01/34627 A1

(54) Title: 28 HUMAN SECRETED PROTEINS

(57) Abstract: The present invention relates to novel human secreted proteins and isolated nucleic acids containing the coding re-
gions of the genes encoding such proteins. Also provided are vectors, host cells, antibodies, and recombinant methods for producing
human secreted proteins. The invention further relates to diagnostic and therapeutic methods useful for diagnosing and treating dis-
eases, disorders, and/or conditions related to these novel human secreted proteins.

28 Human Secreted Proteins

Field of the Invention

This invention relates to newly identified polynucleotides, polypeptides encoded by these polynucleotides, antibodies that bind these polypeptides, uses of
5 such polynucleotides, polypeptides, and antibodies, and their production.

Background of the Invention

Unlike bacterium, which exist as a single compartment surrounded by a membrane, human cells and other eucaryotes are subdivided by membranes into many functionally distinct compartments. Each membrane-bounded compartment, or
10 organelle, contains different proteins essential for the function of the organelle. The cell uses "sorting signals," which are amino acid motifs located within the protein, to target proteins to particular cellular organelles.

One type of sorting signal, called a signal sequence, a signal peptide, or a leader sequence, directs a class of proteins to an organelle called the endoplasmic
15 reticulum (ER). The ER separates the membrane-bounded proteins from all other types of proteins. Once localized to the ER, both groups of proteins can be further directed to another organelle called the Golgi apparatus. Here, the Golgi distributes the proteins to vesicles, including secretory vesicles, the cell membrane, lysosomes, and the other organelles.

20 Proteins targeted to the ER by a signal sequence can be released into the extracellular space as a secreted protein. For example, vesicles containing secreted proteins can fuse with the cell membrane and release their contents into the extracellular space - a process called exocytosis. Exocytosis can occur constitutively or after receipt of a triggering signal. In the latter case, the proteins are stored in
25 secretory vesicles (or secretory granules) until exocytosis is triggered. Similarly, proteins residing on the cell membrane can also be secreted into the extracellular space by proteolytic cleavage of a "linker" holding the protein to the membrane.

Despite the great progress made in recent years, only a small number of genes encoding human secreted proteins have been identified. These secreted proteins
30 include the commercially valuable human insulin, interferon, Factor VIII, human growth hormone, tissue plasminogen activator, and erythropoietin. Thus, in light of

the pervasive role of secreted proteins in human physiology, a need exists for identifying and characterizing novel human secreted proteins and the genes that encode them. This knowledge will allow one to detect, to treat, and to prevent medical diseases, disorders, and/or conditions by using secreted proteins or the genes that encode them.

Summary of the Invention

The present invention relates to novel polynucleotides and the encoded polypeptides. Moreover, the present invention relates to vectors, host cells, antibodies, and recombinant and synthetic methods for producing the polypeptides and polynucleotides. Also provided are diagnostic methods for detecting diseases, disorders, and/or conditions related to the polypeptides and polynucleotides, and therapeutic methods for treating such diseases, disorders, and/or conditions. The invention further relates to screening methods for identifying binding partners of the polypeptides.

Detailed Description

Definitions

The following definitions are provided to facilitate understanding of certain terms used throughout this specification.

In the present invention, "isolated" refers to material removed from its original environment (e.g., the natural environment if it is naturally occurring), and thus is altered "by the hand of man" from its natural state. For example, an isolated polynucleotide could be part of a vector or a composition of matter, or could be contained within a cell, and still be "isolated" because that vector, composition of matter, or particular cell is not the original environment of the polynucleotide. The term "isolated" does not refer to genomic or cDNA libraries, whole cell total or mRNA preparations, genomic DNA preparations (including those separated by electrophoresis and transferred onto blots), sheared whole cell genomic DNA preparations or other compositions where the art demonstrates no distinguishing features of the polynucleotide/sequences of the present invention.

In the present invention, a "secreted" protein refers to those proteins capable of being directed to the ER, secretory vesicles, or the extracellular space as a result of a signal sequence, as well as those proteins released into the extracellular space without necessarily containing a signal sequence. If the secreted protein is released into the extracellular space, the secreted protein can undergo extracellular processing to produce a "mature" protein. Release into the extracellular space can occur by many mechanisms, including exocytosis and proteolytic cleavage.

In specific embodiments, the polynucleotides of the invention are at least 15, at least 30, at least 50, at least 100, at least 125, at least 500, or at least 1000 continuous nucleotides but are less than or equal to 300 kb, 200 kb, 100 kb, 50 kb, 15 kb, 10 kb, 7.5 kb, 5 kb, 2.5 kb, 2.0 kb, or 1 kb, in length. In a further embodiment, polynucleotides of the invention comprise a portion of the coding sequences, as disclosed herein, but do not comprise all or a portion of any intron. In another embodiment, the polynucleotides comprising coding sequences do not contain coding sequences of a genomic flanking gene (i.e., 5' or 3' to the gene of interest in the genome). In other embodiments, the polynucleotides of the invention do not contain the coding sequence of more than 1000, 500, 250, 100, 50, 25, 20, 15, 10, 5, 4, 3, 2, or 1 genomic flanking gene(s).

As used herein, a "polynucleotide" refers to a molecule having a nucleic acid sequence contained in SEQ ID NO:X or the cDNA contained within the clone deposited with the ATCC. For example, the polynucleotide can contain the nucleotide sequence of the full length cDNA sequence, including the 5' and 3' untranslated sequences, the coding region, with or without the signal sequence, the secreted protein coding region, as well as fragments, epitopes, domains, and variants of the nucleic acid sequence. Moreover, as used herein, a "polypeptide" refers to a molecule having the translated amino acid sequence generated from the polynucleotide as broadly defined.

In the present invention, the full length sequence identified as SEQ ID NO:X was often generated by overlapping sequences contained in multiple clones (contig analysis). A representative clone containing all or most of the sequence for SEQ ID NO:X was deposited with the American Type Culture Collection ("ATCC"). As

shown in Table 1, each clone is identified by a cDNA Clone ID (Identifier) and the ATCC Deposit Number. The ATCC is located at 10801 University Boulevard, Manassas, Virginia 20110-2209, USA. The ATCC deposit was made pursuant to the terms of the Budapest Treaty on the international recognition of the deposit of
5 microorganisms for purposes of patent procedure.

A "polynucleotide" of the present invention also includes those polynucleotides capable of hybridizing, under stringent hybridization conditions, to sequences contained in SEQ ID NO:X, the complement thereof, or the cDNA within the clone deposited with the ATCC. "Stringent hybridization conditions" refers to an
10 overnight incubation at 42 degree C in a solution comprising 50% formamide, 5x SSC (750 mM NaCl, 75 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65 degree C.

Also contemplated are nucleic acid molecules that hybridize to the
15 polynucleotides of the present invention at lower stringency hybridization conditions. Changes in the stringency of hybridization and signal detection are primarily accomplished through the manipulation of formamide concentration (lower percentages of formamide result in lowered stringency); salt conditions, or temperature. For example, lower stringency conditions include an overnight
20 incubation at 37 degree C in a solution comprising 6X SSPE (20X SSPE = 3M NaCl; 0.2M NaH₂PO₄; 0.02M EDTA, pH 7.4), 0.5% SDS, 30% formamide, 100 µg/ml salmon sperm blocking DNA; followed by washes at 50 degree C with 1XSSPE, 0.1% SDS. In addition, to achieve even lower stringency, washes performed following stringent hybridization can be done at higher salt concentrations (e.g. 5X
25 SSC).

Note that variations in the above conditions may be accomplished through the inclusion and/or substitution of alternate blocking reagents used to suppress background in hybridization experiments. Typical blocking reagents include Denhardt's reagent, BLOTTO, heparin, denatured salmon sperm DNA, and
30 commercially available proprietary formulations. The inclusion of specific blocking reagents may require modification of the hybridization conditions described above, due to problems with compatibility.

Of course, a polynucleotide which hybridizes only to polyA⁺ sequences (such as any 3' terminal polyA⁺ tract of a cDNA shown in the sequence listing), or to a complementary stretch of T (or U) residues, would not be included in the definition of "polynucleotide," since such a polynucleotide would hybridize to any nucleic acid molecule containing a poly (A) stretch or the complement thereof (e.g., practically any double-stranded cDNA clone generated using oligo dT as a primer).

The polynucleotide of the present invention can be composed of any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. For example, polynucleotides can be composed of single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, the polynucleotide can be composed of triple-stranded regions comprising RNA or DNA or both RNA and DNA. A polynucleotide may also contain one or more modified bases or DNA or RNA backbones modified for stability or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications can be made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically, or metabolically modified forms.

The polypeptide of the present invention can be composed of amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres, and may contain amino acids other than the 20 gene-encoded amino acids. The polypeptides may be modified by either natural processes, such as posttranslational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be branched, for example, as a result of

ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched, and branched cyclic polypeptides may result from posttranslation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cysteine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, pegylation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination. (See, for instance, PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York (1993); POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS, B. C. Johnson, Ed., Academic Press, New York, pgs. 1-12 (1983); Seifter et al., Meth Enzymol 182:626-646 (1990); Rattan et al., Ann NY Acad Sci 663:48-62 (1992).)

"SEQ ID NO:X" refers to a polynucleotide sequence while "SEQ ID NO:Y" refers to a polypeptide sequence, both sequences identified by an integer specified in Table 1.

"A polypeptide having biological activity" refers to polypeptides exhibiting activity similar, but not necessarily identical to, an activity of a polypeptide of the present invention, including mature forms, as measured in a particular biological assay, with or without dose dependency. In the case where dose dependency does exist, it need not be identical to that of the polypeptide, but rather substantially similar to the dose-dependence in a given activity as compared to the polypeptide of the present invention (i.e., the candidate polypeptide will exhibit greater activity or not more than about 25-fold less and, preferably, not more than about tenfold less activity, and most preferably, not more than about three-fold less activity relative to the polypeptide of the present invention.)

Many proteins (and translated DNA sequences) contain regions where the amino acid composition is highly biased toward a small subset of the available

residues. For example, membrane spanning domains and signal peptides (which are also membrane spanning) typically contain long stretches where Leucine (L), Valine (V), Alanine (A), and Isoleucine (I) predominate. Poly-Adenosine tracts (polyA) at the end of cDNAs appear in forward translations as poly-Lysine (poly-K) and poly-Phenylalanine (poly-F) when the reverse complement is translated. These regions are often referred to as “low complexity” regions.

Such regions can cause database similarity search programs such as BLAST to find high-scoring sequence matches that do not imply true homology. The problem is exacerbated by the fact that most weight matrices (used to score the alignments generated by BLAST) give a match between any of a group of hydrophobic amino acids (L,V and I) that are commonly found in certain low complexity regions almost as high a score as for exact matches.

In order to compensate for this, BLASTX.2 (version 2.0a5MP-WashU) employs two filters (“seg” and “xnu”) which “mask” the low complexity regions in a particular sequence. These filters parse the sequence for such regions, and create a new sequence in which the amino acids in the low complexity region have been replaced with the character “X”. This is then used as the input sequence (sometimes referred to herein as “Query” and/or “Q”) to the BLASTX program. While this regime helps to ensure that high-scoring matches represent true homology, there is a negative consequence in that the BLASTX program uses the query sequence that has been masked by the filters to draw alignments.

Thus, a stretch of “X”s in an alignment shown in the following application does not necessarily indicate that either the underlying DNA sequence or the translated protein sequence is unknown or uncertain. Nor is the presence of such stretches meant to indicate that the sequence is identical or not identical to the sequence disclosed in the alignment of the present invention. Such stretches may simply indicate that the BLASTX program masked amino acids in that region due to the detection of a low complexity region, as defined above. In all cases, the reference sequence(s) (sometimes referred to herein as “Subject”, “Sbjct”, and/or “S”) indicated in the specification, sequence table (Table 1), and/or the deposited clone is (are) the definitive embodiment(s) of the present invention, and should not be construed as

limiting the present invention to the partial sequence shown in an alignment, unless specifically noted otherwise herein.

Polynucleotides and Polypeptides of the Invention

5

FEATURES OF PROTEIN ENCODED BY GENE NO: 1

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. dbj|BAA00321.1| (all information available through the recited accession number is incorporated herein by reference) which is described therein as "polyprotein precursor [Homo sapiens]" A partial alignment demonstrating the observed homology is shown immediately below.

15

```
>dbj|BAA00321.1| polyprotein precursor [Homo sapiens] >pir|JX0061|SAHUP saposin
precursor - human
Length = 527
```

20

Plus Strand HSPs:

Score = 509 (179.2 bits), Expect = 5.0e-66, Sum P(2) = 5.0e-66
Identities = 122/345 (35%), Positives = 201/345 (58%), Frame = +1

25

```
Q: 214 NGPLTFHPRQAFEGALCQDCVRQVSRQLQEAVRSNLTADL---NIQEQCESLGPGGLAVLC 384
      +GP + P+ G +CQDC++ V+ +Q AVR+N T +++E+C+ LGPG+A +C
S: 183 DGPRS-KPQPKDNGDVCQDCIQMVTDIQTAVRTNSTFVQALVEHVKEECDRLGPGMADIC 241
```

30

```
Q: 385 KNYLFQFFVPADQALRLLP---PQELCRKGGFCEELGAPARLT---QVAMDGVPSELEL 543
      KNY+ Q+ A Q + + P+E+C GFC+E+ T +V + + +P+LEL
S: 242 KNYISQYSEIAIQMMHMQDQPKKEICALVGFCEVKEKMPMQTLVPAKVASKNVIPALEL 301
```

35

```
Q: 544 GLPRKQSEMQMKGAVTCEVCMNVVQKLDHWMNSSELMITHALERVCSVMPASITKECI 723
      P K+ E+ K+ V CEVC +V+++ + +N +E I A +++CS +P S+++EC
S: 302 VEPIKKHEVPAKSDVYCEVCEFLVKEVTKLIDNNKTEKEILDADFDMCKSLPKSLSEECQ 361
```

40

```
Q: 724 ILVDITYSPSLVQ-LVAKITPEKVCKFIRLC-GNRRRARAVHDAYAIVPSPPEWDAENQGSF 897
      +VDTY S++ L+ +++PE VC + LC G R A VH V P+ G F
S: 362 EVVDITYGSSILSILLEEVSPVLVCSMLHLCSGTRLPALTVH-----VTQPK-----DGGF 411
```

45

```
Q: 898 CNGCKRLLTVSSHNLSEKSTKRDILVAFKGGCSILPLPI*SKCKHFRHPIRSPCSLRVSK 1077
      C CK+L+ NLE STK++IL A + GCS LP P +C F P + +
S: 412 CEVCKKLVGYLDNRNLEKNSTKQEILAALEKGCFLPDYPYQKQCDQFVAEYE-PVLIEILV 470

Q: 1078 DMMGPRG-CVQEGGGPGTAPRPHLLGTDQXGLAQXAGARTKKAASFGNALQHWQKHGW 1248
      ++M P C++ G P +A +P LLGT++ + + + A+ NA++H ++H W
S: 471 EVMDPSFVCLKIGACP-SAHKP-LLGTEKCIWGPSYWCQNTETAAQCNAVEHCKRHVW 526
```

The segment of dbj|BAA00321.1| that is shown as "S" above is set out in the sequence listing as SEQ ID NO: 171. Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein.

- 5 Assays for determining such activities are also known in the art, some of which have been described elsewhere herein. Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out in the sequence listing as SEQ ID NO: 172 which corresponds to the Q sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed). Moreover,
- 10 fragments and variants of these polypeptides (such as, for example, fragments as described herein, polypeptides at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to these polypeptides and polypeptides encoded by the polynucleotide which hybridizes, under stringent conditions, to the polynucleotide encoding these polypeptides) are encompassed by the invention. Antibodies that bind polypeptides
- 15 of the invention are also encompassed by the invention. Polynucleotides encoding these polypeptides are also encompassed by the invention.

This gene is expressed primarily in healing abdomen wound tissue, 15 days post incision.

- Polynucleotides and polypeptides of the invention are useful as reagents for
- 20 differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include but are not limited to: diseases and/or disorders involving wound healing. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of
- 25 disorders of the above tissues or cells, particularly of healing wound tissues, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g., cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or sample taken from an individual having such a disorder, relative to
- 30 the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Based upon the tissue distribution, this gene product may play roles in wound healing, inflammation, arthritis, tissue rejection, fibrosis, and scarring. Based upon the homology, this gene product may have additional roles. Homology searches reveal that translation products corresponding to this gene share sequence similarity with sulfatide activator protein, also known as SAP-1. The physiological function of SAP-1 is to stimulate hydrolysis of sulfatide by arylsulfatase A in vivo. A genetic defect in the sulfatide activator results in a metabolic disorder similar to classical metachromatic leukodystrophy, which is itself caused by a genetic defect in arylsulfatase A. Thus, polynucleotides, translation products and antibodies corresponding to this gene may have utility in ameliorating similar metabolic defects. Homologous proteins called prosaposins have demonstrated effects on promotion of neurite outgrowth in vitro. Prosaponin and saposin-C also promoted nerve cell myelination ex vivo. Additionally, translation products corresponding to this gene, as well as antibodies directed against these translation products, may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

FEATURES OF PROTEIN ENCODED BY GENE NO: 2

This gene is expressed in tissues that are rapidly proliferating and/or differentiating/dedifferentiating, such as various tumor and fetal tissues. Specifically the transcript is found in the following tissues/cDNA libraries: Soares_testis_NHT; Soares_fetal_liver_spleen_1NFLS_S1; Human Endometrial Tumor; Soares infant brain 1NIB; and to a lesser extent in a variety of normal and transformed tissues and cell lines.

Polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include but are not limited to: proliferative diseases and/or disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of proliferating tissues and/or cells, expression of this

gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g., proliferating, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The presence of the transcript for this gene in tissues/cDNA libraries with high rates of proliferation such as tumors, fetal tissue, placenta, testes, and colon indicates a role for this gene in control of cell proliferation and/or differentiation. Thus, polypeptides and polynucleotides of the invention are useful in the diagnosis and treatment of a variety of cancers and other proliferative disorders. Additionally, translation products corresponding to this gene, as well as antibodies directed against these translation products, may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

FEATURES OF PROTEIN ENCODED BY GENE NO: 3

This gene is expressed in immune system cells; specifically the transcript for this gene has been detected in the following tissues/cDNA libraries: primary dendritic cells, activated monocytes, and T helper cells.

Polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include but are not limited to: diseases and/or disorders of the immune system. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g., immune, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or sample taken from an individual having such a disorder, relative to the standard gene

expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The specificity of this transcript for immune system cells (dendritic cells, monocytes, and T-helper cells) indicates that polypeptides and polynucleotides of the invention are involved in the control and/or modulation of the immune response. As such, it (or antagonists to the protein) might be useful in the diagnosis and/or treatment of a variety of immune disorders, either by increasing a deficient immune response (AIDS, age-related immune suppression, chemotherapy, radiation exposure) or by reducing an inappropriate immune response (lupus and other autoimmune disorders, allergic reactions, etc.). Additionally, translation products corresponding to this gene, as well as antibodies directed against these translation products, may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

15 **FEATURES OF PROTEIN ENCODED BY GENE NO: 4**

This gene is expressed in immune system tissues and cells, such as primary dendritic cells.

Polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include but are not limited to: immune system diseases and/or disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g., immune, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in primary dendritic cells indicates that polypeptides and polynucleotides of the invention are useful for the diagnosis and/or treatment of a variety of immune system disorders. Representative uses are described in the "Immune Activity" and "Infectious Disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. Involvement in the regulation of cytokine production, antigen presentation, or other processes indicates a usefulness for treatment of cancer (e.g. by boosting immune responses). Expression in cells of lymphoid origin indicates the natural gene product would be involved in immune functions. Therefore it would also be useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Furthermore, the protein may also be used to determine biological activity, raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

FEATURES OF PROTEIN ENCODED BY GENE NO: 5

The protein product of this gene is preferentially expressed in immune cells, particularly neutrophils, monocytes, Germinal center B-cells, chronic lymphocytic leukemia, activated T-cells, bone marrow and dendritic cells. In addition, it is expressed to a lesser extent in a variety of normal and transformed tissues and cell types.

Polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include but are not limited to: diseases and/or disorders of the immune system. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g., immune, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The distribution of this transcript, primarily in immune and cancerous tissues (which tend to contain immune cells) indicates that polypeptides and polynucleotides of the invention are useful in immune modulation, and hence in the diagnosis and/or treatment of diseases and/or disorders caused by improper immune function. Such diseases and/or disorders include autoimmune disorders such as lupus and tissue rejection; and diseases caused by immune insufficiencies such as AIDS. Furthermore, the protein product of this gene might be useful in the diagnosis and/or treatment of cancer and other proliferative disorders. Additionally, translation products corresponding to this gene, as well as antibodies directed against these translation products, may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

FEATURES OF PROTEIN ENCODED BY GENE NO: 6

This gene is expressed primarily in spleen from chronic lymphocytic leukemia and leukocytes; specifically in the following tissues/cDNA libraries: Spleen tissue from chronic lymphocytic leukemia patients, anergic and helper T-cells, and
5 neutrophils.

Polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include but are not limited to: diseases and/or disorders of the immune system. Similarly, polypeptides and
10 antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g., immune, cancerous and wounded tissues) or bodily fluids
15 (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The presence of this transcript exclusively in spleen and leukocytes indicates
20 that translation products corresponding to this gene play a role in the control of the immune response. The association with Chronic lymphocytic leukemia indicates that translation products corresponding to this gene might be useful for the diagnosis and/or treatment of this disorder and other related disorders. Further, polynucleotides, translation products and antibodies corresponding to this gene might be useful to
25 stimulate a immune response in other conditions (AIDS, age related immune deficiencies, chemotherapy, radiation exposure, etc) that result in immune suppression. Additionally, translation products corresponding to this gene, as well as antibodies directed against these translation products, may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

30

FEATURES OF PROTEIN ENCODED BY GENE NO: 7

This gene is expressed in primary dendritic cells.

Polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include but are not limited to: diseases and/or disorders of the immune system. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g., immune, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Based on its expression in dendritic cells, this gene and its encoded polypeptides may be useful for the treatment and/or diagnosis of immune system disorders, especially those involving defective antigen presentation such as immune deficiency syndromes as well as autoimmune diseases in which self antigens are recognized (i.e., SLE and RA), and also in the treatment and/or diagnosis of cancer in which antigen presentation is thought to be useful in immune monitoring of neoplasia. More generally, the tissue distribution in primary dendritic cells indicates that polynucleotides, translation products and antibodies corresponding to this gene also strongly indicates a role for this protein in immune function and immune surveillance. Additionally, translation products corresponding to this gene, as well as antibodies directed against these translation products, may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

30 FEATURES OF PROTEIN ENCODED BY GENE NO: 8

This gene is expressed in primary dendritic cells.

Polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include but are not limited to: diseases and/or disorders of the immune system. Similarly, polypeptides and
5 antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g., immune, cancerous and wounded tissues) or bodily fluids
10 (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Based on its expression in dendritic cells, polynucleotides, translation
15 products and antibodies corresponding to this gene may be useful for the treatment and/or diagnosis of immune system disorders, especially those involving defective antigen presentation such as immune deficiency syndromes, as well as autoimmune diseases in which self antigens are recognized (i.e., SLE and RA), and also in the treatment and/or diagnosis of cancer in which antigen presentation is thought to be
20 useful in immune monitoring of neoplasia. More generally, the tissue distribution in primary dendritic cells indicates that polynucleotides, translation products and antibodies corresponding to this gene also strongly indicates a role for this protein in immune function and immune surveillance. Additionally, translation products
25 corresponding to this gene, as well as antibodies directed against these translation products, may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

FEATURES OF PROTEIN ENCODED BY GENE NO: 9

30

This gene is expressed primarily in tissues and cell lines that show increased rates of proliferation. Specifically this transcript is found in the following

tissues/cDNA libraries: Human Esophagus, Cancer; NCI_CGAP_Co9; Human Fetal Dura Mater; Human Hypothalamus, Schizophrenia, Human Chondrosarcoma; Soares breast 2NbHbst; Human ovary; Hodgkin's Lymphoma II; Human Cerebellum; and Soares Pregnant uterus, and to a lesser extent in Testes; Whole 6 Week Old Embryo;

5 Human (HCC) cell line liver (mouse) metastasis, remake; Human Colon Carcinoma (HCC) cell line; Human Umbilical Vein Endothelial Cells, fract. A; Adipocytes, re-excision; Human Fetal Bone, Lung, Cancer (4005163 B7): Invasive, Poorly Diff. Adenocarcinoma, Metastatic; Human Normal Breast; Human Fetal Epithelium (Skin); Human Osteosarcoma; Stratagene lung carcinoma 937218; Human Prostate Cancer,

10 Stage B2; re-excision; Human Activated Monocytes; Human Osteoblasts II; Hemangiopericytoma; Synovial Fibroblasts (control); Human Testes Tumor; Human Amagdyla; Human Placenta; Smooth muscle, control; Pancreas Tumor PCA4 Tu; Human Endometrial Tumor; Colon tumor II; Soares fetal hear NbHH19W; and Soares NhHMPu S1.

15 Polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include but are not limited to: proliferative disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential

20 identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly relating to proliferative disorders, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g., proliferating, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or

25 sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The presence of this transcript in tissues and cell lines with high rates of proliferation (such as fetal and cancerous tissues) indicates a role in growth control.

30 Thus, polynucleotides, translation products and antibodies corresponding to this gene might be useful for the diagnosis and/or treatment of cancer and/or other proliferative disorders. Additionally, translation products corresponding to this gene, as well as

antibodies directed against these translation products, may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

5 FEATURES OF PROTEIN ENCODED BY GENE NO: 10

The protein product of this gene is an exact match to: anaphase-promoting complex subunit 2 [Homo sapiens] (Genbank reference gb|AAF05751.1). An alignment generated by the computer program BLASTX is shown below:

10

```
>gb|AAF05751.1|AF191337_1 (AF191337) anaphase-promoting complex subunit 2 [Homo sapiens]
>sp|AAF05751|AAF05751 Anaphase-promoting complex subunit 2.
Length = 822
```

15

Plus Strand HSPs:

Score = 3967 (1817.7 bits), Expect = 0.0, P = 0.0
Identities = 775/822 (94%), Positives = 775/822 (94%), Frame = +3

20

```
Query:   30 MAAVVVAEGDSDSRPGQELLVAWNTVSTGLVPPAALGLVSSRTSGAVPPKEEELRAAVE 209
          MAAVVVAEGDSDSRPGQELLVAWNTVSTGLVPPAALGLVSSRTSGAVPPKEEELRAAVE
Sbjct:   1 MAAVVVAEGDSDSRPGQELLVAWNTVSTGLVPPAALGLVSSRTSGAVPPKEEELRAAVE 60
```

25

```
Query:  210 VLRGHGLHSVLEEFVEVLQNDLQANISPEFWNAISQCENSADEPQXXXXXXXXXXES 389
          VLRGHGLHSVLEEFVEVLQNDLQANISPEFWNAISQCENSADEPQ          ES
Sbjct:  61 VLRGHGLHSVLEEFVEVLQNDLQANISPEFWNAISQCENSADEPQCLLLLDAGLLLES 120
```

30

```
Query:  390 RLDPYLRSLELLEKWTRLGLLMGTGAQGLREEVHTMLRGVLFSTPRTFQEMIQRLYGCF 569
          RLDPYLRSLELLEKWTRLGLLMGTGAQGLREEVHTMLRGVLFSTPRTFQEMIQRLYGCF
Sbjct:  121 RLDPYLRSLELLEKWTRLGLLMGTGAQGLREEVHTMLRGVLFSTPRTFQEMIQRLYGCF 180
```

35

```
Query:  570 LRVMQSKRKGEAGTDPPELEGELDSXXXXXXXXXXLLQSPLCAGCSSDKQQCWCRALEQF 749
          LRVMQSKRKGEAGTDPPELEGELDS          LLQSPLCAGCSSDKQQCWCRALEQF
Sbjct:  181 LRVMQSKRKGEAGTDPPELEGELDSRYARRRYRLLQSPLCAGCSSDKQQCWCRALEQF 240
```

40

```
Query:  750 HQLSQVLHRLSLLERVSAEAVTTTLHQVTRERMEDRCRGEYERSFLREFHKWIERVVGWL 929
          HQLSQVLHRLSLLERVSAEAVTTTLHQVTRERMEDRCRGEYERSFLREFHKWIERVVGWL
Sbjct:  241 HQLSQVLHRLSLLERVSAEAVTTTLHQVTRERMEDRCRGEYERSFLREFHKWIERVVGWL 300
```

45

```
Query:  930 GKVFLQDGPAPASPEAGNLTNRWRCHVQRFYRIYASLRIEELFSIVRDFPDSRPAIED 1109
          GKVFLQDGPAPASPEAGNLTNRWRCHVQRFYRIYASLRIEELFSIVRDFPDSRPAIED
Sbjct:  301 GKVFLQDGPAPASPEAGNLTNRWRCHVQRFYRIYASLRIEELFSIVRDFPDSRPAIED 360
```

50

```
Query:  1110 LKYCLERTDQRQQLLVSLKAALETLLHPGVNTCDIITLYISAIKALRVLDPSMVILEVA 1289
          LKYCLERTDQRQQLLVSLKAALETLLHPGVNTCDIITLYISAIKALRVLDPSMVILEVA
Sbjct:  361 LKYCLERTDQRQQLLVSLKAALETLLHPGVNTCDIITLYISAIKALRVLDPSMVILEVA 420
```

55

```
Query:  1290 CEPIRRRLRTREDTVRQIVAXXXXXXXXXXXAVELSKTDPASLETGQDSEDDSGEPEDW 1469
          CEPIRRRLRTREDTVRQIVA          AVELSKTDPASLETGQDSEDDSGEPEDW
Sbjct:  421 CEPIRRRLRTREDTVRQIVAGLTGDSGTGDLAVELSKTDPASLETGQDSEDDSGEPEDW 480
```

```
Query:  1470 VPDVADADPGXXXXXXXXXXXXXXXXXVSIYGSKDLFINEYRSLADRLLHQFSFSPEIR 1649
          VPDVADADPG          VSIYGSKDLFINEYRSLADRLLHQFSFSPEIR
Sbjct:  481 VPDVADADPGKSSSKRRSSDIISLVSIYGSKDLFINEYRSLADRLLHQFSFSPEIR 540
```

```
Query:  1650 NVELLKLRFGEAPMHFCEVMLKDMADSRINANIREEDEKRPAAEQPPFGVYAVILSSEF 1829
          NVELLKLRFGEAPMHFCEVMLKDMADSRINANIREEDEKRPAAEQPPFGVYAVILSSEF
Sbjct:  541 NVELLKLRFGEAPMHFCEVMLKDMADSRINANIREEDEKRPAAEQPPFGVYAVILSSEF 600
```

5 Query: 1830 WPPFKDEKLEVPEDIRAALAYCKKYEQLKAMRTLSWKHTLGLVTMDVELADRTLSTAVT 2009
 WPPFKDEKLEVPEDIRAALAYCKKYEQLKAMRTLSWKHTLGLVTMDVELADRTLSTAVT
 Sbjct: 601 WPPFKDEKLEVPEDIRAALAYCKKYEQLKAMRTLSWKHTLGLVTMDVELADRTLSTAVT 660
 10 Query: 2010 PVQAVILLYFQDQASWTLEELSKAVKMPVALLRRRMSVWLQQGVLRREEPPGTFSVIEEER 2189
 PVQAVILLYFQDQASWTLEELSKAVKMPVALLRRRMSVWLQQGVLRREEPPGTFSVIEEER
 Sbjct: 661 PVQAVILLYFQDQASWTLEELSKAVKMPVALLRRRMSVWLQQGVLRREEPPGTFSVIEEER 720
 15 Query: 2190 PQDRDNMVLIDSDDSDSGMASQADQKEEELLLFWTYIQAMLTNLESLSLDRIYNMLRMF 2369
 PQDRDNMVLIDSDDSDSGMASQADQKEEELLLFWTYIQAMLTNLESLSLDRIYNMLRMF
 Sbjct: 721 PQDRDNMVLIDSDDSDSGMASQADQKEEELLLFWTYIQAMLTNLESLSLDRIYNMLRMF 780
 Query: 2370 VVTGPALAEIDLQELQGYLQKKVRDQQLVYSAGVYRLPKNCS 2495
 VVTGPALAEIDLQELQGYLQKKVRDQQLVYSAGVYRLPKNCS
 Sbjct: 781 VVTGPALAEIDLQELQGYLQKKVRDQQLVYSAGVYRLPKNCS 822

The segments of Genbank Accession AAF05751.1 that are shown as "S" above are set out in the sequence listing as SEQ ID NO: 173. Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein. Assays for determining such activities are also known in the art, some of which have been described elsewhere herein. Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out in the sequence listing as SEQ ID NO: 174 which correspond to the Q sequences in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed). Moreover, fragments and variants of these polypeptides (such as, for example, fragments as described herein, polypeptides at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to these polypeptides and polypeptides encoded by the polynucleotide which hybridizes, under stringent conditions, to the polynucleotide encoding these polypeptides) are encompassed by the invention. Antibodies that bind polypeptides of the invention are also encompassed by the invention. Polynucleotides encoding these polypeptides are also encompassed by the invention.

This protein is believed to be required for the transition from metaphase to anaphase; and is believed to act (see Yu, et. al, Science, 279:1219-122) by targeting cell-cycle regulators for destruction via the ubiquitin mediated protein degradation pathway. This latter observation is based on this protein's resemblance to cullins; which are proteins that have been implicated in similar controls of cell cycle progression during G1.

This gene is expressed in testes and epididymus tissues.

Polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include but are not limited to:

5 diseases and/or disorders of the reproductive system and proliferative disorders.

Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the testes, expression of this gene at significantly higher or lower levels may be routinely
10 detected in certain tissues or cell types (e.g., testes, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

15 The tissue distribution in testes tissue indicates that polypeptides and polynucleotides of the invention are useful for the diagnosis and/or treatment of cancer and other proliferative disorders. Additional potential uses include diagnosis and treatment of a variety of disorders of the testes which may include (but is not limited to) infertility. Polynucleotides, translation products and antibodies
20 corresponding to this gene are also useful in assays designed to identify binding agents, as such agents (antagonists) are useful as male contraceptive agents. Similarly, polynucleotides, translation products and antibodies corresponding to this gene are useful in the treatment and/or diagnosis of testicular cancer. The testes are also a site of active gene expression of transcripts that may be expressed, particularly at low
25 levels, in other tissues of the body. Therefore, translation products corresponding to this gene may be expressed in other specific tissues or organs where it may play related functional roles in other processes, such as hematopoiesis, inflammation, bone formation, and kidney function, to name a few possible target indications. Additionally, translation products corresponding to this gene, as well as antibodies
30 directed against these translation products, may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

FEATURES OF PROTEIN ENCODED BY GENE NO: 11

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. emb|CAA75729.1| (all information available through the recited accession number is incorporated herein by reference) which is described therein as "ABCR [Homo sapiens]" A partial alignment demonstrating the observed homology is shown immediately below.

```

>emb|CAA75729.1| ABCR [Homo sapiens] >sp|O60915|O60915 ABCR.
      Length = 2273

15      Plus Strand HSPs:

      Score = 312 (109.8 bits), Expect = 2.6e-28, Sum P(3) = 2.6e-28
      Identities = 58/145 (40%), Positives = 86/145 (59%), Frame = +2

20      Q:  206 MAFWTQLMLLLWKNFMYRRRQPQVQXXXXXXXXXXXXXILVAVRHSHPPLEHHECHFPNKP 385
           M F Q+ LLLWKN+ R+RQ ++                      +L+ +R+++P HHECHFPNK
      S:    1 MGFVRQIQLLLWKNWTLRKQKIRFVVELVWPLSLFLVLIWLRNANPLYSHHECHFPNKA 60

25      Q:  386 LPSAGTVPWLQGLICNVNNTCFPQLTPGEEPGRLSNFNDSLVSRLADARTVLGGASAH 565
           +PSAG +PWLQG+ CNVNN CF      TPGE PG +SN+N+S+++R+ D + +L A +
      S:    61 MPSAGMLPWLQGFICNVNNPCFQSPTPGESPGIVSNYNNSILARVYRDFQELLMNAPESQ 120

30      Q:  566 TLAGLGKLIATLRAARSTAQPQPTK 640
           L + + L      T + P +
      S:   121 HLGRIWTELHILSQFMDTLRTHPER 145

      Score = 53 (18.7 bits), Expect = 1.1e-27, Sum P(2) = 1.1e-27
      Identities = 17/50 (34%), Positives = 25/50 (50%), Frame = +2

35      Q:  815 RALLQR--PRGTSGPLELLSEALCSV-RGPSSTVGPSLNWYEASDLMELVG 958
           R L+Q P + + +LS+ LC      G S V S NWYE ++      +G
      S:   301 RPLMQNGGPETFTKLMGILSDLLCGYPEGGGSRV-LSFNWYEDNNYKAFLG 350

```

The segments of emb|CAA75729.1| that are shown as "S" above are set out in the sequence listing as SEQ ID NO: 178 and SEQ ID NO: 176 Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein. Assays for determining such activities are also known in the art, some of which have been described elsewhere herein. Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out in the sequence listing as SEQ ID NO: 175 and/or SEQ ID NO: 177 which correspond to the

Q sequences in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed). Moreover, fragments and variants of these polypeptides (such as, for example, fragments as described herein, polypeptides at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to these polypeptides and polypeptides encoded by the polynucleotide which hybridizes, under stringent conditions, to the polynucleotide encoding these polypeptides) are encompassed by the invention. Antibodies that bind polypeptides of the invention are also encompassed by the invention. Polynucleotides encoding these polypeptides are also encompassed by the invention.

This gene is expressed primarily in a variety of hematopoietic cells and tissues, including B cells, T cells, neutrophils, and dendritic cells, as well as in lymphomas such as Hodgkin's lymphoma.

Polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include but are not limited to: diseases and/or disorders of the immune system. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g., immune, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Based upon the homology to ATP-binding cassette (ABC) transporter proteins, lesions in this gene may correlate with a human disease syndrome, such as mutations in the photoreceptor cell-specific ABC transporter (ABCR) that have been recently shown to underlie Stargardt's disease. Based upon the observed preferential expression of this gene product in hematopoietic cells and tissues, polynucleotides, translation products and antibodies corresponding to this gene may play a critical role in hematopoiesis, and in the survival, proliferation, differentiation, and activation of

various blood cell lineages. Therefore, polynucleotides, translation products and antibodies corresponding to this gene may provide a useful diagnostic and/or therapeutic for a variety of hematological disorders, including inflammation, autoimmunity, allergy, neutropenia, lymphoma, susceptibility to infection, and immune system dysfunction. Additionally, translation products corresponding to this gene, as well as antibodies directed against these translation products, may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

10 FEATURES OF PROTEIN ENCODED BY GENE NO: 12

Translation products corresponding to this gene share sequence homology with a number of plant proteins of unknown function, and to a human semaphorin receptor (See Genbank Accession CAB57277).

15 This gene is expressed in the neural tissues, such as infant brain and cerebellum tissues.

Polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include but are not limited to:

20 Alzheimer's disease, Parkinson's disease, ALS, multiple sclerosis, stroke, and other disorders of the nervous system. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the nervous system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g., cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

30 Based on its expression in brain tissue, polynucleotides and translation products corresponding to this gene are useful for the treatment and/or diagnosis of

neurological conditions such as Alzheimer's disease, Parkinson's disease, ALS, multiple sclerosis and stroke as well as conditions involving problems associated with cognition, learning, memory and psychiatric disorders such as depression and psychosis. In addition, translation products corresponding to this gene may be useful
5 for the treatment of nerve injuries, such as spinal cord injuries where neuroregeneration is necessary. This is further supported by the homology of translation products corresponding to this gene to a semaphorin receptor, a class of protein which are known to be involved in the guidance and repulsion of neuronal axons during the formation of the nervous system.

10

FEATURES OF PROTEIN ENCODED BY GENE NO: 13

Translation products corresponding to this gene share sequence homology to
15 proteins of bacterial homology, such as B. Subtilis proteins (See Genbank Accession CAB16119.1). The gene encoding the disclosed cDNA is thought to reside on chromosome 17. Accordingly, polynucleotides related to this invention have uses, such as, for example, as a marker in linkage analysis for chromosome 17.

This gene is expressed in testes tissues, and to a lesser extent in colon cancer
20 tissue.

Polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include but are not limited to: testicular diseases and/or disorders, as well as colon cancer. Similarly, polypeptides
25 and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the testes, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g., testes, colon, cancerous and wounded tissues) or bodily
30 fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or sample taken from an individual having such a disorder, relative to the

standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Based on the testicular expression, polynucleotides, translation products and antibodies corresponding to this gene may be useful for the diagnosis and/or treatment of male infertility syndromes, in which sperm production is low or non existent and orchiditis. In addition, polynucleotides, translation products and antibodies corresponding to this gene may be useful for the diagnosis and/or treatment of testicular cancer. More generally, the tissue distribution in testes tissue indicates that polynucleotides, translation products and antibodies corresponding to this gene are useful for the treatment and/or diagnosis of conditions concerning proper testicular function (e.g. endocrine function, sperm maturation), as well as cancer. Therefore, polynucleotides, translation products and antibodies corresponding to this gene are useful in the treatment of male infertility and/or impotence. Polynucleotides, translation products and antibodies corresponding to this gene are also useful in assays designed to identify binding agents, as such agents (antagonists) are useful as male contraceptive agents. Similarly, polynucleotides, translation products and antibodies corresponding to this gene are useful in the treatment and/or diagnosis of testicular cancer. The testes are also a site of active gene expression of transcripts that may be expressed, particularly at low levels, in other tissues of the body. Therefore, translation products corresponding to this gene may be expressed in other specific tissues or organs where it may play related functional roles in other processes, such as hematopoiesis, inflammation, bone formation, and kidney function, to name a few possible target indications. Additionally, translation products corresponding to this gene, as well as antibodies directed against these translation products, may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

FEATURES OF PROTEIN ENCODED BY GENE NO: 14

This gene is expressed primarily in wound healing tissues, brain tissue, and Merkel cells.

Polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include but are not limited to: diseases and/or disorders of the neural system and of wound healing. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the neural system and wound healing disorders, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g., neural, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Based upon the elevated expression of this gene product in Merkel cells, it may play a role in the detection of sensations, pain, analgesia, or nerve transmission. Therefore, polynucleotides and translation products corresponding to this gene may even play a diagnostic/therapeutic role in neurodegenerative disorders such as schizophrenia, ALS, and even depression. Expression of this gene product in wound healing tissue indicates a possible involvement in normal wound healing, and may implicate this gene in fibrosis, scarring, and inflammation. Additionally, translation products corresponding to this gene, as well as antibodies directed against these translation products, may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

FEATURES OF PROTEIN ENCODED BY GENE NO: 15

This gene is expressed primarily in helper T cells, and to a lesser extent in healing groin wound and kidney tissues.

Polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include but are not limited to:

disorders of the immune system, including autoimmunity, susceptibility to infection, allergy, inflammation, and immune system dysfunction. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g., cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Based upon the elevated expression of this gene product in helper T cells, polynucleotides, translation products and antibodies corresponding to this gene may play a useful role in the diagnosis and/or treatment of disorders of the immune system, including autoimmunity, susceptibility to infection, allergy, inflammation, and immune system dysfunction. It may also be useful in anti-tumor immunotherapy. Expression of this gene product in healing wounds indicates that polynucleotides, translation products and antibodies corresponding to this gene may also be involved in such clinical conditions as fibrosis, scarring, mesengensis, scleroderma, ARDS, osteoarthritis, and other disorders involving fibroblasts and/or epithelial cells. Similarly, expression of this gene product in kidney indicates that polynucleotides, translation products and antibodies corresponding to this gene may also play a role in normal renal function, and may be an important diagnostic and/or therapeutic in the area of acute renal failure, or other kidney disorders such as kidney transplant rejection. Additionally, translation products corresponding to this gene, as well as antibodies directed against these translation products, may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

FEATURES OF PROTEIN ENCODED BY GENE NO: 16

Translation products corresponding to this gene share sequence homology to androgen/steroid hormone receptors (See Genbank Accession G01627).

This gene is expressed in various tumor tissues, such as breast, ovarian, kidney, uterus, and pancreatic tumor tissues.

5 Polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include but are not limited to: tumors of various bodily tissues. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential
10 identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly pertaining to tumorous tissues, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g., cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or sample taken from
15 an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Based upon the expression pattern, polynucleotides, translation products and antibodies corresponding to this gene may be useful for the treatment and/or diagnosis
20 of neoplasms arising from liver, spleen, breast, kidney, uterus, ovary and pancreatic tissues. Importantly, translation products corresponding to this gene share homology with a sex hormone receptor, and may therefore be most efficacious in the treatment of tumors derived from uterine, mammary, ovarian, testicular, prostate and other reproductive tract tissues. Additionally, translation products corresponding to this
25 gene, as well as antibodies directed against these translation products, may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

FEATURES OF PROTEIN ENCODED BY GENE NO: 17

30 The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no.

emb|CAA68759.1| (all information available through the recited accession number is incorporated herein by reference) which is described therein as "gp210 (AA 1-1886) [Rattus norvegicus]" A partial alignment demonstrating the observed homology is shown immediately below.

5

```
>emb|CAA68759.1| gp210 (AA 1-1886) [Rattus norvegicus] >pir|S04921|S04921
nuclear pore protein gp210 precursor - rat >sp|P11654|GP21_RAT
INTEGRAL MEMBRANE GLYCOPROTEIN GP210 PRECURSOR.
Length = 1886
```

10

Plus Strand HSPs:

Score = 3765 (1325.3 bits), Expect = 0.0, Sum P(2) = 0.0

Identities = 727/833 (87%), Positives = 768/833 (92%), Frame = +2

15

```
Q: 131 AAAXKLNIKPVLPPFTRATRVNFTLEASEGCRYWLSTRPEVASIEPLGLDEQQCSQKAVV 310
A A KLNIPKVLPPFTRATRVNFTLEASEGCRYW STRPEVASIEPLG EQQCSQKAVV
S: 23 AVAAKLNIKPVLPPFTRATRVNFTLEASEGCRYWSSTRPEVASIEPLGSSEQQCSQKAVV 82
```

20

```
Q: 311 QARLTQPARLTSIIFAEDITTGQVLRCDIAVDLIHQIVSTTRELYLEDSPLELKIQA 490
QARLTQPARLTSIIFAEDITTGQVLRCDIAVDLIHQIVSTTRELYLEDSPLELKIQA
S: 83 QARLTQPARLTSIIFAEDITTGQVLRCDIAVDLIHQIVSTTRELYLEDSPLELKIQA 142
```

25

```
Q: 491 DSEGNTFSTLAGLVFEWTIVKDSEADRFSDSHNALRILTFLESTYIPPSYISEMEKAAKQ 670
DSEGNTFSTLAGLVF+WTIVKD+EA+ FSDSHNALRILTFLESTYIPPSYISEMEKAAKQ
S: 143 DSEGNTFSTLAGLVFDWTIVKDTEANGFSDSHNALRILTFLESTYIPPSYISEMEKAAKQ 202
```

30

```
Q: 671 GDTILVSGMKTGSXKLKARIQEAVYKNVRPAXVRLILLENILLNPAYDVYLMVGTSIHYK 850
GDTILVSGMKTGS KLKARIQEAVYKNVRPA VRLILLENILLNPAYDVYL+VGTSIHYK
S: 203 GDTILVSGMKTGSXKLKARIQEAVYKNVRPAEVRLLILLENILLNPAYDVYLLVGTSIHYK 262
```

35

```
Q: 851 VQKIRQKGKITEKXMPDQYELQLQNSIPGPEGDPTRPVAVLAQDTSMTALQLGQSSSLVL 1030
VQKIRQKGKITEK MPDQYELQLQNSIP P+GDP RPVAVL QDTS VTA+Q+GQS+LVL
S: 263 VQKIRQKGKITEKXMPDQYELQLQNSIPDPQGDPPARPVAVLTQDTSRVMTAMQMGQSNLVL 322
```

40

```
Q: 1031 GHRSIQMAGASRLPNSTIYVVEPGYLGFTVHPGDRWVLETGRLYEITIEVFDKFSNKVYV 1210
GHRSIQMAGASRLPNSTIYVVE GYLGFTVHPGDRWVLETG LY +TIEVFD+ SNKVY
S: 323 GHRSIQMAGASRLPNSTIYVVEAGYLGFTVHPGDRWVLETGHLVAVTIEVFDKFSNKVYV 382
```

45

```
Q: 1211 SDNIRIETVLPAEFFEVLSSSQNGSYHRIALRKGQTIDAALTSVVDQDGGVHILQVPV 1390
SDNIRIE V PAEFFEVLSSSQNGSYH +RA++ GQT I A+LTSVVDQDGGVH+LQVPV
S: 383 SDNIRIEAVFPAEFFEVLSSSQNGSYHVRVIAQSGQTTISASLTSVVDQDGGVHVLQVPV 442
```

50

```
Q: 1391 WNQQEVEIHIPITLYPSILTFFPWQPKTGAYQYTI+AH +VATVTVKGV 1570
WNQQEV+IHIPITLYPSILTFFPWQPKTGAYQYTI+AH +VATVTVKGV
S: 443 WNQQEVDIHIPITLYPSILTFFPWQPKTGAYQYTIKAHGGSGNFTWSSSYMVATVTVKGV 502
```

55

```
Q: 1571 MTTGSDIGFSVIQAHQVQNLPHFGEMKVYVIEPHSMEFAPCQVEARVGQALEPLRISGL 1750
MTTG D G SVI+AHQVQNLPHFGEMKVYVIEP SMEFAPCQVEARVG LELPL ISGL
S: 503 MTTGGDTGLSVIRAHQVQNLPHFGEMKVYVIEPSSMEFAPCQVEARVGHTLELPLTISGL 562
```

60

```
Q: 1751 MPGGASEVVTLSDCSHFDLAVEVENQGVFQPLPGRLLPPGSEHSCGVRVKAQAQGSTTLLV 1930
MPGG+SEVVTLSDCSHFDL VEVENQGVFQPLPGRLLPPG EHCAGV+V+A+AQGSTTLLV
S: 563 MPGGSEVVTLSDCSHFDLVVEVENQGVFQPLPGRLLPPGSEHSCGVRVKAQAQGSTTLLV 622
```

65

```
Q: 1931 SYRHHGHVLSAKITIAAYLPLKAVDPSSVALVTLGSSKEMLFEGGPRPWLEPSKFFQNV 2110
SY HGHVHL AKIT+AAAYLPLKAVDPSSVA+VTLGSSKEMLFEGGPRPW+LEPSKFF+NV
S: 623 SYTHGHVHLGAKITLAAAYLPLKAVDPSSVAVVTLGSSKEMLFEGGPRPWLEPSKFFRN 682
```

```
Q: 2111 TAEDTDSIGLALFAPHSRRYQQHWILVTCQALGEQVIALSVGNKPSLTNPFFAVEPAVV 2290
T+EDT SI L+L P +SRNYQQH +LVTCQALGEQVIALSVGN+PSLTNPFFAVEP VV
S: 683 TSEDTSISLSLGLPPASRRYQQHVLVTCQALGEQVIALSVGNRPSLTNPFFAVEPTVV 742
```

```
Q: 2291 KFCVAPP SRLTLVPVYTSPLQDMSCPLLQNKQVVPVSSHRNPLLDLAAYDQEGRRFDNF 2470
K VCAPPSRLTL+PVY PQLD+SCPLLQNKQVVPVSSHRNPLLDL AYDQ+GRRFDNF
```

S: 743 KSVCAPPSRLTLPVYALPQLDLSCPLLQNKQVVPVSSHRLPLDLGAYDQQGRRFDNF 802
 Q: 2471 SSLSIQWESTRPVLASIEPELPMQLVSQDDESGQKKLHGLQAILVHEASGTTA 2629
 SSLSIQWES RP+LASIE + PMQLVSQDD +GQKKLHGLQ + VHEASGTTA
 S: 803 SSLSIQWESFRPLLASIEVDQPMQLVSQDDGNGQKKLHGLQTVSVHEASGTTA 855
 Score = 1296 (456.2 bits), Expect = 0.0, Sum P(2) = 0.0
 Identities = 251/315 (79%), Positives = 286/315 (90%), Frame = +1
 Q: 2626 SITATATGYQESHLSXRTKQPHDPLVPLSASIELILVEDVRVSPPEVTIYNHPGIQAE 2805
 +I+ATATGYQ+SHLS+ KQ DPLVP+SASIELILVEDVRVSPPEVTIYNHPG+Q EL
 S: 855 AISATATGYQQSHLSAAGVKQLRDPLVPVSASIELILVEDVRVSPPEVTIYNHPGVQVEL 914
 Q: 2806 RIREGSGYFFLNTSTADVVKVAYQEARGVAMVHPLPGSSTIMIHDLCVFPAPAKAVVY 2985
 I EGSYFFLNTST D++ VAYQ+ RGVAMVHPL PGSST+M+HDLCL FPAPAKA ++
 S: 915 HITEGSGYFFLNTSTQDIINVAYQDTRGVAMVHPLPGSSTVMVHDLCLTFPAPAKATIH 974
 Q: 2986 VSDIQELYIRVVDKVEIGKTVKAYVRVLDLHKPFLAKYFFPMDLKLRAASPIITLV 3165
 VSDIQELY+RVVDKVEIGK VKAYVRVLD +KKPFLAKYF FMDLKLRAAS IITLV LD
 S: 975 VSDIQELYVRVVDKVEIGKAVKAYVRVLDVFKKPFLLAKYFTFMDLKLRAASQIITLV 1034
 Q: 3166 EALDNYTITFLIRGVAIGQTSLSASVTNKAGQRINXPPQIEVFPFRLMPRKVTLLIGA 3345
 EALDNYT TFL+ GVAIGQTSLS+ASVT+K+GQR++S QQIEVFPFRL+PRKVTLLIGA
 S: 1035 EALDNYTATFLVHGVAIGQTSLSASVTDKSGQRVSSTAQQIEVFPFRLIPRKVTLLIGA 1094
 Q: 3346 TMQVTSEGGPQPQSNILFISISNESVALVSAAGLVQGLAIGNGTVSGLVQAVDAETGKVVI 3525
 +Q+TSEGGPQPQSNILFSI+NESVA VS+AGLV+GL +GNG+V G+VQAVDAETGKV+I
 S: 1095 MIQITSEGGPQPQSNILFSINNESVAAVSSAGLVRLMGNGSVLGVVQAVDAETGKVII 1154
 Q: 3526 ISQDLVQVEVLLRA 3570
 +SQD V+VEVL L+A
 S: 1155 VSQDHVEVEVLQQA 1169
 Score = 45 (15.8 bits), Expect = 0.0, Sum P(2) = 0.0
 Identities = 15/47 (31%), Positives = 21/47 (44%), Frame = +3
 Q: 2595 PFWFTRHQEPQHHCHWLPVPPQ---LXQNKAAA*PSGASVGLHRAH 2732
 P T H PQH H +L P L ++ A+ PSG + +H
 S: 1840 PLALTPHASPHSPH--YLASSPTAFNTLPDRKASPPSGLWSPAYASH 1886

The segments of emb|CAA68759.1| that are shown as "S" above are set out in the sequence listing as SEQ ID NO: 179, SEQ ID NO: 181 and SEQ ID NO: 183 Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein. Assays for determining such activities are also known in the art, some of which have been described elsewhere herein. Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out in the sequence listing as SEQ ID NO: 180, SEQ ID NO: 182 and/or SEQ ID NO: 184 which correspond to the Q sequences in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed). Moreover, fragments and variants of these polypeptides (such as, for example, fragments as described herein, polypeptides at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to these polypeptides and polypeptides encoded by the polynucleotide

which hybridizes, under stringent conditions, to the polynucleotide encoding these polypeptides) are encompassed by the invention. Antibodies that bind polypeptides of the invention are also encompassed by the invention. Polynucleotides encoding these polypeptides are also encompassed by the invention.

5 This gene is expressed primarily in a variety of hematopoietic cells and tissues, including neutrophils, T cells, and dendritic cells, and to a lesser extent in a variety of other tissues.

 Polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample
10 and for diagnosis of diseases and conditions which include but are not limited to: diseases and/or disorders of the immune system. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression
15 of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g., immune, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an
20 individual not having the disorder.

 Based upon the observed tissue distribution, this gene is likely to play a role in the survival, proliferation, differentiation, or activation of hematopoietic cells. Therefore, polynucleotides, translation products and antibodies corresponding to this gene may be useful in the treatment and/or diagnosis of a variety of pathologies or
25 clinical conditions, including autoimmunity, lymphomas, susceptibility to infections, inflammation, neutropenia, and a variety of immune system dysfunctions. Additionally, translation products corresponding to this gene, as well as antibodies directed against these translation products, may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

30

FEATURES OF PROTEIN ENCODED BY GENE NO: 18

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no.

5 gb|AAB47236.1| (all information available through the recited accession number is incorporated herein by reference) which is described therein as "| transporter protein [Homo sapiens]" A partial alignment demonstrating the observed homology is shown immediately below.

```

10  >gb|AAF06800.1|AF196972_5 (AF196972) JM24 protein [Homo sapiens]
    >sp|AAF06800|AAF06800 JM24 protein (fragment).
    Length = 476

    Plus Strand HSPs:

15  Score = 401 (183.7 bits), Expect = 2.3e-83, Sum P(3) = 2.3e-83
    Identities = 83/121 (68%), Positives = 83/121 (68%), Frame = +1

    Query:   343 SVKAEMLMHYSQKXXXXXXXXXXXXXXXXXXXXFPIRRALQQLLPFGKAFSWPRHXX 522
                SVKAEMLMHYSQK                                FPIRRALQQLLPFGKAFSWPRH
20  Sbjet:   319 SVKAEMLMHYSQKDPLILCVRLAVLLAVTLTVPVVLPPIRRALQQLLPFGKAFSWPRHVA 378

    Query:   523 XXXXXXXXXXXXXCVPTIRDIFGVIGSTSAPSLIFILPSIFYLRIVPSEVEPFLSWPKI 702
                CVPTIRDIFGVIGSTSAPSLIFILPSIFYLRIVPSEVEPFLSWPKI
25  Sbjet:   379 IALILLVLVNVLVICVPTIRDIFGVIGSTSAPSLIFILPSIFYLRIVPSEVEPFLSWPKI 438

    Query:   703 Q 705
                Q
30  Sbjet:   439 Q 439

    Score = 182 (83.4 bits), Expect = 2.3e-83, Sum P(3) = 2.3e-83
    Identities = 35/35 (100%), Positives = 35/35 (100%), Frame = +2

    Query:   14 RPSKRRMQAVANVSIGAMFCMYGLTATFGYLTFFYS 118
                RPSKRRMQAVANVSIGAMFCMYGLTATFGYLTFFYS
35  Sbjet:   284 RPSKRRMQAVANVSIGAMFCMYGLTATFGYLTFFYS 318

    Score = 99 (45.4 bits), Expect = 2.3e-83, Sum P(3) = 2.3e-83
    Identities = 16/17 (94%), Positives = 17/17 (100%), Frame = +1

40  Query:   976 KGCQAWIMALIPTTPW 1026
                +GCQAWIMALIPTTPW
    Sbjet:   439 QGCQAWIMALIPTTPW 455

```

45 The segment of gb|AAB47236.1| that is shown as "S" above is set out in the sequence listing as SEQ ID NO: 185 Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein. Assays for determining such activities are also known in the art, some of which have

50 been described elsewhere herein. Preferred polypeptides of the invention comprise a

polypeptide having the amino acid sequence set out in the sequence listing as SEQ ID NO: 186 which corresponds to the Q sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed). Moreover, fragments and variants of these polypeptides (such as, for example, fragments as described herein, polypeptides at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to these polypeptides and polypeptides encoded by the polynucleotide which hybridizes, under stringent conditions, to the polynucleotide encoding these polypeptides) are encompassed by the invention. Antibodies that bind polypeptides of the invention are also encompassed by the invention. Polynucleotides encoding these polypeptides are also encompassed by the invention.

This gene is expressed primarily in kidney cortex, bone, immune/hematopoietic libraries, and to a lesser extent in colon and colon cancer tissues.

Polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include but are not limited to: diseases and/or disorders of the colon and the immune system. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system and the colon, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g., colon, immune, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Translation products corresponding to this gene share sequence similarity with a novel member of a new family of transporter protein genes that resides in the lung cancer TSG region. Based upon the observed tissue distribution, this gene product may be useful as a treatment and/or diagnostic for cancer, particularly colon cancer. It may also play a role in the treatment or progression of acute renal failure, based upon its expression in kidney cortex, or may play a role in other kidney-based clinical

conditions. Expression of this gene product in normal colon also indicates a role in the normal function of colon, and may be involved in inflammatory bowel disease or diverticulitis. Additionally, translation products corresponding to this gene, as well as antibodies directed against these translation products, may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

FEATURES OF PROTEIN ENCODED BY GENE NO: 19

In specific embodiments, polypeptides of the invention comprise, or alternatively consist of, the amino acid set out in the sequence listing as SEQ ID.NO: 187. Moreover, fragments and variants of these polypeptides (such as, for example, fragments as described herein, polypeptides at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to these polypeptides and polypeptides encoded by the polynucleotide which hybridizes, under stringent conditions, to the polynucleotide encoding these polypeptides) are encompassed by the invention. Antibodies that bind polypeptides of the invention are also encompassed by the invention. Polynucleotides encoding these polypeptides are also encompassed by the invention.

This gene is expressed primarily in human endometrial stromal cells (both estrogen and non-estrogen treated), synovium (both normal and transformed), osteoblasts, pregnant uterus, colon tumor and pancreas (as NCI_CGAP_Pan1) tissues, and to a lesser extent in fetal and infant tissues, lung tissue, immune cells, and microvascular endothelial cells.

Polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include but are not limited to: proliferative disorders, as well as pulmonary, developmental, immune, and vascular disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly pertaining to proliferative disorders, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g., proliferating, cancerous and wounded tissues) or bodily fluids (e.g.,

lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

5 The expression profile of this gene (in rapidly proliferating cells) indicates that it is useful for the treatment, detection, and/or prevention of cancer, as well as pulmonary, developmental, immune, and vascular disorders. Furthermore, polynucleotides, translation products and antibodies corresponding to this gene may also be used to determine biological activity, to raise antibodies, as tissue markers, to
10 isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

 Many polynucleotide sequences, such as EST sequences, are publicly
15 available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:36 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention
20 are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1283 of SEQ ID NO:36, b is an integer of 15 to 1297, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:36, and where b is greater than or equal to a + 14.

25

FEATURES OF PROTEIN ENCODED BY GENE NO: 20

 This gene is expressed primarily in primary dendritic cells, kidney and brain
30 tissues, and many different tumor tissues, such as colon and breast tumor tissues.

 Polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample

and for diagnosis of diseases and conditions which include but are not limited to: diseases and/or disorders of the immune, renal and neural systems, as well as for tumors such as colon and breast tumors. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for
5 differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune, renal system and colon and breast tissues, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g., renal, neural, colon, breast, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine,
10 synovial fluid and spinal fluid) or another tissue or sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Based on its expression in brain and kidney, this gene and its encoded
15 polypeptides may be useful for the treatment and diagnosis of disorders of neural or urological conditions involving kidney pathogenesis. Based on its expression in many different types of tumor tissues, this gene and its encoded polypeptides may be useful in the treatment and diagnosis of cancer, especially as a tumor antigen where antibodies or other proteins are directed against this protein and used to specifically
20 kill the tumor cell. More generally, this tissue distribution in neural tissues indicates that polynucleotides, translation products and antibodies corresponding to this gene are useful for the detection and/or treatment of neurodegenerative disease states and behavioural disorders such as Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, schizophrenia, mania, dementia, paranoia,
25 obsessive compulsive disorder, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. Additionally, the tissue distribution in renal tissues indicates that polynucleotides, translation products and antibodies corresponding to this gene are useful for the treatment and/or detection of kidney diseases including renal failure,
30 nephritis, renal tubular acidosis, proteinuria, pyuria, edema, pyelonephritis, hydronephritis, nephrotic syndrome, crush syndrome, glomerulonephritis, hematuria, renal colic and kidney stones, in addition to Wilms Tumor Disease, and congenital

kidney abnormalities such as horseshoe kidney, polycystic kidney, and Falconi's syndrome. This gene and its encoded polypeptides may also be useful for the treatment and diagnosis of immunological disorders. Additionally, translation products corresponding to this gene, as well as antibodies directed against these translation products, may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

FEATURES OF PROTEIN ENCODED BY GENE NO: 21

In specific embodiments, polypeptides of the invention comprise, or alternatively consist of, the following amino acid sequence: SEQ ID NO: 188. Fragments and/or variants of these polypeptides, such as, for example, fragments and/or variants as described herein, are encompassed by the invention.

Polynucleotides encoding these polypeptides (including fragments and/or variants) are also encompassed by the invention, as are antibodies that bind these polypeptides.

This gene is expressed in human adult kidney tissue and primary dendritic cells.

Polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include but are not limited to: diseases and/or disorders of the renal and immune systems. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the renal and immune systems, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g., renal, immune, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Based on its expression in kidney tissue, polynucleotides, translation products and antibodies corresponding to this gene may be useful for the treatment and/or diagnosis of disorders of the kidney, especially nephritis associated with autoimmune disease, such as lupus nephritis. Alternatively, based on its expression in dendritic
5 cells, polynucleotides, translation products and antibodies corresponding to this gene may be useful for the treatment and/or diagnosis of immune system disorders, especially those involving defective antigen presentation such as immune deficiency syndromes, as well as autoimmune diseases in which self antigens are recognized (i.e., SLE and RA), and also in the treatment and/or diagnosis of cancer in which
10 antigen presentation is thought to be useful in immune monitoring of neoplasia. More generally, the tissue distribution in primary dendritic cells indicates that polynucleotides, translation products and antibodies corresponding to this gene also strongly indicates a role for this protein in immune function and immune surveillance. Additionally, translation products corresponding to this gene, as well as antibodies
15 directed against these translation products, may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

FEATURES OF PROTEIN ENCODED BY GENE NO: 22

20 This gene is expressed primarily in fetal heart and germ cell tumors, and to a lesser extent in osteoblasts and placenta.

Polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include but are not limited to:
25 diseases and/or disorders of the cardiovascular and musculoskeletal systems. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the cardiovascular and musculoskeletal systems, expression of this gene at significantly
30 higher or lower levels may be routinely detected in certain tissues or cell types (e.g., cardiovascular, musculoskeletal, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or

sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Based upon the observed tissue distribution, this gene product may be useful
5 as a diagnostic and/or therapeutic in a variety of clinical indications. Elevated expression of this gene product in fetal heart indicates a possible role in cardiovascular function - for example, possibly in cardiomyocyte proliferation - and indicates possible roles for polynucleotides and translation products corresponding to this gene in myocardial infarction, arrhythmias, angiogenesis, coronary disease,
10 restenosis, atherosclerosis, stroke, angina, thrombosis, and wound healing. Expression of this gene product in germ cell tumors indicates a utility as a diagnostic for such cancers, and may even indicate a role in the progression or metastasis of such cancers. Similarly, expression in osteoblasts and placenta suggest possible roles in osteoporosis and other bone mass disorders, and reproductive dysfunction,
15 respectively. Additionally, translation products corresponding to this gene, as well as antibodies directed against these translation products, may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

20 **FEATURES OF PROTEIN ENCODED BY GENE NO: 23**

This gene is expressed primary dendritic cells.

Polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample
25 and for diagnosis of diseases and conditions which include but are not limited to: immune system diseases and/or disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene
30 at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g., immune, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or

sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Based on its expression in dendritic cells, polynucleotides, translation
5 products and antibodies corresponding to this gene may be useful for the treatment and/or diagnosis of immune system disorders, especially those involving defective antigen presentation such as immune deficiency syndromes, as well as autoimmune diseases in which self antigens are recognized (i.e., SLE and RA), and also in the treatment and/or diagnosis of cancer in which antigen presentation is thought to be
10 useful in immune monitoring of neoplasia. More generally, the tissue distribution in primary dendritic cells indicates that polynucleotides, translation products and antibodies corresponding to this gene also strongly indicates a role for this protein in immune function and immune surveillance. Additionally, translation products corresponding to this gene, as well as antibodies directed against these translation
15 products, may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

FEATURES OF PROTEIN ENCODED BY GENE NO: 24

20

This gene is expressed primarily in kidney tissue, in particular rejected kidney tissue.

Polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample
25 and for diagnosis of diseases and conditions which include but are not limited to: diseases and/or disorders of the renal system. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the renal system, expression of this gene at
30 significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g., renal, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or sample

taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Based upon the specific expression of this gene product in kidney,
5 polynucleotides, translation products and antibodies corresponding to this gene may be useful in the diagnosis and/or treatment of disorders of the kidney, most notably in acute renal failure and toxemia. Elevated expression of this gene product in rejected kidney may also point towards an immune function, and possible roles in tissue rejection, tumor immunotherapy, inflammation, susceptibility to infection, and
10 immune system disorders. Additionally, translation products corresponding to this gene, as well as antibodies directed against these translation products, may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

15 FEATURES OF PROTEIN ENCODED BY GENE NO: 25

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no.
20 gb|AAC11543.1| (all information available through the recited accession number is incorporated herein by reference) which is described therein as "| (AC004523) F22329_1 [Homo sapiens]" This protein is a member of Cytochrome P450/leukotriene-B4 omega hydroxylase family of genes that are involved in metabolic degradation/detoxification of a wide variety of chemical and biological
25 compounds. The action of P450 proteins can affect a wide variety of physiological processes; for example, leukotriene B4 omega-hydroxylase helps modulate the inflammatory response by reducing the amount of circulating leukotriene B4, known to be a mediator in this process (see Kikuta, et. al, (1999) DNA Cell Biol. 18:723-730). A partial alignment demonstrating the observed homology is shown
30 immediately below.

```
>gb|AAC11543.1| (AC004523) F22329_1 [Homo sapiens] >sp|O60389|O60389 F22329_1
```

(FRAGMENT).
Length = 458

Plus Strand HSPs:

5
Score = 1634 (575.2 bits), Expect = 4.4e-249, Sum P(3) = 4.4e-249
Identities = 308/312 (98%), Positives = 308/312 (98%), Frame = +3

10
Q: 531 GEGILLSGGDKWSRHRRLTPAFHFNILKSYITIFNKSANIMLDKWQHLASEGSSRLDMF 710
GEGILLSGGDKWSRHRRLTPAFHFNILKSYITIFNKSANIMLDKWQHLASEGSS LDMF
S: 67 GEGILLSGGDKWSRHRRLTPAFHFNILKSYITIFNKSANIMLDKWQHLASEGSSCLDMF 126

15
Q: 711 EHISLMTLDSLQKCIFSFDSHCQERPSEYIATILELSALVEKRSQHILQHMDFLYYLSHD 890
EHISLMTLDSLQKCIFSFDSHCQERPSEYIATILELSALVEKRSQHILQHMDFLYYLSHD
S: 127 EHISLMTLDSLQKCIFSFDSHCQERPSEYIATILELSALVEKRSQHILQHMDFLYYLSHD 186

20
Q: 891 GRRFHRACRLVHDFDTDAVIRERRRLPTQGIDDFKDKAKSKTLD FIDVLLSKDEGKA 1070
GRRFHRACRLVHDFDTDAVIRERRRLPTQGIDDFKDKAKSKTLD FIDVLLSKDEGKA
S: 187 GRRFHRACRLVHDFDTDAVIRERRRLPTQGIDDFKDKAKSKTLD FIDVLLSKDEGKA 246

25
Q: 1071 LSDEDIRAEADTFMFGGHD TTASGLSWVLYNLARHPEYQERCQEVQELLKDRDPKEIEW 1250
LSDEDIRAEADTFMFGGHD TTASGLSWVLYNLARHPEYQERCQEVQELLKDRDPKEIEW
S: 247 LSDEDIRAEADTFMFGGHD TTASGLSWVLYNLARHPEYQERCQEVQELLKDRDPKEIEW 306

30
Q: 1251 DDLAQLPFLTMCVKESLRLHPPAPFISRCCTQDIVLPDGRVIPKGITCLIDIIGVHHNPT 1430
DDL AQLPFLTMCVKESLRLHPPAPFISRCCTQDIVLPDGRVIPKGITCLIDIIGVHHNPT
S: 307 DDLAQLPFLTMCVKESLRLHPPAPFISRCCTQDIVLPDGRVIPKGITCLIDIIGVHHNPT 366

Q: 1431 VWPDP EVLPSPF 1466
VWPDP EV PF
S: 367 VWPDP EVY-DPF 377

The segment of gb|AAC11543.1| that is shown as "S" above is set out in the sequence listing as SEQ ID NO: 189 Based on the structural similarity these

35 homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein. Assays for determining such activities are also known in the art, some of which have been described elsewhere herein. Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out in the sequence listing as SEQ ID

40 NO: 190 which corresponds to the Q sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed). Moreover, fragments and variants of these polypeptides (such as, for example, fragments as described herein, polypeptides at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to these polypeptides and polypeptides encoded by the polynucleotide

45 which hybridizes, under stringent conditions, to the polynucleotide encoding these polypeptides) are encompassed by the invention. Antibodies that bind polypeptides of the invention are also encompassed by the invention. Polynucleotides encoding these polypeptides are also encompassed by the invention.

This gene is expressed primarily in tissues that exhibit rapid proliferation such as colon and a variety of tumors. Specifically this transcript is detected in the following tissues/cDNA libraries: Colon Normal III and to a lesser extent in NCI_CGAP_Co16 (colon); Human Fetal Bone; Human Tonsils, Lib 2; Human Adult Small Intestine; Human Bone Marrow, re-excision; Liver Hepatoma; Healing groin wound, 6.5 hours post incision; Human Pancreas Tumor, Reexcision; Pancreas Islet Cell Tumor; Human Eosinophils; Colon Normal II and Soares fetal liver spleen INFLS.

Polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include but are not limited to: [insert disease condition]. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the [insert system where a related disease state is likely, e.g., immune], expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g., cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution indicates that polynucleotides, translation products and antibodies corresponding to this gene are involved in the control of cellular proliferation. Thus, polynucleotides and translation products corresponding to this gene could be used in the diagnosis and/or treatment of cancer and other proliferative disorders, especially colon cancers. The partial preference for colon indicates that the protein might also be used for the diagnosis and/or treatment of a variety of diseases of that organ, including, but not limited to Crohn's disease and colitis. Additionally, translation products corresponding to this gene, as well as antibodies directed against these translation products, may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

FEATURES OF PROTEIN ENCODED BY GENE NO: 26

This gene is expressed primarily in osteoblasts.

- 5 Polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include but are not limited to: bone mass disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential
- 10 identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the skeletal system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g., skeletal, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or sample
- 15 taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

- Based upon the specific expression of this gene product in osteoblasts, it may be useful in the diagnosis and/or treatment of bone disorders, particularly osteoporosis
- 20 and osteopetrosis. The tissue distribution in osteoblasts also indicates that polynucleotides, translation products and antibodies corresponding to this gene may play a role in the survival, proliferation, and/or growth of osteoblasts, and therefore may be useful in influencing diseases and/or disorders involving bone mass. Additionally, translation products corresponding to this gene, as well as antibodies
- 25 directed against these translation products, may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

FEATURES OF PROTEIN ENCODED BY GENE NO: 27

30

This gene is expressed primarily in kidney tissue, and to a lesser extent in lung and heart tissues.

Polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include but are not limited to: diseases and/or disorders of the renal, cardiovascular, and pulmonary systems.

5 Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the renal, cardiovascular, and pulmonary systems, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g.,
10 renal, cardiovascular, pulmonary, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

15 Based upon the observed tissue distribution, polynucleotides and translation products corresponding to this gene may play a role in a variety of clinical conditions. Expression of this gene product in kidney tissue indicates a possible role in kidney function, and may implicate polynucleotides and translation products corresponding to this gene in acute renal failure, kidney dysfunction, and septicemia. Expression of
20 this gene product in lung indicates possible involvement in pulmonary function, ARDS, inflammation, lung fibrosis, angiogenesis, and circulatory function. Elevated expression of this gene product in heart indicates possible roles in congestive heart failure, myocardial infarction, cardiovascular function, and arrhythmias. Additionally, translation products corresponding to this gene, as well as antibodies directed against
25 these translation products, may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

FEATURES OF PROTEIN ENCODED BY GENE NO: 28

30

This gene is expressed primarily in placenta and lung tissues, and to a lesser extent in a variety of epithelial and hematopoietic cell types.

Polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include but are not limited to: diseases and/or disorders of the placenta and lungs. Similarly, polypeptides and
5 antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the placenta and lungs, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g., placenta, lungs, cancerous and wounded
10 tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Based upon the tissue distribution, polynucleotides and translation products
15 corresponding to this gene may be useful in the diagnosis and/or treatment of a variety of disorders. Elevated expression within placenta may be reflective of enriched expression in endothelial cells, suggesting that polynucleotides and translation products corresponding to this gene may play a role in circulatory disorders, cardiovascular disease, and angiogenesis. Enriched expression of this gene
20 in lung indicates possible involvement in pulmonary function, inflammation, or lung fibrosis, and is again consistent with enriched expression in endothelium. Expression in hematopoietic cells indicates possible roles in the survival, proliferation, differentiation, or activation of all blood cell lineages. Additionally, translation products corresponding to this gene, as well as antibodies directed against these
25 translation products, may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Table 1

| Gene No. | cDNA Clone ID | ATCC Deposit No.:Z and Date | Vector | NT SEQ ID NO: X | Total NT Seq. | 5' NT of Clone Seq. | 3' NT of Clone Seq. | 5' NT of Start Codon | 5' NT of First AA of Signal Pep | AA SEQ ID NO: Y | First AA of Sig Pep | Last AA of Sig Pep | First AA of Secreted Portion | Last AA of ORF |
|----------|---------------|-----------------------------|------------------|-----------------|---------------|---------------------|---------------------|----------------------|---------------------------------|-----------------|---------------------|--------------------|------------------------------|----------------|
| 1 | HWEAC77 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 11 | 2670 | 1 | 2670 | 32 | 32 | 91 | 1 | 17 | 18 | 531 |
| 1 | HWEAC77 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 39 | 1312 | 1 | 1312 | 1256 | 1256 | 119 | 1 | 10 | 11 | 19 |
| 2 | HRAEH37 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 12 | 2187 | 1 | 2187 | 88 | 88 | 92 | 1 | 26 | 27 | 186 |
| 2 | HRAEH37 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 40 | 1253 | 1 | 1253 | 7 | 7 | 120 | 1 | 32 | 33 | 245 |
| 2 | HRAEH37 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 41 | 958 | 549 | 958 | | 732 | 121 | 1 | 25 | 26 | 75 |
| 3 | HDQGO29 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 13 | 2664 | 1 | 2664 | 219 | 219 | 93 | 1 | 16 | 17 | 96 |
| 3 | HDQGO29 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 42 | 2197 | 1 | 2197 | 209 | 209 | 122 | 1 | 16 | 17 | 96 |
| 3 | HDQGO29 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 43 | 403 | 1 | 403 | | 214 | 123 | 1 | 8 | 9 | 9 |
| 3 | HDQGO29 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 44 | 440 | 1 | 375 | | 438 | 124 | 1 | 1 | 2 | 44 |
| 4 | HDQHY04 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 14 | 3393 | 1 | 3334 | 211 | 211 | 94 | 1 | 33 | 34 | 77 |

| Gene No. | cDNA Clone ID | ATCC Deposit No:Z and Date | Vector | NT SEQ ID NO: X | Total NT Seq. | 5' NT of Clone Seq. | 3' NT of Clone Seq. | 5' NT of Start Codon | 5' NT of First AA of Signal Pep | AA SEQ ID NO: Y | First AA of Sig Pep | Last AA of Sig Pep | First AA of Secreted Portion | Last AA of ORF |
|----------|---------------|----------------------------|------------------|-----------------|---------------|---------------------|---------------------|----------------------|---------------------------------|-----------------|---------------------|--------------------|------------------------------|----------------|
| 4 | HDQHY04 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 45 | 3288 | 1 | 3265 | 154 | 154 | 125 | 1 | 33 | 34 | 77 |
| 4 | HDQHY04 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 46 | 1080 | 495 | 1080 | 462 | 462 | 126 | 1 | | | 26 |
| 5 | HWDHAH38 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 15 | 1604 | 1 | 1604 | 255 | 255 | 95 | 1 | 20 | 21 | 40 |
| 5 | HWDHAH38 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 47 | 796 | 1 | 796 | 319 | 319 | 127 | 1 | 20 | 21 | 40 |
| 6 | HHESQ62 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 16 | 3541 | 1 | 3541 | 29 | 29 | 96 | 1 | 25 | 26 | 58 |
| 6 | HHESQ62 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 48 | 657 | 1 | 657 | 23 | 23 | 128 | 1 | 25 | 26 | 58 |
| 6 | HHESQ62 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 49 | 1082 | 1 | 1082 | | 308 | 129 | 1 | | | 6 |
| 6 | HHESQ62 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 50 | 690 | 1 | 690 | | 323 | 130 | 1 | 23 | 24 | 34 |
| 6 | HHESQ62 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 51 | 1618 | 1 | 1618 | | 720 | 131 | 1 | 1 | 2 | 90 |
| 7 | HDPSH53 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 17 | 1687 | 1 | 1687 | 153 | 153 | 97 | 1 | 19 | 20 | 127 |
| 7 | HDPSH53 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 52 | 570 | 1 | 570 | 212 | 212 | 132 | 1 | 19 | 20 | 90 |

| Gene No. | cDNA Clone ID | ATCC Deposit No.:Z and Date | Vector | NT SEQ ID NO: X | Total NT Seq. | 5' NT of Clone Seq. | 3' NT of Clone Seq. | 5' NT of Start Codon | 5' NT of First AA of Signal Pep | AA SEQ ID NO: Y | First AA of Sig Pep | Last AA of Sig Pep | First AA of Secreted Portion | Last AA of ORF |
|----------|---------------|-----------------------------|---------------|-----------------|---------------|---------------------|---------------------|----------------------|---------------------------------|-----------------|---------------------|--------------------|------------------------------|----------------|
| 8 | HDPRN70 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 18 | 2075 | 1 | 2075 | 103 | 103 | 98 | 1 | 30 | 31 | 86 |
| 8 | HDPRN70 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 53 | 355 | 1 | 355 | 155 | 155 | 133 | 1 | 30 | 31 | 67 |
| 9 | HLWCP78 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 19 | 4699 | 1 | 4699 | 218 | 218 | 99 | 1 | | | 23 |
| 9 | HLWCP78 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 54 | 1149 | 274 | 1149 | 481 | 481 | 134 | 1 | | | 23 |
| 9 | HLWCP78 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 55 | 1767 | 1 | 1767 | | 471 | 135 | 1 | | | 7 |
| 9 | HLWCP78 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 56 | 555 | 1 | 555 | | 1 | 136 | 1 | 14 | 15 | 73 |
| 10 | HTJML75 | PTA-868 10/26/99 | pCMVSPORT 2.0 | 20 | 2762 | 1 | 2762 | 30 | 30 | 100 | 1 | 1 | 2 | 822 |
| 10 | HTJML75 | PTA-868 10/26/99 | pCMVSPORT 2.0 | 57 | 2694 | 21 | 2694 | | 335 | 137 | 1 | 21 | 22 | 64 |
| 11 | HWBFES7 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 21 | 1133 | 36 | 1133 | 227 | 227 | 101 | 1 | 37 | 38 | 302 |
| 11 | HWBFES7 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 58 | 5811 | 3302 | 5811 | | 3300 | 138 | 1 | 16 | 17 | 37 |
| 11 | HWBFES7 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 59 | 1012 | 1 | 1012 | | 622 | 139 | 1 | | | 16 |

| Gene No. | cDNA Clone ID | ATCC Deposit No:Z and Date | Vector | NT SEQ ID NO: X | Total NT Seq. | 5' NT of Clone Seq. | 3' NT of Clone Seq. | 5' NT of Start Codon | 5' NT of First AA of Signal Pep | AA SEQ ID NO: Y | First AA of Sig Pep | Last AA of Sig Pep | First AA of Secreted Portion | Last AA of ORF |
|----------|---------------|----------------------------|---------------|-----------------|---------------|---------------------|---------------------|----------------------|---------------------------------|-----------------|---------------------|--------------------|------------------------------|----------------|
| 12 | HDPLO25 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 22 | 1714 | 1 | 1714 | 153 | 153 | 102 | 1 | | | 39 |
| 12 | HDPLO25 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 60 | 1680 | 26 | 1680 | 179 | 179 | 140 | 1 | 43 | 44 | 66 |
| 12 | HDPLO25 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 61 | 1386 | 225 | 972 | | 181 | 141 | 1 | 1 | 2 | 124 |
| 13 | HDPWN93 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 23 | 2679 | 1 | 2669 | 45 | 45 | 103 | 1 | 19 | 20 | 802 |
| 13 | HDPWN93 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 62 | 716 | 1 | 716 | 35 | 35 | 142 | 1 | 19 | 20 | 214 |
| 13 | HDPWN93 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 63 | 2716 | 26 | 2716 | 27 | 27 | 143 | 1 | 19 | 20 | 43 |
| 14 | HWHPM16 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 24 | 1786 | 1 | 1786 | 283 | 283 | 104 | 1 | 24 | 25 | 89 |
| 14 | HWHPM16 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 64 | 1011 | 1 | 1011 | | 878 | 144 | 1 | 1 | 2 | 99 |
| 15 | HRAEE45 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 25 | 3431 | 1 | 3431 | 49 | 49 | 105 | 1 | 30 | 31 | 79 |
| 15 | HRAEE45 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 65 | 940 | 1 | 940 | 43 | 43 | 145 | 1 | 30 | 31 | 60 |
| 15 | HRAEE45 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 66 | 490 | 238 | 359 | | 489 | 146 | 1 | 1 | 2 | 101 |

| Gene No. | cDNA Clone ID | ATCC Deposit No.:Z and Date | Vector | NT SEQ ID NO: X | Total NT Seq. | 5' NT of Clone Seq. | 3' NT of Clone Seq. | 5' NT of Start Codon | 5' NT of First AA of Signal Pep | AA SEQ ID NO: Y | First AA of Sig Pep | Last AA of Sig Pep | First AA of Secreted Portion | Last AA of ORF |
|----------|---------------|-----------------------------|---------------|-----------------|---------------|---------------------|---------------------|----------------------|---------------------------------|-----------------|---------------------|--------------------|------------------------------|----------------|
| 16 | HDPXY01 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 26 | 766 | 1 | 766 | 23 | 23 | 106 | 1 | 37 | 38 | 98 |
| 16 | HDPXY01 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 67 | 2409 | 1 | 2409 | 33 | 33 | 147 | 1 | 37 | 38 | 98 |
| 16 | HDPXY01 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 68 | 737 | 1 | 423 | | 539 | 148 | 1 | | | 22 |
| 16 | HDPXY01 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 69 | 1471 | 105 | 1471 | | 1190 | 149 | 1 | | | 25 |
| 17 | HWBEM18 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 27 | 6729 | 1 | 6729 | 75 | 75 | 107 | 1 | 25 | 26 | 1887 |
| 17 | HWBEM18 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 70 | 3599 | 1 | 3599 | 65 | 65 | 150 | 1 | 25 | 26 | 886 |
| 17 | HWBEM18 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 71 | 2924 | 429 | 2924 | | 2924 | 151 | 1 | 1 | 2 | 498 |
| 18 | HDTFE17 | PTA-868 10/26/99 | pCMVSPORT 2.0 | 28 | 1242 | 1 | 1242 | 260 | 260 | 108 | 1 | 20 | 21 | 29 |
| 18 | HDTFE17 | PTA-868 10/26/99 | pCMVSPORT 2.0 | 72 | 628 | 1 | 628 | 251 | 251 | 152 | 1 | 20 | 21 | 29 |
| 18 | HDTFE17 | PTA-868 10/26/99 | pCMVSPORT 2.0 | 73 | 923 | 29 | 903 | | 101 | 153 | 1 | 6 | 7 | 80 |
| 19 | HBXAB02 | PTA-868 10/26/99 | ZAP Express | 29 | 2032 | 1 | 1994 | | 2 | 109 | 1 | 1 | 2 | 378 |

| Gene No. | cDNA Clone ID | ATCC Deposit No.:Z and Date | Vector | NT SEQ ID NO: X | Total NT Seq. | 5' NT of Clone Seq. | 3' NT of Clone Seq. | 5' NT of Start Codon | 5' NT of First AA of Signal Pep | AA SEQ ID NO: Y | First AA of Sig Pep | Last AA of Sig Pep | First AA of Secreted Portion | Last AA of ORF |
|----------|---------------|-----------------------------|---------------|-----------------|---------------|---------------------|---------------------|----------------------|---------------------------------|-----------------|---------------------|--------------------|------------------------------|----------------|
| 19 | HBXAB02 | PTA-868 10/26/99 | ZAP Express | 74 | 1297 | 1 | 1297 | 131 | 131 | 154 | 1 | | | 11 |
| 19 | HBXAB02 | PTA-868 10/26/99 | ZAP Express | 75 | 2550 | 688 | 2518 | 818 | 818 | 155 | 1 | | | 11 |
| 20 | HDPSB18 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 30 | 3408 | 1 | 3408 | 123 | 123 | 110 | 1 | 18 | 19 | 66 |
| 20 | HDPSB18 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 76 | 308 | 1 | 308 | | 116 | 156 | 1 | 17 | 18 | 64 |
| 20 | HDPSB18 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 77 | 1568 | 1 | 1568 | | 1525 | 157 | 1 | 9 | 10 | 14 |
| 20 | HDPSB18 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 78 | 865 | 1 | 865 | | 521 | 158 | 1 | 1 | 2 | 107 |
| 21 | HDPTW65 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 31 | 1917 | 1 | 1917 | 236 | 236 | 111 | 1 | 15 | 16 | 48 |
| 21 | HDPTW65 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 79 | 1182 | 180 | 1182 | 407 | 407 | 159 | 1 | 15 | 16 | 48 |
| 21 | HDPTW65 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 80 | 1074 | 1 | 1074 | | 404 | 160 | 1 | 1 | 2 | 73 |
| 22 | HOHDF66 | PTA-868 10/26/99 | pCMVSPORT 2.0 | 32 | 2359 | 1 | 2359 | 25 | 25 | 112 | 1 | 21 | 22 | 778 |
| 22 | HOHDF66 | PTA-868 10/26/99 | pCMVSPORT 2.0 | 81 | 546 | 47 | 546 | 63 | 63 | 161 | 1 | 21 | 22 | 161 |

| Gene No. | cDNA Clone ID | ATCC Deposit No:Z and Date | Vector | NT SEQ ID NO: X | Total NT Seq. | 5' NT of Clone Seq. | 3' NT of Clone Seq. | 5' NT of Start Codon | 5' NT of First AA of Signal Pep | AA SEQ ID NO: Y | First AA of Sig Pep | Last AA of Sig Pep | First AA of Secreted Portion | Last AA of ORF |
|----------|---------------|----------------------------|---------------|-----------------|---------------|---------------------|---------------------|----------------------|---------------------------------|-----------------|---------------------|--------------------|------------------------------|----------------|
| 22 | HOHDF66 | PTA-868 10/26/99 | pCMVSPORT 2.0 | 82 | 557 | 1 | 557 | | 557 | 162 | 1 | 1 | 2 | 181 |
| 23 | HDPTW24 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 33 | 527 | 1 | 527 | 100 | 100 | 113 | 1 | 22 | 23 | 55 |
| 24 | HRADO01 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 34 | 1951 | 1 | 1951 | 284 | 284 | 114 | 1 | 18 | 19 | 33 |
| 24 | HRADO01 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 83 | 932 | 1 | 932 | 278 | 278 | 163 | 1 | 18 | 19 | 33 |
| 24 | HRADO01 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 84 | 661 | 1 | 661 | | 398 | 164 | 1 | 1 | 2 | 85 |
| 25 | HTJNX29 | PTA-868 10/26/99 | pCMVSPORT 2.0 | 35 | 2024 | 1 | 2024 | 86 | 86 | 115 | 1 | 25 | 26 | 157 |
| 25 | HTJNX29 | PTA-868 10/26/99 | pCMVSPORT 2.0 | 85 | 2224 | 30 | 2224 | 108 | 108 | 165 | 1 | 25 | 26 | 126 |
| 25 | HTJNX29 | PTA-868 10/26/99 | pCMVSPORT 2.0 | 86 | 1669 | 21 | 1669 | | 1629 | 166 | 1 | | | 13 |
| 26 | HOHDC86 | PTA-868 10/26/99 | pCMVSPORT 2.0 | 36 | 3710 | 1 | 3710 | 449 | 449 | 116 | 1 | 21 | 22 | 102 |
| 26 | HOHDC86 | PTA-868 10/26/99 | pCMVSPORT 2.0 | 87 | 1158 | 1 | 1158 | 309 | 309 | 167 | 1 | 21 | 22 | 33 |
| 26 | HOHDC86 | PTA-868 10/26/99 | pCMVSPORT 2.0 | 88 | 963 | 1 | 963 | | 962 | 168 | 1 | 1 | 2 | 132 |

| Gene No. | cDNA Clone ID | ATCC Deposit No:Z and Date | Vector | NT SEQ ID NO: X | Total NT Seq. | 5' NT of Clone Seq. | 3' NT of Clone Seq. | 5' NT of Start Codon | 5' NT of First AA of Signal Pep | AA SEQ ID NO: Y | First AA of Sig Pep | Last AA of Sig Pep | First AA of Secreted Portion | Last AA of ORF |
|----------|---------------|----------------------------|------------------|-----------------|---------------|---------------------|---------------------|----------------------|---------------------------------|-----------------|---------------------|--------------------|------------------------------|----------------|
| 27 | HLDQA07 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 37 | 343 | 1 | 343 | 127 | 127 | 117 | 1 | 29 | 30 | 72 |
| 27 | HLDQA07 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 89 | 579 | 241 | 579 | 359 | 359 | 169 | 1 | 29 | 30 | 73 |
| 28 | HWDAD17 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 38 | 445 | 1 | 443 | 99 | 99 | 118 | 1 | 26 | 27 | 46 |
| 28 | HWDAD17 | PTA-868 10/26/99 | pCMVSPORT 3.0 | 90 | 1316 | 890 | 1316 | 974 | 974 | 170 | 1 | 26 | 27 | 53 |

Table 1 summarizes the information corresponding to each "Gene No." described above. The nucleotide sequence identified as "NT SEQ ID NO:X" was assembled from partially homologous ("overlapping") sequences obtained from the "cDNA clone ID" identified in Table 1 and, in some cases, from additional related DNA clones. The overlapping sequences were assembled into a single contiguous sequence of high redundancy (usually three to five overlapping sequences at each nucleotide position), resulting in a final sequence identified as SEQ ID NO:X.

The cDNA Clone ID was deposited on the date and given the corresponding deposit number listed in "ATCC Deposit No:Z and Date." Some of the deposits contain multiple different clones corresponding to the same gene. "Vector" refers to the type of vector contained in the cDNA Clone ID.

"Total NT Seq." refers to the total number of nucleotides in the contig identified by "Gene No." The deposited clone may contain all or most of these sequences, reflected by the nucleotide position indicated as "5' NT of Clone Seq." and the "3' NT of Clone Seq." of SEQ ID NO:X. The nucleotide position of SEQ ID NO:X of the putative start codon (methionine) is identified as "5' NT of Start Codon." Similarly, the nucleotide position of SEQ ID NO:X of the predicted signal sequence is identified as "5' NT of First AA of Signal Pep."

The translated amino acid sequence, beginning with the methionine, is identified as "AA SEQ ID NO:Y," although other reading frames can also be easily translated using known molecular biology techniques. The polypeptides produced by these alternative open reading frames are specifically contemplated by the present invention.

The first and last amino acid position of SEQ ID NO:Y of the predicted signal peptide is identified as "First AA of Sig Pep" and "Last AA of Sig Pep." The predicted first amino acid position of SEQ ID NO:Y of the secreted portion is identified as "Predicted First AA of Secreted Portion." Finally, the amino acid position of SEQ ID NO:Y of the last amino acid in the open reading frame is identified as "Last AA of ORF."

SEQ ID NO:X (where X may be any of the polynucleotide sequences disclosed in the sequence listing) and the translated SEQ ID NO:Y (where Y may be any of the polypeptide sequences disclosed in the sequence listing) are sufficiently

accurate and otherwise suitable for a variety of uses well known in the art and described further below. For instance, SEQ ID NO:X is useful for designing nucleic acid hybridization probes that will detect nucleic acid sequences contained in SEQ ID NO:X or the cDNA contained in the deposited clone. These probes will also

5 hybridize to nucleic acid molecules in biological samples, thereby enabling a variety of forensic and diagnostic methods of the invention. Similarly, polypeptides identified from SEQ ID NO:Y may be used, for example, to generate antibodies which bind specifically to proteins containing the polypeptides and the secreted proteins encoded by the cDNA clones identified in Table 1.

10 Nevertheless, DNA sequences generated by sequencing reactions can contain sequencing errors. The errors exist as misidentified nucleotides, or as insertions or deletions of nucleotides in the generated DNA sequence. The erroneously inserted or deleted nucleotides cause frame shifts in the reading frames of the predicted amino acid sequence. In these cases, the predicted amino acid sequence diverges from the

15 actual amino acid sequence, even though the generated DNA sequence may be greater than 99.9% identical to the actual DNA sequence (for example, one base insertion or deletion in an open reading frame of over 1000 bases).

Accordingly, for those applications requiring precision in the nucleotide sequence or the amino acid sequence, the present invention provides not only the

20 generated nucleotide sequence identified as SEQ ID NO:X and the predicted translated amino acid sequence identified as SEQ ID NO:Y, but also a sample of plasmid DNA containing a human cDNA of the invention deposited with the ATCC, as set forth in Table 1. The nucleotide sequence of each deposited clone can readily be determined by sequencing the deposited clone in accordance with known methods.

25 The predicted amino acid sequence can then be verified from such deposits. Moreover, the amino acid sequence of the protein encoded by a particular clone can also be directly determined by peptide sequencing or by expressing the protein in a suitable host cell containing the deposited human cDNA, collecting the protein, and determining its sequence.

30 The present invention also relates to the genes corresponding to SEQ ID NO:X, SEQ ID NO:Y, or the deposited clone. The corresponding gene can be isolated in accordance with known methods using the sequence information disclosed

herein. Such methods include preparing probes or primers from the disclosed sequence and identifying or amplifying the corresponding gene from appropriate sources of genomic material.

Also provided in the present invention are allelic variants, orthologs, and/or species homologs. Procedures known in the art can be used to obtain full-length genes, allelic variants, splice variants, full-length coding portions, orthologs, and/or species homologs of genes corresponding to SEQ ID NO:X, SEQ ID NO:Y, or a deposited clone, using information from the sequences disclosed herein or the clones deposited with the ATCC. For example, allelic variants and/or species homologs may be isolated and identified by making suitable probes or primers from the sequences provided herein and screening a suitable nucleic acid source for allelic variants and/or the desired homologue.

Table 2 provides preferred epitopes contained in certain embodiments of the invention and polynucleotide sequences that may be disclaimed according to certain embodiments of the invention. The first column refers to each "Gene No." described above in Table 1. The second column provides the sequence identifier, "NT SEQ ID NO:X", for polynucleotide sequences disclosed in Table 1. The third column provides the sequence identifier, "AA SEQ ID NO:Y", for polypeptide sequences disclosed in Table 1. The fourth column provides a unique integer "ntA" where "ntA" is any integer between 1 and the final nucleotide minus 15 of SEQ ID NO:X, and the fifth column provides a unique integer "ntB" where "ntB" is any integer between 15 and the final nucleotide of SEQ ID NO:X, where both ntA and ntB correspond to the positions of nucleotide residues shown in SEQ ID NO:X, and where ntB is greater than or equal to a + 14. For each of the polynucleotides shown as SEQ ID NO:X, the uniquely defined integers can be substituted into the general formula of a-b, and used to describe polynucleotides which may be preferably excluded from the invention. Column 6 lists residues comprising predicted epitopes contained in the polypeptides encoded by each of the preferred ORFs (SEQ ID NO:Y). Identification of potential immunogenic regions was performed according to the method of Jameson and Wolf ((1988) CABIOS, 4; 181-186); specifically, the Genetics Computer Group (GCG) implementation of this algorithm, embodied in the program PEPTIDESTRUCTURE

(Wisconsin Package v10.0, Genetics Computer Group (GCG), Madison, Wisc.). This method returns a measure of the probability that a given residue is found on the surface of the protein. Regions where the antigenic index score is greater than 0.9 over at least 6 amino acids are indicated in Table 2 as "Preferred Epitopes".

- 5 Polypeptides of the invention may possess one, two, three, four, five or more antigenic epitopes comprising residues described in Table 2. It will be appreciated that depending on the analytical criteria used to predict antigenic determinants, the exact address of the determinant may vary slightly.

- 10 Table 3 summarizes the expression profile of polynucleotides corresponding to the clones disclosed in Table 1. The first column provides a unique clone identifier, "Clone ID", for a cDNA clone related to each contig sequence disclosed in Table 1. Column 2, "Library Codes" shows the expression profile of tissue and/or cell line libraries which express the polynucleotides of the invention. Each Library Code
15 in column 2 represents a tissue/cell source identifier code corresponding to the Library Code and Library description provided in Table 5. Expression of these polynucleotides was not observed in the other tissues and/or cell libraries tested. One of skill in the art could routinely use this information to identify tissues which show a predominant expression pattern of the corresponding polynucleotide of the invention
20 or to identify polynucleotides which show predominant and/or specific tissue expression.

- Table 4, column 1, provides a nucleotide sequence identifier, "SEQ ID NO:X," that matches a nucleotide SEQ ID NO:X disclosed in Table 1, column 5. Table 4, column 2, provides the chromosomal location, "Cytologic Band or
25 Chromosome," of polynucleotides corresponding to SEQ ID NO:X. Chromosomal location was determined by finding exact matches to EST and cDNA sequences contained in the NCBI (National Center for Biotechnology Information) UniGene database. Given a presumptive chromosomal location, disease locus association was determined by comparison with the Morbid Map, derived from Online Mendelian
30 Inheritance in Man (Online Mendelian Inheritance in Man, OMIM™. McKusick-Nathans Institute for Genetic Medicine, Johns Hopkins University (Baltimore, MD) and National Center for Biotechnology Information, National Library of Medicine

(Bethesda, MD) 2000. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>).

If the putative chromosomal location of the Query overlapped with the chromosomal location of a Morbid Map entry, the OMIM reference identification number of the morbid map entry is provided in Table 4, column 3, labelled "OMIM ID." A key to the OMIM reference identification numbers is provided in Table 6.

Table 5 provides a key to the Library Code disclosed in Table 3. Column 1 provides the Library Code disclosed in Table 3, column 2. Column 2 provides a description of the tissue or cell source from which the corresponding library was derived. Library codes corresponding to diseased Tissues are indicated in column 3 with the word "disease".

Table 6 provides a key to the OMIM reference identification numbers disclosed in Table 4, column 3. OMIM reference identification numbers (Column 1) were derived from Online Mendelian Inheritance in Man (Online Mendelian Inheritance in Man, OMIM. McKusick-Nathans Institute for Genetic Medicine, Johns Hopkins University (Baltimore, MD) and National Center for Biotechnology Information, National Library of Medicine, (Bethesda, MD) 2000. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>). Column 2 provides diseases associated with the cytologic band disclosed in Table 4, column 2, as determined using the Morbid Map database.

Table 2

| Gene # | NT SEQ ID NO: X | AA SEQ ID NO: Y | nt A | nt B | Preferred Epitopes |
|--------|-----------------|-----------------|----------|-----------|--|
| 1 | 11 | 91 | 1 - 2656 | 15 - 2670 | Arg-16 to Gly-22 Leu-98 to Gly-106 Pro-152 to Thr-158 Pro-175 to Glu-180 Pro-280 to Met-286 Ala-389 to Phe-395 His-408 to Arg-417 Ala-505 to Thr-514 Gly-518 to Ala-530. |
| 1 | 39 | 119 | 1 - 1298 | 15 - 1312 | |
| 2 | 12 | 92 | 1 - 2173 | 15 - 2187 | Asp-111 to Gln-116 Glu-154 to Val-160 Ala-162 to Cys-178. |
| 2 | 40 | 120 | 1 - 1239 | 15 - 1253 | Ala-6 to Gln-11 Asp-136 to Gln-141 Glu-179 to Val-185 Ala-187 to Glu-192 Ala-218 to Lys-224. |
| 2 | 41 | 121 | 1 - 944 | 15 - 958 | Leu-28 to Ala-35 Phe-39 to Pro-46. |
| 3 | 13 | 93 | 1 - 2650 | 15 - 2664 | Ala-28 to Ser-34 Gln-54 to Asp-62 Pro-71 to Arg-89. |
| 3 | 42 | 122 | 1 - 2183 | 15 - 2197 | Ala-28 to Ser-34 Gln-54 to Asp-62 Pro-71 to Arg-89. |
| 3 | 43 | 123 | 1 - 389 | 15 - 403 | |
| 3 | 44 | 124 | 1 - 426 | 15 - 440 | |
| 4 | 14 | 94 | 1 - 3379 | 15 - 3393 | |
| 4 | 45 | 125 | 1 - 3274 | 15 - 3288 | |
| 4 | 46 | 126 | 1 - 1066 | 15 - 1080 | |
| 5 | 15 | 95 | 1 - 1590 | 15 - 1604 | |
| 5 | 47 | 127 | 1 - 782 | 15 - 796 | |
| 6 | 16 | 96 | 1 - 3527 | 15 - 3541 | |
| 6 | 48 | 128 | 1 - 643 | 15 - 657 | |
| 6 | 49 | 129 | 1 - 1068 | 15 - 1082 | |
| 6 | 50 | 130 | 1 - 676 | 15 - 690 | |
| 6 | 51 | 131 | 1 - 1604 | 15 - 1618 | |
| 7 | 17 | 97 | 1 - 1673 | 15 - 1687 | Met-1 to Trp-6 Leu-22 to Thr-27 Pro-44 to Gly-58 Ala-61 to Glu-74 Pro-99 to Gly-111 Cys-121 to Ser-127. |
| 7 | 52 | 132 | 1 - 556 | 15 - 570 | Met-1 to Trp-6 Leu-22 to Thr-27. |
| 8 | 18 | 98 | 1 - 2061 | 15 - 2075 | |
| 8 | 53 | 133 | 1 - 341 | 15 - 355 | |
| 9 | 19 | 99 | 1 - 4685 | 15 - 4699 | |

| | | | | | |
|----|----|-----|----------|-----------|--|
| 9 | 54 | 134 | 1 - 1135 | 15 - 1149 | |
| 9 | 55 | 135 | 1 - 1753 | 15 - 1767 | |
| 9 | 56 | 136 | 1 - 541 | 15 - 555 | Leu-9 to Tyr-14 Arg-36 to Arg-48. |
| 10 | 20 | 100 | 1 - 2748 | 15 - 2762 | Gly-10 to Gly-17 Pro-49 to Glu-54 Gln-97 to Asp-103 Ser-120 to Tyr-125 Gln-186 to Leu-199 Glu-202 to Tyr-213 Ser-225 to Cys-233 Thr-269 to Ser-284 Gly-308 to Val-328 Asp-350 to Ala-357 Arg-367 to Gln-372 Arg-429 to Thr-434 Gly-444 to Thr-449 Thr-466 to Val-481 Val-485 to Ser-499 Ser-534 to Arg-540 Met-564 to Ile-570 Asn-573 to Phe-589 Pro-603 to Val-611 Arg-706 to Gly-711 Glu-717 to Asp-725 Ser-732 to Ser-738 Gln-743 to Glu-749 Leu-799 to Asp-805. |
| 10 | 57 | 137 | 1 - 2680 | 15 - 2694 | Gly-49 to His-56. |
| 11 | 21 | 101 | 1 - 1119 | 15 - 1133 | Phe-8 to Pro-15 Glu-43 to Leu-54 Thr-79 to Arg-86 Arg-129 to Leu-142 Gln-201 to Gly-208 Ser-249 to Gln-254. |
| 11 | 58 | 138 | 1 - 5797 | 15 - 5811 | |
| 11 | 59 | 139 | 1 - 998 | 15 - 1012 | |
| 12 | 22 | 102 | 1 - 1700 | 15 - 1714 | |
| 12 | 60 | 140 | 1 - 1666 | 15 - 1680 | |
| 12 | 61 | 141 | 1 - 1372 | 15 - 1386 | Ser-8 to Leu-13. |
| 13 | 23 | 103 | 1 - 2665 | 15 - 2679 | Pro-36 to Ser-52 Ala-63 to Pro-78 Ala-106 to Lys-115 Glu-134 to Glu-141 Val-155 to Asp-164 Phe-199 to Gly-204 Arg-218 to Leu-228 Glu-230 to Val-235 Val-247 to Pro-253 Arg-262 to Gly-276 Thr-303 to Gln-310 Arg-335 to Trp-342 Glu-399 to Ala-415 Ser-458 to Glu-466 Arg-508 to Asp-517 Glu-580 to Pro-585 |

| | | | | | |
|----|----|-----|----------|-----------|---|
| | | | | | Gln-620 to Trp-628 Lys-651 to Ala-657 Gly-677 to Met-682 Ala-712 to Leu-717 Gly-724 to Thr-731 Arg-770 to Gln-775. |
| 13 | 62 | 142 | 1 - 702 | 15 - 716 | Pro-36 to Ser-52 Ala-63 to Pro-78 Ala-106 to Lys-115 Glu-134 to Glu-141 Val-155 to Asp-164. |
| 13 | 63 | 143 | 1 - 2702 | 15 - 2716 | |
| 14 | 24 | 104 | 1 - 1772 | 15 - 1786 | Ser-31 to Asp-40 Arg-80 to Gln-88. |
| 14 | 64 | 144 | 1 - 997 | 15 - 1011 | Ser-1 to Gly-12 Gln-88 to Ser-95. |
| 15 | 25 | 105 | 1 - 3417 | 15 - 3431 | Arg-26 to Asn-34. |
| 15 | 65 | 145 | 1 - 926 | 15 - 940 | Arg-26 to Asn-34. |
| 15 | 66 | 146 | 1 - 476 | 15 - 490 | Pro-3 to Arg-12. |
| 16 | 26 | 106 | 1 - 752 | 15 - 766 | Pro-39 to Trp-44. |
| 16 | 67 | 147 | 1 - 2395 | 15 - 2409 | Pro-39 to Trp-44. |
| 16 | 68 | 148 | 1 - 723 | 15 - 737 | |
| 16 | 69 | 149 | 1 - 1457 | 15 - 1471 | |
| 17 | 27 | 107 | 1 - 6715 | 15 - 6729 | Ser-165 to His-174 Met-196 to Asp-204 Lys-212 to Leu-218 Pro-277 to Leu-285 Pro-290 to Arg-298 Ser-402 to Ser-407 Trp-465 to Gly-470 His-698 to His-706 Asp-793 to Asn-801 Gln-830 to Lys-838 Gly-862 to His-867 Ala-871 to His-877 Lys-1063 to Asn-1069 Ser-1100 to Ser-1108 Asn-1194 to Ser-1200 Leu-1308 to Gly-1314 Lys-1437 to Asn-1442 Asp-1583 to Val-1599 Thr-1651 to Lys-1656 Lys-1735 to Gly-1740 Arg-1789 to Tyr-1795 Arg-1846 to His-1854 Pro-1869 to Pro-1875. |
| 17 | 70 | 150 | 1 - 3585 | 15 - 3599 | |
| 17 | 71 | 151 | 1 - 2910 | 15 - 2924 | |
| 18 | 28 | 108 | 1 - 1228 | 15 - 1242 | |
| 18 | 72 | 152 | 1 - 614 | 15 - 628 | |
| 18 | 73 | 153 | 1 - 909 | 15 - 923 | |
| 19 | 29 | 109 | 1 - 2018 | 15 - 2032 | Ala-1 to Tyr-6 Cys-65 to Tyr-70 Asp-111 to Gly-118 Pro-126 to Thr-132 Tyr-145 to Lys-152 |

| | | | | | |
|----|----|-----|----------|-----------|--|
| | | | | | Gly-181 to His-198 Glu-262 to Phe-268 Ser-297 to Gln-302 Ala-344 to Phe-350 Ala-360 to Cys-378. |
| 19 | 74 | 154 | 1 - 1283 | 15 - 1297 | |
| 19 | 75 | 155 | 1 - 2536 | 15 - 2550 | |
| 20 | 30 | 110 | 1 - 3394 | 15 - 3408 | Lys-23 to Lys-31 Ala-38 to Ser-43. |
| 20 | 76 | 156 | 1 - 294 | 15 - 308 | |
| 20 | 77 | 157 | 1 - 1554 | 15 - 1568 | |
| 20 | 78 | 158 | 1 - 851 | 15 - 865 | Lys-57 to Gly-64. |
| 21 | 31 | 111 | 1 - 1903 | 15 - 1917 | |
| 21 | 79 | 159 | 1 - 1168 | 15 - 1182 | |
| 21 | 80 | 160 | 1 - 1060 | 15 - 1074 | Leu-8 to Tyr-13 Ser-18 to His-31 Ser-67 to Ser-73. |
| 22 | 32 | 112 | 1 - 2345 | 15 - 2359 | Gln-22 to Asn-31 Asn-66 to Gln-71 Ala-99 to Pro-117 Arg-161 to Lys-168 Ile-222 to His-230 His-255 to Pro-266 Glu-314 to Val-320 Thr-334 to Thr-343 Lys-358 to Thr-363 Lys-416 to Leu-424 Pro-437 to Ser-453 Pro-461 to Arg-472 Pro-493 to Thr-506 Ser-510 to Thr-529 Ile-538 to Lys-553 Val-569 to Gln-596 Val-607 to Val-619 His-633 to Glu-648 Ser-661 to Leu-667 Thr-670 to Phe-682 Thr-696 to Phe-702 Val-706 to His-715 Arg-722 to Gly-734 Gln-747 to Arg-756 Asp-761 to Pro-777. |
| 22 | 81 | 161 | 1 - 532 | 15 - 546 | Gln-22 to Asn-31 Asn-66 to Gln-71 Ala-99 to Pro-117. |
| 22 | 82 | 162 | 1 - 543 | 15 - 557 | Pro-1 to Val-9 His-23 to Glu-38 Ser-51 to Leu-57 Thr-60 to Phe-72. |
| 23 | 33 | 113 | 1 - 513 | 15 - 527 | |
| 24 | 34 | 114 | 1 - 1937 | 15 - 1951 | |
| 24 | 83 | 163 | 1 - 918 | 15 - 932 | |
| 24 | 84 | 164 | 1 - 647 | 15 - 661 | Ile-53 to Glu-67. |
| 25 | 35 | 115 | 1 - 2010 | 15 - 2024 | Gln-38 to Phe-45 Thr-55 to Gln-64 Pro-147 to Arg-157. |

| | | | | | |
|----|----|-----|----------|-----------|--|
| 25 | 85 | 165 | 1 - 2210 | 15 - 2224 | Gln-38 to Phe-45 Thr-55 to Gln-64. |
| 25 | 86 | 166 | 1 - 1655 | 15 - 1669 | Trp-6 to Arg-13. |
| 26 | 36 | 116 | 1 - 3696 | 15 - 3710 | His-29 to Ser-39 Lys-44 to Thr-56 Gly-87 to Gly-93 Leu-96 to Asp-102. |
| 26 | 87 | 167 | 1 - 1144 | 15 - 1158 | |
| 26 | 88 | 168 | 1 - 949 | 15 - 963 | Arg-3 to Ser-14 His-64 to Lys-69. |
| 27 | 37 | 117 | 1 - 329 | 15 - 343 | Leu-46 to Lys-72. |
| 27 | 89 | 169 | 1 - 565 | 15 - 579 | Leu-46 to Ile-73. |
| 28 | 38 | 118 | 1 - 431 | 15 - 445 | Glu-34 to Gly-39. |
| 28 | 90 | 170 | 1 - 1302 | 15 - 1316 | Glu-34 to Gly-39. |

Table 3

| Clone ID | Library Codes |
|----------|---|
| HWEAC77 | H0081 H0264 H0601 L1290 S0434 |
| HRAEH37 | H0013 H0036 H0039 H0046 H0081 H0135 H0156 H0265 H0457 H0486 H0494 H0520 H0521 H0539 H0551 H0553 H0555 H0556 H0572 H0575 H0593 H0618 H0645 H0653 H0658 H0668 H0670 H0676 L1290 S0007 S0011 S0036 S0038 S0126 S0132 S0278 S0328 S0354 S0374 S0394 T0006 |
| HDQGO29 | H0521 H0543 S0426 |
| HDQHY04 | H0013 H0014 H0032 H0040 H0056 H0170 H0244 H0265 H0328 H0380 H0427 H0435 H0519 H0520 H0521 H0522 H0539 H0553 H0575 H0583 H0587 H0625 H0638 L1290 S0032 S0242 S0250 S0358 S0376 S0436 S0460 S3014 T0041 T0114 |
| HWDAH38 | H0069 H0097 H0179 H0250 H0271 H0416 H0575 H0580 H0581 H0600 H0660 H0670 H0689 L1290 S0052 S0053 S0216 S0360 S0426 S0428 |
| HHESQ62 | H0445 H0543 S0052 S0114 |
| HDP SH53 | H0521 |
| HDPRN70 | H0521 |
| HLWCP78 | H0008 H0040 H0041 H0045 H0046 H0052 H0081 H0123 H0188 H0250 H0251 H0268 H0333 H0428 H0486 H0529 H0547 H0553 H0647 H0673 L1290 S0010 S0028 S0042 S0051 S0192 S0250 S0342 S0360 S0380 T0109 T0115 |
| HTJML75 | H0012 H0013 H0052 H0123 H0253 H0295 H0341 H0413 H0436 H0457 H0483 H0488 H0494 H0506 H0519 H0521 H0522 H0529 H0539 H0542 H0547 H0550 H0618 H0619 H0623 H0687 H0690 L1290 S0002 S0027 S0037 S0052 S0114 S0150 S0190 S0222 S0328 S0378 S0424 |
| HWBFE57 | H0069 H0255 H0264 H0457 H0543 H0580 H0593 H0620 H0650 |
| HDPLO25 | H0009 H0014 H0036 H0052 H0090 H0135 H0163 H0265 H0268 H0270 H0275 H0411 H0423 H0427 H0445 H0457 H0486 H0506 H0517 H0521 H0543 H0548 H0549 H0556 H0574 H0580 H0581 H0587 H0593 H0597 H0600 H0615 H0649 H0656 H0657 H0661 H0662 H0663 H0670 H0690 L1290 S0003 S0026 S0031 S0040 S0051 S0142 S0144 S0192 S0196 S0222 S0280 S0354 S0356 S0358 S0360 S0376 S0408 T0067 |
| HDPWN93 | H0038 H0046 H0052 H0063 H0087 H0131 H0135 H0194 H0207 H0251 H0253 H0255 H0263 H0309 H0333 H0409 H0412 H0417 H0424 H0494 H0506 H0521 H0543 H0545 H0555 H0580 H0589 H0590 H0597 H0616 H0618 H0622 H0646 H0648 H0672 L1290 S0010 S0044 S0045 S0150 S0152 S0222 S0340 S0356 S0358 S0624 T0010 T0060 T0082 |
| HWHPM16 | H0038 H0222 H0373 H0521 H0551 H0556 H0586 H0587 H0590 H0592 S0386 |
| HRAEE45 | H0543 H0555 H0592 |
| HDPXY01 | H0521 H0539 H0660 |
| HWBEM18 | H0009 H0050 H0051 H0052 H0069 H0083 H0134 H0216 H0264 H0265 H0290 H0309 H0318 H0333 H0341 H0393 H0422 H0424 H0445 H0457 H0520 H0521 H0522 H0529 H0543 H0556 H0580 H0581 H0610 H0617 H0620 H0634 H0635 H0637 H0645 H0650 H0656 H0670 H0701 L1290 S0038 S0426 |
| HDTFE17 | H0013 H0024 H0050 H0052 H0063 H0087 H0123 H0188 H0208 H0218 H0250 H0252 H0254 H0264 H0265 H0272 H0306 H0316 H0318 H0369 H0370 H0372 H0381 H0402 H0417 H0418 H0423 H0424 H0441 H0445 H0478 H0486 H0519 H0521 H0522 H0539 H0542 H0543 H0545 H0550 H0555 H0556 H0561 H0575 H0580 H0617 H0620 H0634 H0638 H0652 H0656 H0657 H0659 H0665 H0670 H0675 H0677 L1290 S0002 S0132 S0354 S0356 S0358 S0360 S0376 S0380 S0408 S0426 S0458 |

| | |
|---------|--|
| HBXAB02 | H0013 H0014 H0024 H0031 H0032 H0038 H0039 H0040 H0041 H0046 H0050 H0051 H0052 H0056 H0057 H0081 H0085 H0086 H0087 H0100 H0105 H0116 H0123 H0124 H0135 H0144 H0150 H0163 H0171 H0179 H0188 H0196 H0208 H0242 H0251 H0252 H0253 H0264 H0266 H0268 H0269 H0271 H0284 H0286 H0290 H0292 H0294 H0309 H0316 H0318 H0333 H0343 H0352 H0381 H0392 H0402 H0411 H0412 H0413 H0427 H0428 H0437 H0444 H0445 H0484 H0485 H0486 H0506 H0518 H0519 H0520 H0521 H0522 H0529 H0539 H0542 H0543 H0544 H0545 H0546 H0547 H0549 H0550 H0551 H0553 H0559 H0575 H0580 H0586 H0587 H0590 H0592 H0593 H0594 H0597 H0598 H0599 H0600 H0602 H0618 H0619 H0620 H0622 H0623 H0624 H0626 H0628 H0631 H0638 H0647 H0648 H0653 H0656 H0657 H0660 H0664 H0665 H0667 H0682 H0684 H0687 H0688 H0689 H0690 H0691 H0696 L1290 S0002 S0011 S0022 S0026 S0027 S0028 S0036 S0037 S0038 S0040 S0044 S0045 S0046 S0116 S0126 S0192 S0194 S0196 S0208 S0210 S0212 S0222 S0242 S0250 S0294 S0330 S0332 S0342 S0352 S0354 S0356 S0358 S0360 S0364 S0366 S0376 S0406 S0418 S0432 S0446 S3012 S3014 S6024 T0003 T0004 T0010 T0040 T0049 |
| HDPSB18 | H0100 H0149 H0178 H0181 H0204 H0295 H0318 H0381 H0412 H0483 H0484 H0521 H0529 H0543 H0620 H0623 H0633 H0638 H0646 H0656 H0690 L1290 S0045 S0278 S0358 S0360 S0420 S3014 T0049 |
| HDPTW65 | H0053 H0521 |
| HOHDF66 | H0628 L1290 S0126 S0250 |
| HDPTW24 | H0521 |
| HRADO01 | H0555 |
| HTJNX29 | H0036 H0041 H0488 H0509 H0586 H0622 S0434 |
| HOHDC86 | S0250 |
| HLDQA07 | H0510 |
| HWDAD17 | H0036 H0163 H0270 H0275 H0411 H0427 H0445 H0457 H0506 H0517 H0521 H0574 H0593 H0600 H0615 H0649 H0663 L1290 S0003 S0026 S0031 S0040 S0051 S0142 S0196 S0222 S0280 S0354 S0358 S0360 S0376 S0408 T0067 |

Table 4

| SEQ ID NO: X | Cytologic Band or Chromosome: | OMIM Reference(s): | | | | | |
|-----------------|----------------------------------|--------------------|--------|--------|--------|--------|--------|
| 12 | 16p12-p11 | 108730 | 147781 | 172471 | 182381 | 186580 | 266600 |
| | | 600760 | 600761 | 602066 | | | |

Table 5

| Library Code | Library Description | Disease |
|--------------|--|---------|
| H0008 | Whole 6 Week Old Embryo | |
| H0009 | Human Fetal Brain | |
| H0012 | Human Fetal Kidney | |
| H0013 | Human 8 Week Whole Embryo | |
| H0014 | Human Gall Bladder | |
| H0024 | Human Fetal Lung III | |
| H0031 | Human Placenta | |
| H0032 | Human Prostate | |
| H0036 | Human Adult Small Intestine | |
| H0038 | Human Testes | |
| H0039 | Human Pancreas Tumor | disease |
| H0040 | Human Testes Tumor | disease |
| H0041 | Human Fetal Bone | |
| H0045 | Human Esophagus, Cancer | disease |
| H0046 | Human Endometrial Tumor | disease |
| H0050 | Human Fetal Heart | |
| H0051 | Human Hippocampus | |
| H0052 | Human Cerebellum | |
| H0053 | Human Adult Kidney | |
| H0056 | Human Umbilical Vein, Endo. remake | |
| H0057 | Human Fetal Spleen | |
| H0063 | Human Thymus | |
| H0069 | Human Activated T-Cells | |
| H0081 | Human Fetal Epithelium (Skin) | |
| H0083 | HUMAN JURKAT MEMBRANE BOUND POLYSOMES | |
| H0085 | Human Colon | |
| H0086 | Human epithelioid sarcoma | disease |
| H0087 | Human Thymus | |
| H0090 | Human T-Cell Lymphoma | disease |
| H0097 | Human Adult Heart, subtracted | |
| H0100 | Human Whole Six Week Old Embryo | |
| H0105 | Human Fetal Heart, subtracted | |
| H0116 | Human Thymus Tumor, subtracted | |
| H0123 | Human Fetal Dura Mater | |
| H0124 | Human Rhabdomyosarcoma | disease |
| H0131 | LNCAP + 0.3nM R1881 | |
| H0134 | Raji Cells, cyclohexamide treated | |
| H0135 | Human Synovial Sarcoma | |
| H0144 | Nine Week Old Early Stage Human | |
| H0149 | 7 Week Old Early Stage Human, subtracted | |
| H0150 | Human Epididymus | |
| H0156 | Human Adrenal Gland Tumor | disease |
| H0163 | Human Synovium | |
| H0170 | 12 Week Old Early Stage Human | |
| H0171 | 12 Week Old Early Stage Human, II | |
| H0178 | Human Fetal Brain | |
| H0179 | Human Neutrophil | |
| H0181 | Human Primary Breast Cancer | disease |
| H0188 | Human Normal Breast | |
| H0194 | Human Cerebellum, subtracted | |

| | | |
|-------|--|---------|
| H0196 | Human Cardiomyopathy, subtracted | |
| H0204 | Human Colon Cancer, subtracted | |
| H0207 | LNCAP, differential expression | |
| H0208 | Early Stage Human Lung, subtracted | |
| H0216 | Supt cells, cyclohexamide treated, subtracted | |
| H0218 | Activated T-Cells, 0hrs, subtracted | |
| H0222 | Activated T-Cells, 8 hrs, subtracted | |
| H0242 | Human Fetal Heart, Differential (Fetal-Specific) | |
| H0244 | Human 8 Week Whole Embryo, subtracted | |
| H0250 | Human Activated Monocytes | |
| H0251 | Human Chondrosarcoma | disease |
| H0252 | Human Osteosarcoma | disease |
| H0253 | Human adult testis, large inserts | |
| H0254 | Breast Lymph node cDNA library | |
| H0255 | breast lymph node CDNA library | |
| H0263 | human colon cancer | disease |
| H0264 | human tonsils | |
| H0265 | Activated T-Cell (12hs)/Thiouridine labelledEco | |
| H0266 | Human Microvascular Endothelial Cells, fract. A | |
| H0268 | Human Umbilical Vein Endothelial Cells, fract. A | |
| H0269 | Human Umbilical Vein Endothelial Cells, fract. B | |
| H0270 | HPAS (human pancreas, subtracted) | |
| H0271 | Human Neutrophil, Activated | |
| H0272 | HUMAN TONSILS, FRACTION 2 | |
| H0275 | Human Infant Adrenal Gland, Subtracted | |
| H0284 | Human OB MG63 control fraction I | |
| H0286 | Human OB MG63 treated (10 nM E2) fraction I | |
| H0290 | Human OB HOS treated (1 nM E2) fraction I | |
| H0292 | Human OB HOS treated (10 nM E2) fraction I | |
| H0294 | Amniotic Cells - TNF induced | |
| H0295 | Amniotic Cells - Primary Culture | |
| H0306 | CD34 depleted Buffy Coat (Cord Blood) | |
| H0309 | Human Chronic Synovitis | disease |
| H0316 | HUMAN STOMACH | |
| H0318 | HUMAN B CELL LYMPHOMA | disease |
| H0328 | human ovarian cancer | disease |
| H0333 | Hemangiopericytoma | disease |
| H0341 | Bone Marrow Cell Line (RS4,11) | |
| H0343 | stomach cancer (human) | disease |
| H0352 | wilm's tumor | disease |
| H0369 | H. Atrophic Endometrium | |
| H0370 | H. Lymph node breast Cancer | disease |
| H0372 | Human Testes | |
| H0373 | Human Heart | |
| H0380 | Human Tongue, frac 2 | |
| H0381 | Bone Cancer | disease |
| H0392 | H. Meningioma, M1 | |
| H0393 | Fetal Liver, subtraction II | |
| H0402 | CD34 depleted Buffy Coat (Cord Blood), re-excision | |
| H0409 | H. Striatum Depression, subtracted | |
| H0411 | H Female Bladder, Adult | |
| H0412 | Human umbilical vein endothelial cells, IL-4 induced | |
| H0413 | Human Umbilical Vein Endothelial Cells, uninduced | |

| | | |
|-------|---|---------|
| H0416 | Human Neutrophils, Activated, re-excision | |
| H0417 | Human Pituitary, subtracted VIII | |
| H0418 | Human Pituitary, subtracted VII | |
| H0422 | T-Cell PHA 16 hrs | |
| H0423 | T-Cell PHA 24 hrs | |
| H0424 | Human Pituitary, subt IX | |
| H0427 | Human Adipose | |
| H0428 | Human Ovary | |
| H0435 | Ovarian Tumor 10-3-95 | |
| H0436 | Resting T-Cell Library,II | |
| H0437 | H Umbilical Vein Endothelial Cells, frac A, re-excision | |
| H0441 | H. Kidney Cortex, subtracted | |
| H0444 | Spleen metastatic melanoma | disease |
| H0445 | Spleen, Chronic lymphocytic leukemia | disease |
| H0457 | Human Eosinophils | |
| H0478 | Salivary Gland, Lib 2 | |
| H0483 | Breast Cancer cell line, MDA 36 | |
| H0484 | Breast Cancer Cell line, angiogenic | |
| H0485 | Hodgkin's Lymphoma I | disease |
| H0486 | Hodgkin's Lymphoma II | disease |
| H0488 | Human Tonsils, Lib 2 | |
| H0494 | Keratinocyte | |
| H0506 | Ulcerative Colitis | |
| H0509 | Liver, Hepatoma | disease |
| H0510 | Human Liver, normal | |
| H0517 | Nasal polyps | |
| H0518 | pBMC stimulated w/ poly I/C | |
| H0519 | NTERA2, control | |
| H0520 | NTERA2 + retinoic acid, 14 days | |
| H0521 | Primary Dendritic Cells, lib 1 | |
| H0522 | Primary Dendritic cells, frac 2 | |
| H0529 | Myeloid Progenitor Cell Line | |
| H0539 | Pancreas Islet Cell Tumor | disease |
| H0542 | T Cell helper I | |
| H0543 | T cell helper II | |
| H0544 | Human endometrial stromal cells | |
| H0545 | Human endometrial stromal cells-treated with progesterone | |
| H0546 | Human endometrial stromal cells-treated with estradiol | |
| H0547 | NTERA2 teratocarcinoma cell line+retinoic acid (14 days) | |
| H0548 | Human Skin Fibroblasts, normal | |
| H0549 | H. Epididymus, caput & corpus | |
| H0550 | H. Epididymus, cauda | |
| H0551 | Human Thymus Stromal Cells | |
| H0553 | Human Placenta | |
| H0555 | Rejected Kidney, lib 4 | disease |
| H0556 | Activated T-cell(12h)/Thiouridine-re-excision | |
| H0559 | HL-60, PMA 4H, re-excision | |
| H0561 | L428 | |
| H0572 | Human Fetal Brain, normalized AC5002 | |
| H0574 | Hepatocellular Tumor, re-excision | disease |
| H0575 | Human Adult Pulmonary, re-excision | |
| H0580 | Dendritic cells, pooled | |
| H0581 | Human Bone Marrow, treated | |

| | | |
|-------|---|---------|
| H0583 | B Cell lymphoma | disease |
| H0586 | Healing groin wound, 6.5 hours post incision | disease |
| H0587 | Healing groin wound, 7.5 hours post incision | disease |
| H0589 | CD34 positive cells (cord blood),re-ex | |
| H0590 | Human adult small intestine,re-excision | |
| H0592 | Healing groin wound - zero hr post-incision (control) | disease |
| H0593 | Olfactory epithelium,nasalcavity | |
| H0594 | Human Lung Cancer,re-excision | disease |
| H0597 | Human Colon, re-excision | |
| H0598 | Human Stomach,re-excision | |
| H0599 | Human Adult Heart,re-excision | |
| H0600 | Healing Abdomen wound,70&90 min post incision | disease |
| H0601 | Healing Abdomen Wound,15 days post incision | disease |
| H0602 | Healing Abdomen Wound,21&29 days post incision | disease |
| H0610 | H. Leukocytes, normalized cot 5A | |
| H0615 | Human Ovarian Cancer Reexcision | disease |
| H0616 | Human Testes, Reexcision | |
| H0617 | Human Primary Breast Cancer Reexcision | disease |
| H0618 | Human Adult Testes, Large Inserts, Reexcision | |
| H0619 | Fetal Heart | |
| H0620 | Human Fetal Kidney, Reexcision | |
| H0622 | Human Pancreas Tumor, Reexcision | disease |
| H0623 | Human Umbilical Vein, Reexcision | |
| H0624 | 12 Week Early Stage Human II, Reexcision | |
| H0625 | Ku 812F Basophils Line | |
| H0626 | Saos2 Cells, Untreated | |
| H0628 | Human Pre-Differentiated Adipocytes | |
| H0631 | Saos2, Dexamethosome Treated | |
| H0633 | Lung Carcinoma A549 TNFalpha activated | disease |
| H0634 | Human Testes Tumor, re-excision | disease |
| H0635 | Human Activated T-Cells, re-excision | |
| H0637 | Dendritic Cells From CD34 Cells | |
| H0638 | CD40 activated monocyte dendritic cells | |
| H0645 | Fetal Heart, re-excision | |
| H0646 | Lung, Cancer (4005313 A3): Invasive Poorly Differentiated Lung Adenocarcinoma, | |
| H0647 | Lung, Cancer (4005163 B7): Invasive, Poorly Diff. Adenocarcinoma, Metastatic | disease |
| H0648 | Ovary, Cancer: (4004562 B6) Papillary Serous Cystic Neoplasm, Low Malignant Pot | disease |
| H0649 | Lung, Normal: (4005313 B1) | |
| H0650 | B-Cells | |
| H0652 | Lung, Normal: (4005313 B1) | |
| H0653 | Stromal Cells | |
| H0656 | B-cells (unstimulated) | |
| H0657 | B-cells (stimulated) | |
| H0658 | Ovary, Cancer (9809C332): Poorly differentiated adenocarcinoma | disease |
| H0659 | Ovary, Cancer (15395A1F): Grade II Papillary Carcinoma | disease |
| H0660 | Ovary, Cancer: (15799A1F) Poorly differentiated carcinoma | disease |
| H0661 | Breast, Cancer: (4004943 A5) | disease |
| H0662 | Breast, Normal: (4005522B2) | |
| H0663 | Breast, Cancer: (4005522 A2) | disease |
| H0664 | Breast, Cancer: (9806C012R) | disease |

| | | |
|-------|--|---------|
| H0665 | Stromal cells 3.88 | |
| H0667 | Stromal cells(HBM3.18) | |
| H0668 | stromal cell clone 2.5 | |
| H0670 | Ovary, Cancer(4004650 A3): Well-Differentiated Micropapillary Serous Carcinoma | |
| H0672 | Ovary, Cancer: (4004576 A8) | |
| H0673 | Human Prostate Cancer, Stage B2, re-excision | |
| H0675 | Colon, Cancer: (9808C064R) | |
| H0676 | Colon, Cancer: (9808C064R)-total RNA | |
| H0677 | TNFR degenerate oligo | |
| H0682 | Ovarian cancer, Serous Papillary Adenocarcinoma | |
| H0684 | Ovarian cancer, Serous Papillary Adenocarcinoma | |
| H0687 | Human normal ovary(#9610G215) | |
| H0688 | Human Ovarian Cancer(#9807G017) | |
| H0689 | Ovarian Cancer | |
| H0690 | Ovarian Cancer, # 9702G001 | |
| H0691 | Normal Ovary, #9710G208 | |
| H0696 | Prostate Adenocarcinoma | |
| H0701 | NKyao15(control) | |
| L1290 | Soares total fetus Nb2HF8 9w | |
| S0002 | Monocyte activated | |
| S0003 | Human Osteoclastoma | disease |
| S0007 | Early Stage Human Brain | |
| S0010 | Human Amygdala | |
| S0011 | STROMAL -OSTEOCLASTOMA | disease |
| S0022 | Human Osteoclastoma Stromal Cells - unamplified | |
| S0026 | Stromal cell TF274 | |
| S0027 | Smooth muscle, serum treated | |
| S0028 | Smooth muscle,control | |
| S0031 | Spinal cord | |
| S0032 | Smooth muscle-ILb induced | |
| S0036 | Human Substantia Nigra | |
| S0037 | Smooth muscle, IL1b induced | |
| S0038 | Human Whole Brain #2 - Oligo dT > 1.5Kb | |
| S0040 | Adipocytes | |
| S0042 | Testes | |
| S0044 | Prostate BPH | disease |
| S0045 | Endothelial cells-control | |
| S0046 | Endothelial-induced | |
| S0051 | Human Hypothalamus,Schizophrenia | disease |
| S0052 | neutrophils control | |
| S0053 | Neutrophils IL-1 and LPS induced | |
| S0114 | Anergic T-cell | |
| S0116 | Bone marrow | |
| S0126 | Osteoblasts | |
| S0132 | Epithelial-TNFa and INF induced | |
| S0142 | Macrophage-oxLDL | |
| S0144 | Macrophage (GM-CSF treated) | |
| S0150 | LNCAP prostate cell line | |
| S0152 | PC3 Prostate cell line | |
| S0190 | Prostate BPH,Lib 2, subtracted | |
| S0192 | Synovial Fibroblasts (control) | |
| S0194 | Synovial hypoxia | |
| S0196 | Synovial IL-1/TNF stimulated | |

| | | |
|-------|--|---------|
| S0208 | Messangial cell, frac 1 | |
| S0210 | Messangial cell, frac 2 | |
| S0212 | Bone Marrow Stromal Cell, untreated | |
| S0216 | Neutrophils IL-1 and LPS induced | |
| S0222 | H. Frontal cortex,epileptic,re-excision | disease |
| S0242 | Synovial Fibroblasts (Il1/TNF), subt | |
| S0250 | Human Osteoblasts II | disease |
| S0278 | H Macrophage (GM-CSF treated), re-excision | |
| S0280 | Human Adipose Tissue, re-excision | |
| S0294 | Larynx tumor | disease |
| S0328 | Palate carcinoma | disease |
| S0330 | Palate normal | |
| S0332 | Pharynx carcinoma | |
| S0340 | Human Osteoarthritic Cartilage Fraction IV | disease |
| S0342 | Adipocytes,re-excision | |
| S0352 | Larynx Carcinoma | disease |
| S0354 | Colon Normal II | |
| S0356 | Colon Carcinoma | disease |
| S0358 | Colon Normal III | |
| S0360 | Colon Tumor II | disease |
| S0364 | Human Quadriceps | |
| S0366 | Human Soleus | |
| S0374 | Normal colon | |
| S0376 | Colon Tumor | disease |
| S0378 | Pancreas normal PCA4 No | |
| S0380 | Pancreas Tumor PCA4 Tu | disease |
| S0386 | Human Whole Brain, re-excision | |
| S0394 | Stomach,normal | |
| S0406 | Rectum tumour | |
| S0408 | Colon, normal | |
| S0418 | CHME Cell Line,treated 5 hrs | |
| S0420 | CHME Cell Line,untreated | |
| S0424 | TF-1 Cell Line GM-CSF Treated | |
| S0426 | Monocyte activated, re-excision | |
| S0428 | Neutrophils control, re-excision | |
| S0432 | Sinus piriformis Tumour | |
| S0434 | Stomach Normal | disease |
| S0436 | Stomach Tumour | disease |
| S0446 | Tongue Tumour | |
| S0458 | Thyroid Normal (SDCA2 No) | |
| S0460 | Thyroid Tumour | |
| S3012 | Smooth Muscle Serum Treated, Norm | |
| S3014 | Smooth muscle, serum induced,re-exc | |
| S6024 | Alzheimers, spongy change | disease |
| T0003 | Human Fetal Lung | |
| T0004 | Human White Fat | |
| T0006 | Human Pineal Gland | |
| T0010 | Human Infant Brain | |
| T0040 | HSC172 cells | |
| T0041 | Jurkat T-cell G1 phase | |
| T0049 | Aorta endothelial cells + TNF-a | |
| T0060 | Human White Adipose | |
| T0067 | Human Thyroid | |

| | | |
|-------|---|--|
| T0082 | Human Adult Retina | |
| T0109 | Human (HCC) cell line liver (mouse) metastasis, remake | |
| T0114 | Human (Caco-2) cell line, adenocarcinoma, colon, remake | |
| T0115 | Human Colon Carcinoma (HCC) cell line | |

Table 6

| OMIM ID | OMIM Description |
|---------|--|
| 108730 | Brody myopathy, 601003 (3) |
| 147781 | Atopy, susceptibility to (3) |
| 172471 | Glycogenosis, hepatic, autosomal (3) |
| 182381 | Renal glucosuria, 253100 (1) (?) |
| 186580 | Arthrocutaneous uveitis (2) |
| 266600 | Inflammatory bowel disease-1 (2) |
| 600760 | Liddle syndrome, 177200 (3) Pseudohypoaldosteronism, type I, 264350 (3) |
| 600761 | Liddle syndrome, 177200 (3) Pseudohypoaldosteronism, type I, 264350 (3) |
| 602066 | Convulsions, infantile and paroxysmal choreoathetosis (2) |

The polypeptides of the invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides
5 are well understood in the art.

The polypeptides may be in the form of the secreted protein, including the mature form, or may be a part of a larger protein, such as a fusion protein (see below). It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, pro-sequences, sequences which aid in purification ,
10 such as multiple histidine residues, or an additional sequence for stability during recombinant production.

The polypeptides of the present invention are preferably provided in an isolated form, and preferably are substantially purified. A recombinantly produced version of a polypeptide, including the secreted polypeptide, can be substantially
15 purified using techniques described herein or otherwise known in the art, such as, for example, by the one-step method described in Smith and Johnson, Gene 67:31-40 (1988). Polypeptides of the invention also can be purified from natural, synthetic or recombinant sources using techniques described herein or otherwise known in the art, such as, for example, antibodies of the invention raised against the secreted protein.

20 The present invention provides a polynucleotide comprising, or alternatively consisting of, the nucleic acid sequence of SEQ ID NO:X, and/or a cDNA contained in ATCC deposit Z. The present invention also provides a polypeptide comprising, or alternatively, consisting of, the polypeptide sequence of SEQ ID NO:Y and/or a polypeptide encoded by the cDNA contained in ATCC deposit Z. Polynucleotides
25 encoding a polypeptide comprising, or alternatively consisting of the polypeptide sequence of SEQ ID NO:Y and/or a polypeptide sequence encoded by the cDNA contained in ATCC deposit Z are also encompassed by the invention.

Signal Sequences

The present invention also encompasses mature forms of the polypeptide
30 having the polypeptide sequence of SEQ ID NO:Y and/or the polypeptide sequence encoded by the cDNA in a deposited clone. Polynucleotides encoding the mature forms (such as, for example, the polynucleotide sequence in SEQ ID NO:X and/or the

polynucleotide sequence contained in the cDNA of a deposited clone) are also encompassed by the invention. According to the signal hypothesis, proteins secreted by mammalian cells have a signal or secretory leader sequence which is cleaved from the mature protein once export of the growing protein chain across the rough
5 endoplasmic reticulum has been initiated. Most mammalian cells and even insect cells cleave secreted proteins with the same specificity. However, in some cases, cleavage of a secreted protein is not entirely uniform, which results in two or more mature species of the protein. Further, it has long been known that cleavage specificity of a secreted protein is ultimately determined by the primary structure of
10 the complete protein, that is, it is inherent in the amino acid sequence of the polypeptide.

Methods for predicting whether a protein has a signal sequence, as well as the cleavage point for that sequence, are available. For instance, the method of McGeoch, *Virus Res.* 3:271-286 (1985), uses the information from a short N-terminal
15 charged region and a subsequent uncharged region of the complete (uncleaved) protein. The method of von Heinje, *Nucleic Acids Res.* 14:4683-4690 (1986) uses the information from the residues surrounding the cleavage site, typically residues -13 to +2, where +1 indicates the amino terminus of the secreted protein. The accuracy of predicting the cleavage points of known mammalian secretory proteins for each of
20 these methods is in the range of 75-80%. (von Heinje, *supra.*) However, the two methods do not always produce the same predicted cleavage point(s) for a given protein.

In the present case, the deduced amino acid sequence of the secreted polypeptide was analyzed by a computer program called SignalP (Henrik Nielsen et
25 al., *Protein Engineering* 10:1-6 (1997)), which predicts the cellular location of a protein based on the amino acid sequence. As part of this computational prediction of localization, the methods of McGeoch and von Heinje are incorporated. The analysis of the amino acid sequences of the secreted proteins described herein by this program provided the results shown in Table 1.

30 As one of ordinary skill would appreciate, however, cleavage sites sometimes vary from organism to organism and cannot be predicted with absolute certainty. Accordingly, the present invention provides secreted polypeptides having a sequence

shown in SEQ ID NO:Y which have an N-terminus beginning within 5 residues (i.e., + or - 5 residues) of the predicted cleavage point. Similarly, it is also recognized that in some cases, cleavage of the signal sequence from a secreted protein is not entirely uniform, resulting in more than one secreted species. These polypeptides, and the
5 polynucleotides encoding such polypeptides, are contemplated by the present invention.

Moreover, the signal sequence identified by the above analysis may not necessarily predict the naturally occurring signal sequence. For example, the naturally occurring signal sequence may be further upstream from the predicted signal
10 sequence. However, it is likely that the predicted signal sequence will be capable of directing the secreted protein to the ER. Nonetheless, the present invention provides the mature protein produced by expression of the polynucleotide sequence of SEQ ID NO:X and/or the polynucleotide sequence contained in the cDNA of a deposited clone, in a mammalian cell (e.g., COS cells, as described below). These polypeptides,
15 and the polynucleotides encoding such polypeptides, are contemplated by the present invention.

Polynucleotide and Polypeptide Variants

The present invention is directed to variants of the polynucleotide sequence disclosed in SEQ ID NO:X, the complementary strand thereto, and/or the cDNA
20 sequence contained in a deposited clone.

The present invention also encompasses variants of the polypeptide sequence disclosed in SEQ ID NO:Y and/or encoded by a deposited clone.

"Variant" refers to a polynucleotide or polypeptide differing from the
25 polynucleotide or polypeptide of the present invention, but retaining essential properties thereof. Generally, variants are overall closely similar, and, in many regions, identical to the polynucleotide or polypeptide of the present invention.

The present invention is also directed to nucleic acid molecules which comprise, or alternatively consist of, a nucleotide sequence which is at least 80%,
30 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to, for example, the nucleotide coding sequence in SEQ ID NO:X or the complementary strand thereto, the nucleotide coding sequence contained in a deposited cDNA clone or the

complementary strand thereto, a nucleotide sequence encoding the polypeptide of SEQ ID NO:Y, a nucleotide sequence encoding the polypeptide encoded by the cDNA contained in a deposited clone, and/or polynucleotide fragments of any of these nucleic acid molecules (e.g., those fragments described herein).

- 5 Polynucleotides which hybridize to these nucleic acid molecules under stringent hybridization conditions or lower stringency conditions are also encompassed by the invention, as are polypeptides encoded by these polynucleotides.

The present invention is also directed to polypeptides which comprise, or alternatively consist of, an amino acid sequence which is at least 80%, 85%, 90%, 10 95%, 96%, 97%, 98%, 99% identical to, for example, the polypeptide sequence shown in SEQ ID NO:Y, the polypeptide sequence encoded by the cDNA contained in a deposited clone, and/or polypeptide fragments of any of these polypeptides (e.g., those fragments described herein).

By a nucleic acid having a nucleotide sequence at least, for example, 95% 15 "identical" to a reference nucleotide sequence of the present invention, it is intended that the nucleotide sequence of the nucleic acid is identical to the reference sequence except that the nucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence encoding the polypeptide. In other words, to obtain a nucleic acid having a nucleotide sequence at least 95% 20 identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. The query sequence may be an entire sequence shown in Table 1, the ORF (open reading frame), or any fragment specified 25 as described herein.

As a practical matter, whether any particular nucleic acid molecule or polypeptide is at least 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to a nucleotide sequence of the present invention can be determined conventionally using known computer programs. A preferred method for determining the best 30 overall match between a query sequence (a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag et al. (Comp.

App. Biosci. 6:237-245(1990)). In a sequence alignment the query and subject sequences are both DNA sequences. An RNA sequence can be compared by converting U's to T's. The result of said global sequence alignment is in percent identity. Preferred parameters used in a FASTDB alignment of DNA sequences to
5 calculate percent identity are: Matrix=Unitary, k-tuple=4, Mismatch Penalty=1, Joining Penalty=30, Randomization Group Length=0, Cutoff Score=1, Gap Penalty=5, Gap Size Penalty 0.05, Window Size=500 or the length of the subject nucleotide sequence, whichever is shorter.

If the subject sequence is shorter than the query sequence because of 5' or 3'
10 deletions, not because of internal deletions, a manual correction must be made to the results. This is because the FASTDB program does not account for 5' and 3' truncations of the subject sequence when calculating percent identity. For subject sequences truncated at the 5' or 3' ends, relative to the query sequence, the percent identity is corrected by calculating the number of bases of the query sequence that are
15 5' and 3' of the subject sequence, which are not matched/aligned, as a percent of the total bases of the query sequence. Whether a nucleotide is matched/aligned is determined by results of the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This corrected
20 score is what is used for the purposes of the present invention. Only bases outside the 5' and 3' bases of the subject sequence, as displayed by the FASTDB alignment, which are not matched/aligned with the query sequence, are calculated for the purposes of manually adjusting the percent identity score.

For example, a 90 base subject sequence is aligned to a 100 base query
25 sequence to determine percent identity. The deletions occur at the 5' end of the subject sequence and therefore, the FASTDB alignment does not show a matched/alignment of the first 10 bases at 5' end. The 10 unpaired bases represent 10% of the sequence (number of bases at the 5' and 3' ends not matched/total number of bases in the query sequence) so 10% is subtracted from the percent identity score
30 calculated by the FASTDB program. If the remaining 90 bases were perfectly matched the final percent identity would be 90%. In another example, a 90 base subject sequence is compared with a 100 base query sequence. This time the

deletions are internal deletions so that there are no bases on the 5' or 3' of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only bases 5' and 3' of the subject sequence which are not matched/aligned with the query sequence are manually corrected for. No other manual corrections are to be made for the purposes of the present invention.

By a polypeptide having an amino acid sequence at least, for example, 95% "identical" to a query amino acid sequence of the present invention, it is intended that the amino acid sequence of the subject polypeptide is identical to the query sequence except that the subject polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the query amino acid sequence. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a query amino acid sequence, up to 5% of the amino acid residues in the subject sequence may be inserted, deleted, (indels) or substituted with another amino acid. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

As a practical matter, whether any particular polypeptide is at least 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to, for instance, an amino acid sequences shown in Table 1 (SEQ ID NO:Y) or to the amino acid sequence encoded by cDNA contained in a deposited clone can be determined conventionally using known computer programs. A preferred method for determining the best overall match between a query sequence (a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag et al. (Comp. App. Biosci. 6:237-245(1990)). In a sequence alignment the query and subject sequences are either both nucleotide sequences or both amino acid sequences. The result of said global sequence alignment is in percent identity. Preferred parameters used in a FASTDB amino acid alignment are: Matrix=PAM 0, k-tuple=2, Mismatch Penalty=1, Joining Penalty=20, Randomization Group Length=0, Cutoff Score=1, Window Size=sequence length, Gap Penalty=5, Gap Size Penalty=0.05, Window

Size=500 or the length of the subject amino acid sequence, whichever is shorter.

If the subject sequence is shorter than the query sequence due to N- or C-terminal deletions, not because of internal deletions, a manual correction must be made to the results. This is because the FASTDB program does not account for N- and C-terminal truncations of the subject sequence when calculating global percent identity. For subject sequences truncated at the N- and C-termini, relative to the query sequence, the percent identity is corrected by calculating the number of residues of the query sequence that are N- and C-terminal of the subject sequence, which are not matched/aligned with a corresponding subject residue, as a percent of the total bases of the query sequence. Whether a residue is matched/aligned is determined by results of the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This final percent identity score is what is used for the purposes of the present invention. Only residues to the N- and C-termini of the subject sequence, which are not matched/aligned with the query sequence, are considered for the purposes of manually adjusting the percent identity score. That is, only query residue positions outside the farthest N- and C-terminal residues of the subject sequence.

For example, a 90 amino acid residue subject sequence is aligned with a 100 residue query sequence to determine percent identity. The deletion occurs at the N-terminus of the subject sequence and therefore, the FASTDB alignment does not show a matching/alignment of the first 10 residues at the N-terminus. The 10 unpaired residues represent 10% of the sequence (number of residues at the N- and C-termini not matched/total number of residues in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 residues were perfectly matched the final percent identity would be 90%. In another example, a 90 residue subject sequence is compared with a 100 residue query sequence. This time the deletions are internal deletions so there are no residues at the N- or C-termini of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only residue positions outside the N- and C-terminal ends of the subject sequence, as displayed in the FASTDB alignment, which are not

matched/aligned with the query sequence are manually corrected for. No other manual corrections are to be made for the purposes of the present invention.

The variants may contain alterations in the coding regions, non-coding regions, or both. Especially preferred are polynucleotide variants containing alterations which produce silent substitutions, additions, or deletions, but do not alter the properties or activities of the encoded polypeptide. Nucleotide variants produced by silent substitutions due to the degeneracy of the genetic code are preferred. Moreover, variants in which 5-10, 1-5, or 1-2 amino acids are substituted, deleted, or added in any combination are also preferred. Polynucleotide variants can be produced for a variety of reasons, e.g., to optimize codon expression for a particular host (change codons in the human mRNA to those preferred by a bacterial host such as *E. coli*).

Naturally occurring variants are called "allelic variants," and refer to one of several alternate forms of a gene occupying a given locus on a chromosome of an organism. (Genes II, Lewin, B., ed., John Wiley & Sons, New York (1985).) These allelic variants can vary at either the polynucleotide and/or polypeptide level and are included in the present invention. Alternatively, non-naturally occurring variants may be produced by mutagenesis techniques or by direct synthesis.

Using known methods of protein engineering and recombinant DNA technology, variants may be generated to improve or alter the characteristics of the polypeptides of the present invention. For instance, one or more amino acids can be deleted from the N-terminus or C-terminus of the secreted protein without substantial loss of biological function. The authors of Ron et al., *J. Biol. Chem.* 268: 2984-2988 (1993), reported variant KGF proteins having heparin binding activity even after deleting 3, 8, or 27 amino-terminal amino acid residues. Similarly, Interferon gamma exhibited up to ten times higher activity after deleting 8-10 amino acid residues from the carboxy terminus of this protein. (Dobeli et al., *J. Biotechnology* 7:199-216 (1988).)

Moreover, ample evidence demonstrates that variants often retain a biological activity similar to that of the naturally occurring protein. For example, Gayle and coworkers (*J. Biol. Chem* 268:22105-22111 (1993)) conducted extensive mutational analysis of human cytokine IL-1a. They used random mutagenesis to generate over

3,500 individual IL-1a mutants that averaged 2.5 amino acid changes per variant over the entire length of the molecule. Multiple mutations were examined at every possible amino acid position. The investigators found that "[m]ost of the molecule could be altered with little effect on either [binding or biological activity]." (See, 5 Abstract.) In fact, only 23 unique amino acid sequences, out of more than 3,500 nucleotide sequences examined, produced a protein that significantly differed in activity from wild-type.

Furthermore, even if deleting one or more amino acids from the N-terminus or C-terminus of a polypeptide results in modification or loss of one or more biological 10 functions, other biological activities may still be retained. For example, the ability of a deletion variant to induce and/or to bind antibodies which recognize the secreted form will likely be retained when less than the majority of the residues of the secreted form are removed from the N-terminus or C-terminus. Whether a particular polypeptide lacking N- or C-terminal residues of a protein retains such immunogenic 15 activities can readily be determined by routine methods described herein and otherwise known in the art.

Thus, the invention further includes polypeptide variants which show substantial biological activity. Such variants include deletions, insertions, inversions, repeats, and substitutions selected according to general rules known in the 20 art so as have little effect on activity. For example, guidance concerning how to make phenotypically silent amino acid substitutions is provided in Bowie et al., Science 247:1306-1310 (1990), wherein the authors indicate that there are two main strategies for studying the tolerance of an amino acid sequence to change.

The first strategy exploits the tolerance of amino acid substitutions by natural 25 selection during the process of evolution. By comparing amino acid sequences in different species, conserved amino acids can be identified. These conserved amino acids are likely important for protein function. In contrast, the amino acid positions where substitutions have been tolerated by natural selection indicates that these positions are not critical for protein function. Thus, positions tolerating amino acid 30 substitution could be modified while still maintaining biological activity of the protein.

The second strategy uses genetic engineering to introduce amino acid changes at specific positions of a cloned gene to identify regions critical for protein function. For example, site directed mutagenesis or alanine-scanning mutagenesis (introduction of single alanine mutations at every residue in the molecule) can be used.

5 (Cunningham and Wells, Science 244:1081-1085 (1989).) The resulting mutant molecules can then be tested for biological activity.

As the authors state, these two strategies have revealed that proteins are surprisingly tolerant of amino acid substitutions. The authors further indicate which amino acid changes are likely to be permissive at certain amino acid positions in the
10 protein. For example, most buried (within the tertiary structure of the protein) amino acid residues require nonpolar side chains, whereas few features of surface side chains are generally conserved. Moreover, tolerated conservative amino acid substitutions involve replacement of the aliphatic or hydrophobic amino acids Ala, Val, Leu and Ile; replacement of the hydroxyl residues Ser and Thr; replacement of the acidic
15 residues Asp and Glu; replacement of the amide residues Asn and Gln, replacement of the basic residues Lys, Arg, and His; replacement of the aromatic residues Phe, Tyr, and Trp, and replacement of the small-sized amino acids Ala, Ser, Thr, Met, and Gly.

Besides conservative amino acid substitution, variants of the present invention include (i) substitutions with one or more of the non-conserved amino acid residues,
20 where the substituted amino acid residues may or may not be one encoded by the genetic code, or (ii) substitution with one or more of amino acid residues having a substituent group, or (iii) fusion of the mature polypeptide with another compound, such as a compound to increase the stability and/or solubility of the polypeptide (for example, polyethylene glycol), or (iv) fusion of the polypeptide with additional amino
25 acids, such as, for example, an IgG Fc fusion region peptide, or leader or secretory sequence, or a sequence facilitating purification or (v) fusion of the polypeptide with another compound, such as albumin (including, but not limited to, recombinant albumin (see, e.g., U.S. Patent No. 5,876,969, issued March 2, 1999, EP Patent 0 413 622, and U.S. Patent No. 5,766,883, issued June 16, 1998, herein incorporated by
30 reference in their entirety)). Such variant polypeptides are deemed to be within the scope of those skilled in the art from the teachings herein.

For example, polypeptide variants containing amino acid substitutions of charged amino acids with other charged or neutral amino acids may produce proteins with improved characteristics, such as less aggregation. Aggregation of pharmaceutical formulations both reduces activity and increases clearance due to the aggregate's immunogenic activity. (Pinckard et al., Clin. Exp. Immunol. 2:331-340 (1967); Robbins et al., Diabetes 36: 838-845 (1987); Cleland et al., Crit. Rev. Therapeutic Drug Carrier Systems 10:307-377 (1993).)

A further embodiment of the invention relates to a polypeptide which comprises the amino acid sequence of the present invention having an amino acid sequence which contains at least one amino acid substitution, but not more than 50 amino acid substitutions, even more preferably, not more than 40 amino acid substitutions, still more preferably, not more than 30 amino acid substitutions, and still even more preferably, not more than 20 amino acid substitutions. Of course, in order of ever-increasing preference, it is highly preferable for a peptide or polypeptide to have an amino acid sequence which comprises the amino acid sequence of the present invention, which contains at least one, but not more than 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1 amino acid substitutions. In specific embodiments, the number of additions, substitutions, and/or deletions in the amino acid sequence of the present invention or fragments thereof (e.g., the mature form and/or other fragments described herein), is 1-5, 5-10, 5-25, 5-50, 10-50 or 50-150, conservative amino acid substitutions are preferable.

Polynucleotide and Polypeptide Fragments

The present invention is also directed to polynucleotide fragments of the polynucleotides of the invention.

In the present invention, a "polynucleotide fragment" refers to a short polynucleotide having a nucleic acid sequence which: is a portion of that contained in a deposited clone, or encoding the polypeptide encoded by the cDNA in a deposited clone; is a portion of that shown in SEQ ID NO:X or the complementary strand thereto, or is a portion of a polynucleotide sequence encoding the polypeptide of SEQ ID NO:Y. The nucleotide fragments of the invention are preferably at least about 15 nt, and more preferably at least about 20 nt, still more preferably at least about 30 nt,

and even more preferably, at least about 40 nt, at least about 50 nt, at least about 75 nt, or at least about 150 nt in length. A fragment "at least 20 nt in length," for example, is intended to include 20 or more contiguous bases from the cDNA sequence contained in a deposited clone or the nucleotide sequence shown in SEQ ID NO:X. In this context "about" includes the particularly recited value, a value larger or smaller by several (5, 4, 3, 2, or 1) nucleotides, at either terminus or at both termini. These nucleotide fragments have uses that include, but are not limited to, as diagnostic probes and primers as discussed herein. Of course, larger fragments (e.g., 50, 150, 500, 600, 2000 nucleotides) are preferred.

Moreover, representative examples of polynucleotide fragments of the invention, include, for example, fragments comprising, or alternatively consisting of, a sequence from about nucleotide number 1-50, 51-100, 101-150, 151-200, 201-250, 251-300, 301-350, 351-400, 401-450, 451-500, 501-550, 551-600, 651-700, 701-750, 751-800, 800-850, 851-900, 901-950, 951-1000, 1001-1050, 1051-1100, 1101-1150, 1151-1200, 1201-1250, 1251-1300, 1301-1350, 1351-1400, 1401-1450, 1451-1500, 1501-1550, 1551-1600, 1601-1650, 1651-1700, 1701-1750, 1751-1800, 1801-1850, 1851-1900, 1901-1950, 1951-2000, or 2001 to the end of SEQ ID NO:X, or the complementary strand thereto, or the cDNA contained in a deposited clone. In this context "about" includes the particularly recited ranges, and ranges larger or smaller by several (5, 4, 3, 2, or 1) nucleotides, at either terminus or at both termini. Preferably, these fragments encode a polypeptide which has biological activity. More preferably, these polynucleotides can be used as probes or primers as discussed herein. Polynucleotides which hybridize to these nucleic acid molecules under stringent hybridization conditions or lower stringency conditions are also encompassed by the invention, as are polypeptides encoded by these polynucleotides.

In the present invention, a "polypeptide fragment" refers to an amino acid sequence which is a portion of that contained in SEQ ID NO:Y or encoded by the cDNA contained in a deposited clone. Protein (polypeptide) fragments may be "free-standing," or comprised within a larger polypeptide of which the fragment forms a part or region, most preferably as a single continuous region. Representative examples of polypeptide fragments of the invention; include, for example, fragments comprising, or alternatively consisting of, from about amino acid number 1-20, 21-40,

41-60, 61-80, 81-100, 102-120, 121-140, 141-160, or 161 to the end of the coding region. Moreover, polypeptide fragments can be about 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, or 150 amino acids in length. In this context "about" includes the particularly recited ranges or values, and ranges or values larger or smaller by several (5, 4, 3, 2, or 1) amino acids, at either extreme or at both extremes. Polynucleotides encoding these polypeptides are also encompassed by the invention.

Preferred polypeptide fragments include the secreted protein as well as the mature form. Further preferred polypeptide fragments include the secreted protein or the mature form having a continuous series of deleted residues from the amino or the carboxy terminus, or both. For example, any number of amino acids, ranging from 1-60, can be deleted from the amino terminus of either the secreted polypeptide or the mature form. Similarly, any number of amino acids, ranging from 1-30, can be deleted from the carboxy terminus of the secreted protein or mature form. Furthermore, any combination of the above amino and carboxy terminus deletions are preferred. Similarly, polynucleotides encoding these polypeptide fragments are also preferred.

Also preferred are polypeptide and polynucleotide fragments characterized by structural or functional domains, such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions. Polypeptide fragments of SEQ ID NO:Y falling within conserved domains are specifically contemplated by the present invention. Moreover, polynucleotides encoding these domains are also contemplated.

Other preferred polypeptide fragments are biologically active fragments. Biologically active fragments are those exhibiting activity similar, but not necessarily identical, to an activity of the polypeptide of the present invention. The biological activity of the fragments may include an improved desired activity, or a decreased undesirable activity. Polynucleotides encoding these polypeptide fragments are also encompassed by the invention.

Preferably, the polynucleotide fragments of the invention encode a polypeptide which demonstrates a functional activity. By a polypeptide demonstrating a "functional activity" is meant, a polypeptide capable of displaying one or more known functional activities associated with a full-length (complete) polypeptide of invention protein. Such functional activities include, but are not limited to, biological activity, antigenicity [ability to bind (or compete with a polypeptide of the invention for binding) to an antibody to the polypeptide of the invention], immunogenicity (ability to generate antibody which binds to a polypeptide of the invention), ability to form multimers with polypeptides of the invention, and ability to bind to a receptor or ligand for a polypeptide of the invention.

The functional activity of polypeptides of the invention, and fragments, variants derivatives, and analogs thereof, can be assayed by various methods.

For example, in one embodiment where one is assaying for the ability to bind or compete with full-length polypeptide of the invention for binding to an antibody of the polypeptide of the invention, various immunoassays known in the art can be used, including but not limited to, competitive and non-competitive assay systems using techniques such as radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoradiometric assays, gel diffusion precipitation reactions, immunodiffusion assays, in situ immunoassays (using colloidal gold, enzyme or radioisotope labels, for example), western blots, precipitation reactions, agglutination assays (e.g., gel agglutination assays, hemagglutination assays), complement fixation assays, immunofluorescence assays, protein A assays, and immunoelectrophoresis assays, etc. In one embodiment, antibody binding is detected by detecting a label on the primary antibody. In another embodiment, the primary antibody is detected by detecting binding of a secondary antibody or reagent to the primary antibody. In a further embodiment, the secondary antibody is labeled. Many means are known in the art for detecting binding in an immunoassay and are within the scope of the present invention.

In another embodiment, where a ligand for a polypeptide of the invention identified, or the ability of a polypeptide fragment, variant or derivative of the invention to multimerize is being evaluated, binding can be assayed, e.g., by means well-known in the art, such as, for example, reducing and non-reducing gel

chromatography, protein affinity chromatography, and affinity blotting. See generally, Phizicky, E., et al., 1995, Microbiol. Rev. 59:94-123. In another embodiment, physiological correlates of binding of a polypeptide of the invention to its substrates (signal transduction) can be assayed.

5 In addition, assays described herein (see Examples) and otherwise known in the art may routinely be applied to measure the ability of polypeptides of the invention and fragments, variants derivatives and analogs thereof to elicit related biological activity related to that of the polypeptide of the invention (either in vitro or in vivo). Other methods will be known to the skilled artisan and are within the scope
10 of the invention.

Epitopes and Antibodies

The present invention encompasses polypeptides comprising, or alternatively consisting of, an epitope of the polypeptide having an amino acid sequence of SEQ ID
15 NO:Y, or an epitope of the polypeptide sequence encoded by a polynucleotide sequence contained in ATCC deposit No. Z or encoded by a polynucleotide that hybridizes to the complement of the sequence of SEQ ID NO:X or contained in ATCC deposit No. Z under stringent hybridization conditions or lower stringency hybridization conditions as defined supra. The present invention further encompasses
20 polynucleotide sequences encoding an epitope of a polypeptide sequence of the invention (such as, for example, the sequence disclosed in SEQ ID NO:X), polynucleotide sequences of the complementary strand of a polynucleotide sequence encoding an epitope of the invention, and polynucleotide sequences which hybridize to the complementary strand under stringent hybridization conditions or lower
25 stringency hybridization conditions defined supra.

The term "epitopes," as used herein, refers to portions of a polypeptide having antigenic or immunogenic activity in an animal, preferably a mammal, and most preferably in a human. In a preferred embodiment, the present invention encompasses a polypeptide comprising an epitope, as well as the polynucleotide
30 encoding this polypeptide. An "immunogenic epitope," as used herein, is defined as a portion of a protein that elicits an antibody response in an animal, as determined by any method known in the art, for example, by the methods for generating antibodies

described *infra*. (See, for example, Geysen et al., Proc. Natl. Acad. Sci. USA 81:3998- 4002 (1983)). The term "antigenic epitope," as used herein, is defined as a portion of a protein to which an antibody can immunospecifically bind its antigen as determined by any method well known in the art, for example, by the immunoassays described herein. Immunospecific binding excludes non-specific binding but does not necessarily exclude cross- reactivity with other antigens. Antigenic epitopes need not necessarily be immunogenic.

Fragments which function as epitopes may be produced by any conventional means. (See, e.g., Houghten, Proc. Natl. Acad. Sci. USA 82:5131-5135 (1985), further described in U.S. Patent No. 4,631,211).

In the present invention, antigenic epitopes preferably contain a sequence of at least 4, at least 5, at least 6, at least 7, more preferably at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 20, at least 25, at least 30, at least 40, at least 50, and, most preferably, between about 15 to about 30 amino acids. Preferred polypeptides comprising immunogenic or antigenic epitopes are at least 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 amino acid residues in length. Additional non-exclusive preferred antigenic epitopes include the antigenic epitopes disclosed herein, as well as portions thereof. Antigenic epitopes are useful, for example, to raise antibodies, including monoclonal antibodies, that specifically bind the epitope. Preferred antigenic epitopes include the antigenic epitopes disclosed herein, as well as any combination of two, three, four, five or more of these antigenic epitopes. Antigenic epitopes can be used as the target molecules in immunoassays. (See, for instance, Wilson et al., Cell 37:767-778 (1984); Sutcliffe et al., Science 219:660-666 (1983)).

Similarly, immunogenic epitopes can be used, for example, to induce antibodies according to methods well known in the art. (See, for instance, Sutcliffe et al., *supra*; Wilson et al., *supra*; Chow et al., Proc. Natl. Acad. Sci. USA 82:910-914; and Bittle et al., J. Gen. Virol. 66:2347-2354 (1985)). Preferred immunogenic epitopes include the immunogenic epitopes disclosed herein, as well as any combination of two, three, four, five or more of these immunogenic epitopes. The polypeptides comprising one or more immunogenic epitopes may be presented for eliciting an antibody response together with a carrier protein, such as an albumin, to

an animal system (such as rabbit or mouse), or, if the polypeptide is of sufficient length (at least about 25 amino acids), the polypeptide may be presented without a carrier. However, immunogenic epitopes comprising as few as 8 to 10 amino acids have been shown to be sufficient to raise antibodies capable of binding to, at the very least, linear epitopes in a denatured polypeptide (e.g., in Western blotting).

Epitope-bearing polypeptides of the present invention may be used to induce antibodies according to methods well known in the art including, but not limited to, in vivo immunization, in vitro immunization, and phage display methods. See, e.g., Sutcliffe et al., *supra*; Wilson et al., *supra*, and Bittle et al., *J. Gen. Virol.*, 66:2347-2354 (1985). If in vivo immunization is used, animals may be immunized with free peptide; however, anti-peptide antibody titer may be boosted by coupling the peptide to a macromolecular carrier, such as keyhole limpet hemacyanin (KLH) or tetanus toxoid. For instance, peptides containing cysteine residues may be coupled to a carrier using a linker such as maleimidobenzoyl- N-hydroxysuccinimide ester (MBS), while other peptides may be coupled to carriers using a more general linking agent such as glutaraldehyde. Animals such as rabbits, rats and mice are immunized with either free or carrier- coupled peptides, for instance, by intraperitoneal and/or intradermal injection of emulsions containing about 100 µg of peptide or carrier protein and Freund's adjuvant or any other adjuvant known for stimulating an immune response. Several booster injections may be needed, for instance, at intervals of about two weeks, to provide a useful titer of anti-peptide antibody which can be detected, for example, by ELISA assay using free peptide adsorbed to a solid surface. The titer of anti-peptide antibodies in serum from an immunized animal may be increased by selection of anti-peptide antibodies, for instance, by adsorption to the peptide on a solid support and elution of the selected antibodies according to methods well known in the art.

As one of skill in the art will appreciate, and as discussed above, the polypeptides of the present invention comprising an immunogenic or antigenic epitope can be fused to other polypeptide sequences. For example, the polypeptides of the present invention may be fused with the constant domain of immunoglobulins (IgA, IgE, IgG, IgM), or portions thereof (CH1, CH2, CH3, or any combination thereof and portions thereof), or albumin (including but not limited to recombinant

albumin (see, e.g., U.S. Patent No. 5,876,969, issued March 2, 1999, EP Patent 0 413 622, and U.S. Patent No. 5,766,883, issued June 16, 1998, herein incorporated by reference in their entirety)), resulting in chimeric polypeptides. Such fusion proteins may facilitate purification and may increase half-life in vivo. This has been shown for chimeric proteins consisting of the first two domains of the human CD4-
5 polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins. See, e.g., EP 394,827; Traunecker et al., *Nature*, 331:84-86 (1988). Enhanced delivery of an antigen across the epithelial barrier to the immune system has been demonstrated for antigens (e.g., insulin) conjugated to an
10 FcRn binding partner such as IgG or Fc fragments (see, e.g., PCT Publications WO 96/22024 and WO 99/04813). IgG Fusion proteins that have a disulfide-linked dimeric structure due to the IgG portion disulfide bonds have also been found to be more efficient in binding and neutralizing other molecules than monomeric polypeptides or fragments thereof alone. See, e.g., Fountoulakis et al., *J. Biochem.*,
15 270:3958-3964 (1995). Nucleic acids encoding the above epitopes can also be recombined with a gene of interest as an epitope tag (e.g., the hemagglutinin ("HA") tag or flag tag) to aid in detection and purification of the expressed polypeptide. For example, a system described by Janknecht et al. allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht et al., 1991,
20 *Proc. Natl. Acad. Sci. USA* 88:8972- 897). In this system, the gene of interest is subcloned into a vaccinia recombination plasmid such that the open reading frame of the gene is translationally fused to an amino-terminal tag consisting of six histidine residues. The tag serves as a matrix binding domain for the fusion protein. Extracts from cells infected with the recombinant vaccinia virus are loaded onto Ni²⁺
25 nitriloacetic acid-agarose column and histidine-tagged proteins can be selectively eluted with imidazole-containing buffers.

Additional fusion proteins of the invention may be generated through the techniques of gene-shuffling, motif-shuffling, exon-shuffling, and/or codon-shuffling (collectively referred to as "DNA shuffling"). DNA shuffling may be employed to
30 modulate the activities of polypeptides of the invention, such methods can be used to generate polypeptides with altered activity, as well as agonists and antagonists of the polypeptides. See, generally, U.S. Patent Nos. 5,605,793; 5,811,238; 5,830,721;

5,834,252; and 5,837,458, and Patten et al., Curr. Opinion Biotechnol. 8:724-33 (1997); Harayama, Trends Biotechnol. 16(2):76-82 (1998); Hansson, et al., J. Mol. Biol. 287:265-76 (1999); and Lorenzo and Blasco, Biotechniques 24(2):308- 13 (1998) (each of these patents and publications are hereby incorporated by reference in its entirety). In one embodiment, alteration of polynucleotides corresponding to SEQ ID NO:X and the polypeptides encoded by these polynucleotides may be achieved by DNA shuffling. DNA shuffling involves the assembly of two or more DNA segments by homologous or site-specific recombination to generate variation in the polynucleotide sequence. In another embodiment, polynucleotides of the invention, or the encoded polypeptides, may be altered by being subjected to random mutagenesis by error-prone PCR, random nucleotide insertion or other methods prior to recombination. In another embodiment, one or more components, motifs, sections, parts, domains, fragments, etc., of a polynucleotide encoding a polypeptide of the invention may be recombined with one or more components, motifs, sections, parts, domains, fragments, etc. of one or more heterologous molecules.

Antibodies

Further polypeptides of the invention relate to antibodies and T-cell antigen receptors (TCR) which immunospecifically bind a polypeptide, polypeptide fragment, or variant of SEQ ID NO:Y, and/or an epitope, of the present invention (as determined by immunoassays well known in the art for assaying specific antibody-antigen binding). Antibodies of the invention include, but are not limited to, polyclonal, monoclonal, multispecific, human, humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab') fragments, fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies (including, e.g., anti-Id antibodies to antibodies of the invention), and epitope-binding fragments of any of the above. The term "antibody," as used herein, refers to immunoglobulin molecules and immunologically active portions of immunoglobulin molecules, i.e., molecules that contain an antigen binding site that immunospecifically binds an antigen. The immunoglobulin molecules of the invention can be of any type (e.g., IgG, IgE, IgM, IgD, IgA and IgY), class (e.g., IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2) or subclass of immunoglobulin molecule. In preferred embodiments, the immunoglobulin

molecules of the invention are IgG1. In other preferred embodiments, the immunoglobulin molecules of the invention are IgG4.

Most preferably the antibodies are human antigen-binding antibody fragments of the present invention and include, but are not limited to, Fab, Fab' and F(ab')₂, Fd, single-chain Fvs (scFv), single-chain antibodies, disulfide-linked Fvs (sdFv) and fragments comprising either a VL or VH domain. Antigen-binding antibody fragments, including single-chain antibodies, may comprise the variable region(s) alone or in combination with the entirety or a portion of the following: hinge region, CH1, CH2, and CH3 domains. Also included in the invention are antigen-binding fragments also comprising any combination of variable region(s) with a hinge region, CH1, CH2, and CH3 domains. The antibodies of the invention may be from any animal origin including birds and mammals. Preferably, the antibodies are human, murine (e.g., mouse and rat), donkey, ship rabbit, goat, guinea pig, camel, horse, or chicken. As used herein, "human" antibodies include antibodies having the amino acid sequence of a human immunoglobulin and include antibodies isolated from human immunoglobulin libraries or from animals transgenic for one or more human immunoglobulin and that do not express endogenous immunoglobulins, as described infra and, for example in, U.S. Patent No. 5,939,598 by Kucherlapati et al.

The antibodies of the present invention may be monospecific, bispecific, trispecific or of greater multispecificity. Multispecific antibodies may be specific for different epitopes of a polypeptide of the present invention or may be specific for both a polypeptide of the present invention as well as for a heterologous epitope, such as a heterologous polypeptide or solid support material. See, e.g., PCT publications WO 93/17715; WO 92/08802; WO 91/00360; WO 92/05793; Tutt, et al., J. Immunol. 147:60-69 (1991); U.S. Patent Nos. 4,474,893; 4,714,681; 4,925,648; 5,573,920; 5,601,819; Kostelny et al., J. Immunol. 148:1547-1553 (1992).

Antibodies of the present invention may be described or specified in terms of the epitope(s) or portion(s) of a polypeptide of the present invention which they recognize or specifically bind. The epitope(s) or polypeptide portion(s) may be specified as described herein, e.g., by N-terminal and C-terminal positions, by size in contiguous amino acid residues, or listed in the Tables and Figures. Antibodies which specifically bind any epitope or polypeptide of the present invention may also be

excluded. Therefore, the present invention includes antibodies that specifically bind polypeptides of the present invention, and allows for the exclusion of the same.

Antibodies of the present invention may also be described or specified in terms of their cross-reactivity. Antibodies that do not bind any other analog, ortholog, or homolog of a polypeptide of the present invention are included. Antibodies that bind polypeptides with at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 65%, at least 60%, at least 55%, and at least 50% identity (as calculated using methods known in the art and described herein) to a polypeptide of the present invention are also included in the present invention. In specific embodiments, antibodies of the present invention cross-react with murine, rat and/or rabbit homologs of human proteins and the corresponding epitopes thereof. Antibodies that do not bind polypeptides with less than 95%, less than 90%, less than 85%, less than 80%, less than 75%, less than 70%, less than 65%, less than 60%, less than 55%, and less than 50% identity (as calculated using methods known in the art and described herein) to a polypeptide of the present invention are also included in the present invention. In a specific embodiment, the above-described cross-reactivity is with respect to any single specific antigenic or immunogenic polypeptide, or combination(s) of 2, 3, 4, 5, or more of the specific antigenic and/or immunogenic polypeptides disclosed herein. Further included in the present invention are antibodies which bind polypeptides encoded by polynucleotides which hybridize to a polynucleotide of the present invention under stringent hybridization conditions (as described herein). Antibodies of the present invention may also be described or specified in terms of their binding affinity to a polypeptide of the invention. Preferred binding affinities include those with a dissociation constant or K_d less than 5×10^{-2} M, 10^{-2} M, 5×10^{-3} M, 10^{-3} M, 5×10^{-4} M, 10^{-4} M, 5×10^{-5} M, 10^{-5} M, 5×10^{-6} M, 10^{-6} M, 5×10^{-7} M, 10^{-7} M, 5×10^{-8} M, 10^{-8} M, 5×10^{-9} M, 10^{-9} M, 5×10^{-10} M, 10^{-10} M, 5×10^{-11} M, 10^{-11} M, 5×10^{-12} M, 10^{-12} M, 5×10^{-13} M, 10^{-13} M, 5×10^{-14} M, 10^{-14} M, 5×10^{-15} M, or 10^{-15} M.

The invention also provides antibodies that competitively inhibit binding of an antibody to an epitope of the invention as determined by any method known in the art for determining competitive binding, for example, the immunoassays described herein. In preferred embodiments, the antibody competitively inhibits binding to the

epitope by at least 95%, at least 90%, at least 85 %, at least 80%, at least 75%, at least 70%, at least 60%, or at least 50%.

Antibodies of the present invention may act as agonists or antagonists of the polypeptides of the present invention. For example, the present invention includes
5 antibodies which disrupt the receptor/ligand interactions with the polypeptides of the invention either partially or fully. Preferably, antibodies of the present invention bind an antigenic epitope disclosed herein, or a portion thereof. The invention features both receptor-specific antibodies and ligand-specific antibodies. The invention also features receptor-specific antibodies which do not prevent ligand
10 binding but prevent receptor activation. Receptor activation (i.e., signaling) may be determined by techniques described herein or otherwise known in the art. For example, receptor activation can be determined by detecting the phosphorylation (e.g., tyrosine or serine/threonine) of the receptor or its substrate by immunoprecipitation followed by western blot analysis (for example, as described
15 supra). In specific embodiments, antibodies are provided that inhibit ligand activity or receptor activity by at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, or at least 50% of the activity in absence of the antibody.

The invention also features receptor-specific antibodies which both prevent
20 ligand binding and receptor activation as well as antibodies that recognize the receptor-ligand complex, and, preferably, do not specifically recognize the unbound receptor or the unbound ligand. Likewise, included in the invention are neutralizing antibodies which bind the ligand and prevent binding of the ligand to the receptor, as well as antibodies which bind the ligand, thereby preventing receptor activation, but
25 do not prevent the ligand from binding the receptor. Further included in the invention are antibodies which activate the receptor. These antibodies may act as receptor agonists, i.e., potentiate or activate either all or a subset of the biological activities of the ligand-mediated receptor activation, for example, by inducing dimerization of the receptor. The antibodies may be specified as agonists, antagonists or inverse agonists
30 for biological activities comprising the specific biological activities of the peptides of the invention disclosed herein. The above antibody agonists can be made using methods known in the art. See, e.g., PCT publication WO 96/40281; U.S. Patent No.

5,811,097; Deng et al., Blood 92(6):1981-1988 (1998); Chen et al., Cancer Res. 58(16):3668-3678 (1998); Harrop et al., J. Immunol. 161(4):1786-1794 (1998); Zhu et al., Cancer Res. 58(15):3209-3214 (1998); Yoon et al., J. Immunol. 160(7):3170-3179 (1998); Prat et al., J. Cell. Sci. 111(Pt2):237-247 (1998); Pitard et al., J. Immunol. Methods 205(2):177-190 (1997); Liautard et al., Cytokine 9(4):233-241 (1997); Carlson et al., J. Biol. Chem. 272(17):11295-11301 (1997); Taryman et al., Neuron 14(4):755-762 (1995); Muller et al., Structure 6(9):1153-1167 (1998); Bartunek et al., Cytokine 8(1):14-20 (1996) (which are all incorporated by reference herein in their entireties).

10 Antibodies of the present invention may be used, for example, but not limited to, to purify, detect, and target the polypeptides of the present invention, including both in vitro and in vivo diagnostic and therapeutic methods. For example, the antibodies have use in immunoassays for qualitatively and quantitatively measuring levels of the polypeptides of the present invention in biological samples. See, e.g.,
15 Harlow et al., Antibodies: A Laboratory Manual, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988) (incorporated by reference herein in its entirety).

 As discussed in more detail below, the antibodies of the present invention may be used either alone or in combination with other compositions. The antibodies may further be recombinantly fused to a heterologous polypeptide at the N- or C-terminus
20 or chemically conjugated (including covalently and non-covalently conjugations) to polypeptides or other compositions. For example, antibodies of the present invention may be recombinantly fused or conjugated to molecules useful as labels in detection assays and effector molecules such as heterologous polypeptides, drugs, radionuclides, or toxins. See, e.g., PCT publications WO 92/08495; WO 91/14438;
25 WO 89/12624; U.S. Patent No. 5,314,995; and EP 396,387.

 The antibodies of the invention include derivatives that are modified, i.e., by the covalent attachment of any type of molecule to the antibody such that covalent attachment does not prevent the antibody from generating an anti-idiotypic response. For example, but not by way of limitation, the antibody derivatives include
30 antibodies that have been modified, e.g., by glycosylation, acetylation, pegylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to a cellular ligand or other protein, etc. Any of

numerous chemical modifications may be carried out by known techniques, including, but not limited to specific chemical cleavage, acetylation, formylation, metabolic synthesis of tunicamycin, etc. Additionally, the derivative may contain one or more non-classical amino acids.

5 The antibodies of the present invention may be generated by any suitable method known in the art. Polyclonal antibodies to an antigen-of- interest can be produced by various procedures well known in the art. For example, a polypeptide of the invention can be administered to various host animals including, but not limited to, rabbits, mice, rats, etc. to induce the production of sera containing polyclonal
10 antibodies specific for the antigen. Various adjuvants may be used to increase the immunological response, depending on the host species, and include but are not limited to, Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanins, dinitrophenol, and
15 potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and corynebacterium parvum. Such adjuvants are also well known in the art.

Monoclonal antibodies can be prepared using a wide variety of techniques known in the art including the use of hybridoma, recombinant, and phage display technologies, or a combination thereof. For example, monoclonal antibodies can be
20 produced using hybridoma techniques including those known in the art and taught, for example, in Harlow et al., *Antibodies: A Laboratory Manual*, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988); Hammerling, et al., in: *Monoclonal Antibodies and T-Cell Hybridomas* 563-681 (Elsevier, N.Y., 1981) (said references incorporated by reference in their entireties). The term "monoclonal antibody" as used herein is not
25 limited to antibodies produced through hybridoma technology. The term "monoclonal antibody" refers to an antibody that is derived from a single clone, including any eukaryotic, prokaryotic, or phage clone, and not the method by which it is produced.

Methods for producing and screening for specific antibodies using hybridoma
30 technology are routine and well known in the art and are discussed in detail in the Examples (e.g., Example 16). In a non-limiting example, mice can be immunized with a polypeptide of the invention or a cell expressing such peptide. Once an

immune response is detected, e.g., antibodies specific for the antigen are detected in the mouse serum, the mouse spleen is harvested and splenocytes isolated. The splenocytes are then fused by well known techniques to any suitable myeloma cells, for example cells from cell line SP20 available from the ATCC. Hybridomas are
5 selected and cloned by limited dilution. The hybridoma clones are then assayed by methods known in the art for cells that secrete antibodies capable of binding a polypeptide of the invention. Ascites fluid, which generally contains high levels of antibodies, can be generated by immunizing mice with positive hybridoma clones.

Accordingly, the present invention provides methods of generating
10 monoclonal antibodies as well as antibodies produced by the method comprising culturing a hybridoma cell secreting an antibody of the invention wherein, preferably, the hybridoma is generated by fusing splenocytes isolated from a mouse immunized with an antigen of the invention with myeloma cells and then screening the hybridomas resulting from the fusion for hybridoma clones that secrete an antibody
15 able to bind a polypeptide of the invention.

Antibody fragments which recognize specific epitopes may be generated by known techniques. For example, Fab and F(ab')₂ fragments of the invention may be produced by proteolytic cleavage of immunoglobulin molecules, using enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab')₂ fragments).
20 F(ab')₂ fragments contain the variable region, the light chain constant region and the CH1 domain of the heavy chain.

For example, the antibodies of the present invention can also be generated using various phage display methods known in the art. In phage display methods, functional antibody domains are displayed on the surface of phage particles which
25 carry the polynucleotide sequences encoding them. In a particular embodiment, such phage can be utilized to display antigen binding domains expressed from a repertoire or combinatorial antibody library (e.g., human or murine). Phage expressing an antigen binding domain that binds the antigen of interest can be selected or identified with antigen, e.g., using labeled antigen or antigen bound or captured to a solid
30 surface or bead. Phage used in these methods are typically filamentous phage including fd and M13 binding domains expressed from phage with Fab, Fv or disulfide stabilized Fv antibody domains recombinantly fused to either the phage

gene III or gene VIII protein. Examples of phage display methods that can be used to make the antibodies of the present invention include those disclosed in Brinkman et al., J. Immunol. Methods 182:41-50 (1995); Ames et al., J. Immunol. Methods 184:177-186 (1995); Kettleborough et al., Eur. J. Immunol. 24:952-958 (1994); Persic
5 et al., Gene 187 9-18 (1997); Burton et al., Advances in Immunology 57:191-280 (1994); PCT application No. PCT/GB91/01134; PCT publications WO 90/02809; WO 91/10737; WO 92/01047; WO 92/18619; WO 93/11236; WO 95/15982; WO 95/20401; and U.S. Patent Nos. 5,698,426; 5,223,409; 5,403,484; 5,580,717; 5,427,908; 5,750,753; 5,821,047; 5,571,698; 5,427,908; 5,516,637; 5,780,225;
10 5,658,727; 5,733,743 and 5,969,108; each of which is incorporated herein by reference in its entirety.

As described in the above references, after phage selection, the antibody coding regions from the phage can be isolated and used to generate whole antibodies, including human antibodies, or any other desired antigen binding fragment, and
15 expressed in any desired host, including mammalian cells, insect cells, plant cells, yeast, and bacteria, e.g., as described in detail below. For example, techniques to recombinantly produce Fab, Fab' and F(ab')₂ fragments can also be employed using methods known in the art such as those disclosed in PCT publication WO 92/22324; Mullinax et al., BioTechniques 12(6):864-869 (1992); and Sawai et al., AJRI 34:26-
20 34 (1995); and Better et al., Science 240:1041-1043 (1988) (said references incorporated by reference in their entireties).

Examples of techniques which can be used to produce single-chain Fvs and antibodies include those described in U.S. Patents 4,946,778 and 5,258,498; Huston et al., Methods in Enzymology 203:46-88 (1991); Shu et al., PNAS 90:7995-7999
25 (1993); and Skerra et al., Science 240:1038-1040 (1988). For some uses, including in vivo use of antibodies in humans and in vitro detection assays, it may be preferable to use chimeric, humanized, or human antibodies. A chimeric antibody is a molecule in which different portions of the antibody are derived from different animal species, such as antibodies having a variable region derived from a murine monoclonal
30 antibody and a human immunoglobulin constant region. Methods for producing chimeric antibodies are known in the art. See e.g., Morrison, Science 229:1202 (1985); Oi et al., BioTechniques 4:214 (1986); Gillies et al., (1989) J. Immunol.

Methods 125:191-202; U.S. Patent Nos. 5,807,715; 4,816,567; and 4,816,397, which are incorporated herein by reference in their entirety. Humanized antibodies are antibody molecules from non-human species antibody that binds the desired antigen having one or more complementarity determining regions (CDRs) from the non-human species and a framework regions from a human immunoglobulin molecule. Often, framework residues in the human framework regions will be substituted with the corresponding residue from the CDR donor antibody to alter, preferably improve, antigen binding. These framework substitutions are identified by methods well known in the art, e.g., by modeling of the interactions of the CDR and framework residues to identify framework residues important for antigen binding and sequence comparison to identify unusual framework residues at particular positions. (See, e.g., Queen et al., U.S. Patent No. 5,585,089; Riechmann et al., Nature 332:323 (1988), which are incorporated herein by reference in their entirety.) Antibodies can be humanized using a variety of techniques known in the art including, for example, CDR-grafting (EP 239,400; PCT publication WO 91/09967; U.S. Patent Nos. 5,225,539; 5,530,101; and 5,585,089), veneering or resurfacing (EP 592,106; EP 519,596; Padlan, Molecular Immunology 28(4/5):489-498 (1991); Studnicka et al., Protein Engineering 7(6):805-814 (1994); Roguska. et al., PNAS 91:969-973 (1994)), and chain shuffling (U.S. Patent No. 5,565,332).

Completely human antibodies are particularly desirable for therapeutic treatment of human patients. Human antibodies can be made by a variety of methods known in the art including phage display methods described above using antibody libraries derived from human immunoglobulin sequences. See also, U.S. Patent Nos. 4,444,887 and 4,716,111; and PCT publications WO 98/46645, WO 98/50433, WO 98/24893, WO 98/16654, WO 96/34096, WO 96/33735, and WO 91/10741; each of which is incorporated herein by reference in its entirety.

Human antibodies can also be produced using transgenic mice which are incapable of expressing functional endogenous immunoglobulins, but which can express human immunoglobulin genes. For example, the human heavy and light chain immunoglobulin gene complexes may be introduced randomly or by homologous recombination into mouse embryonic stem cells. Alternatively, the human variable region, constant region, and diversity region may be introduced into

mouse embryonic stem cells in addition to the human heavy and light chain genes. The mouse heavy and light chain immunoglobulin genes may be rendered non-functional separately or simultaneously with the introduction of human immunoglobulin loci by homologous recombination. In particular, homozygous deletion of the JH region prevents endogenous antibody production. The modified embryonic stem cells are expanded and microinjected into blastocysts to produce chimeric mice. The chimeric mice are then bred to produce homozygous offspring which express human antibodies. The transgenic mice are immunized in the normal fashion with a selected antigen, e.g., all or a portion of a polypeptide of the invention. Monoclonal antibodies directed against the antigen can be obtained from the immunized, transgenic mice using conventional hybridoma technology. The human immunoglobulin transgenes harbored by the transgenic mice rearrange during B cell differentiation, and subsequently undergo class switching and somatic mutation. Thus, using such a technique, it is possible to produce therapeutically useful IgG, IgA, IgM and IgE antibodies. For an overview of this technology for producing human antibodies, see Lonberg and Huszar, *Int. Rev. Immunol.* 13:65-93 (1995). For a detailed discussion of this technology for producing human antibodies and human monoclonal antibodies and protocols for producing such antibodies, see, e.g., PCT publications WO 98/24893; WO 92/01047; WO 96/34096; WO 96/33735; European Patent No. 0 598 877; U.S. Patent Nos. 5,413,923; 5,625,126; 5,633,425; 5,569,825; 5,661,016; 5,545,806; 5,814,318; 5,885,793; 5,916,771; and 5,939,598, which are incorporated by reference herein in their entirety. In addition, companies such as Abgenix, Inc. (Freemont, CA) and Genpharm (San Jose, CA) can be engaged to provide human antibodies directed against a selected antigen using technology similar to that described above.

Completely human antibodies which recognize a selected epitope can be generated using a technique referred to as "guided selection." In this approach a selected non-human monoclonal antibody, e.g., a mouse antibody, is used to guide the selection of a completely human antibody recognizing the same epitope. (Jespers et al., *Bio/technology* 12:899-903 (1988)).

Further, antibodies to the polypeptides of the invention can, in turn, be utilized to generate anti-idiotypic antibodies that "mimic" polypeptides of the invention using

techniques well known to those skilled in the art. (See, e.g., Greenspan & Bona, FASEB J. 7(5):437-444; (1989) and Nissinoff, J. Immunol. 147(8):2429-2438 (1991)). For example, antibodies which bind to and competitively inhibit polypeptide multimerization and/or binding of a polypeptide of the invention to a ligand can be used to generate anti-idiotypes that "mimic" the polypeptide multimerization and/or binding domain and, as a consequence, bind to and neutralize polypeptide and/or its ligand. Such neutralizing anti-idiotypes or Fab fragments of such anti-idiotypes can be used in therapeutic regimens to neutralize polypeptide ligand. For example, such anti-idiotypic antibodies can be used to bind a polypeptide of the invention and/or to bind its ligands/receptors, and thereby block its biological activity.

Polynucleotides Encoding Antibodies

The invention further provides polynucleotides comprising a nucleotide sequence encoding an antibody of the invention and fragments thereof. The invention also encompasses polynucleotides that hybridize under stringent or lower stringency hybridization conditions, e.g., as defined supra, to polynucleotides that encode an antibody, preferably, that specifically binds to a polypeptide of the invention, preferably, an antibody that binds to a polypeptide having the amino acid sequence of SEQ ID NO:Y.

The polynucleotides may be obtained, and the nucleotide sequence of the polynucleotides determined, by any method known in the art. For example, if the nucleotide sequence of the antibody is known, a polynucleotide encoding the antibody may be assembled from chemically synthesized oligonucleotides (e.g., as described in Kutmeier et al., BioTechniques 17:242 (1994)), which, briefly, involves the synthesis of overlapping oligonucleotides containing portions of the sequence encoding the antibody, annealing and ligating of those oligonucleotides, and then amplification of the ligated oligonucleotides by PCR.

Alternatively, a polynucleotide encoding an antibody may be generated from nucleic acid from a suitable source. If a clone containing a nucleic acid encoding a particular antibody is not available, but the sequence of the antibody molecule is known, a nucleic acid encoding the immunoglobulin may be chemically synthesized or obtained from a suitable source (e.g., an antibody cDNA library, or a cDNA library

generated from, or nucleic acid, preferably poly A+ RNA, isolated from, any tissue or cells expressing the antibody, such as hybridoma cells selected to express an antibody of the invention) by PCR amplification using synthetic primers hybridizable to the 3' and 5' ends of the sequence or by cloning using an oligonucleotide probe specific for the particular gene sequence to identify, e.g., a cDNA clone from a cDNA library that encodes the antibody. Amplified nucleic acids generated by PCR may then be cloned into replicable cloning vectors using any method well known in the art.

Once the nucleotide sequence and corresponding amino acid sequence of the antibody is determined, the nucleotide sequence of the antibody may be manipulated using methods well known in the art for the manipulation of nucleotide sequences, e.g., recombinant DNA techniques, site directed mutagenesis, PCR, etc. (see, for example, the techniques described in Sambrook et al., 1990, Molecular Cloning, A Laboratory Manual, 2d Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY and Ausubel et al., eds., 1998, Current Protocols in Molecular Biology, John Wiley & Sons, NY, which are both incorporated by reference herein in their entireties), to generate antibodies having a different amino acid sequence, for example to create amino acid substitutions, deletions, and/or insertions.

In a specific embodiment, the amino acid sequence of the heavy and/or light chain variable domains may be inspected to identify the sequences of the complementarity determining regions (CDRs) by methods that are well known in the art, e.g., by comparison to known amino acid sequences of other heavy and light chain variable regions to determine the regions of sequence hypervariability. Using routine recombinant DNA techniques, one or more of the CDRs may be inserted within framework regions, e.g., into human framework regions to humanize a non-human antibody, as described supra. The framework regions may be naturally occurring or consensus framework regions, and preferably human framework regions (see, e.g., Chothia et al., J. Mol. Biol. 278: 457-479 (1998) for a listing of human framework regions). Preferably, the polynucleotide generated by the combination of the framework regions and CDRs encodes an antibody that specifically binds a polypeptide of the invention. Preferably, as discussed supra, one or more amino acid substitutions may be made within the framework regions, and, preferably, the amino

acid substitutions improve binding of the antibody to its antigen. Additionally, such methods may be used to make amino acid substitutions or deletions of one or more variable region cysteine residues participating in an intrachain disulfide bond to generate antibody molecules lacking one or more intrachain disulfide bonds. Other alterations to the polynucleotide are encompassed by the present invention and within the skill of the art.

In addition, techniques developed for the production of "chimeric antibodies" (Morrison et al., Proc. Natl. Acad. Sci. 81:851-855 (1984); Neuberger et al., Nature 312:604-608 (1984); Takeda et al., Nature 314:452-454 (1985)) by splicing genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity can be used. As described supra, a chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived from a murine mAb and a human immunoglobulin constant region, e.g., humanized antibodies.

Alternatively, techniques described for the production of single chain antibodies (U.S. Patent No. 4,946,778; Bird, Science 242:423-42 (1988); Huston et al., Proc. Natl. Acad. Sci. USA 85:5879-5883 (1988); and Ward et al., Nature 334:544-54 (1989)) can be adapted to produce single chain antibodies. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region via an amino acid bridge, resulting in a single chain polypeptide. Techniques for the assembly of functional Fv fragments in E. coli may also be used (Skerra et al., Science 242:1038-1041 (1988)).

Methods of Producing Antibodies

The antibodies of the invention can be produced by any method known in the art for the synthesis of antibodies, in particular, by chemical synthesis or preferably, by recombinant expression techniques.

Recombinant expression of an antibody of the invention, or fragment, derivative or analog thereof, (e.g., a heavy or light chain of an antibody of the invention or a single chain antibody of the invention), requires construction of an expression vector containing a polynucleotide that encodes the antibody. Once a

polynucleotide encoding an antibody molecule or a heavy or light chain of an antibody, or portion thereof (preferably containing the heavy or light chain variable domain), of the invention has been obtained, the vector for the production of the antibody molecule may be produced by recombinant DNA technology using techniques well known in the art. Thus, methods for preparing a protein by expressing a polynucleotide containing an antibody encoding nucleotide sequence are described herein. Methods which are well known to those skilled in the art can be used to construct expression vectors containing antibody coding sequences and appropriate transcriptional and translational control signals. These methods include, for example, in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. The invention, thus, provides replicable vectors comprising a nucleotide sequence encoding an antibody molecule of the invention, or a heavy or light chain thereof, or a heavy or light chain variable domain, operably linked to a promoter. Such vectors may include the nucleotide sequence encoding the constant region of the antibody molecule (see, e.g., PCT Publication WO 86/05807; PCT Publication WO 89/01036; and U.S. Patent No. 5,122,464) and the variable domain of the antibody may be cloned into such a vector for expression of the entire heavy or light chain.

The expression vector is transferred to a host cell by conventional techniques and the transfected cells are then cultured by conventional techniques to produce an antibody of the invention. Thus, the invention includes host cells containing a polynucleotide encoding an antibody of the invention, or a heavy or light chain thereof, or a single chain antibody of the invention, operably linked to a heterologous promoter. In preferred embodiments for the expression of double-chained antibodies, vectors encoding both the heavy and light chains may be co-expressed in the host cell for expression of the entire immunoglobulin molecule, as detailed below.

A variety of host-expression vector systems may be utilized to express the antibody molecules of the invention. Such host-expression systems represent vehicles by which the coding sequences of interest may be produced and subsequently purified, but also represent cells which may, when transformed or transfected with the appropriate nucleotide coding sequences, express an antibody molecule of the invention in situ. These include but are not limited to microorganisms such as

bacteria (e.g., *E. coli*, *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing antibody coding sequences; yeast (e.g., *Saccharomyces*, *Pichia*) transformed with recombinant yeast expression vectors containing antibody coding sequences; insect cell systems
5 infected with recombinant virus expression vectors (e.g., baculovirus) containing antibody coding sequences; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing antibody coding sequences; or mammalian cell systems (e.g., COS, CHO,
10 BHK, 293, 3T3 cells) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter). Preferably, bacterial cells such as *Escherichia coli*, and more preferably, eukaryotic cells, especially for the expression of whole
15 recombinant antibody molecule, are used for the expression of a recombinant antibody molecule. For example, mammalian cells such as Chinese hamster ovary cells (CHO), in conjunction with a vector such as the major intermediate early gene promoter element from human cytomegalovirus is an effective expression system for antibodies (Foecking et al., *Gene* 45:101 (1986); Cockett et al., *Bio/Technology* 8:2
20 (1990)).

In bacterial systems, a number of expression vectors may be advantageously selected depending upon the use intended for the antibody molecule being expressed. For example, when a large quantity of such a protein is to be produced, for the generation of pharmaceutical compositions of an antibody molecule, vectors which
25 direct the expression of high levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not limited, to the *E. coli* expression vector pUR278 (Ruther et al., *EMBO J.* 2:1791 (1983)), in which the antibody coding sequence may be ligated individually into the vector in frame with the lac Z coding region so that a fusion protein is produced; pIN vectors (Inouye & Inouye, *Nucleic
30 Acids Res.* 13:3101-3109 (1985); Van Heeke & Schuster, *J. Biol. Chem.* 24:5503-5509 (1989)); and the like. pGEX vectors may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such

fusion proteins are soluble and can easily be purified from lysed cells by adsorption and binding to matrix glutathione-agarose beads followed by elution in the presence of free glutathione. The pGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene product can be released from the GST moiety.

In an insect system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in *Spodoptera frugiperda* cells. The antibody coding sequence may be cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter).

In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, the antibody coding sequence of interest may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by in vitro or in vivo recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing the antibody molecule in infected hosts. (e.g., see Logan & Shenk, Proc. Natl. Acad. Sci. USA 81:355-359 (1984)). Specific initiation signals may also be required for efficient translation of inserted antibody coding sequences. These signals include the ATG initiation codon and adjacent sequences. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (see Bittner et al., Methods in Enzymol. 153:51-544 (1987)).

In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein.

Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and gene products. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end, eukaryotic host cells which
5 possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such mammalian host cells include but are not limited to CHO, VERO, BHK, HeLa, COS, MDCK, 293, 3T3, WI38, and in particular, breast cancer cell lines such as, for example, BT483, Hs578T, HTB2, BT20 and T47D, and normal mammary gland cell
10 line such as, for example, CRL7030 and Hs578Bst.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express the antibody molecule may be engineered. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with DNA controlled by
15 appropriate expression control elements (e.g., promoter, enhancer, sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection
20 and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines which express the antibody molecule. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that interact directly or indirectly with the antibody molecule.

25 A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler et al., Cell 11:223 (1977)), hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, Proc. Natl. Acad. Sci. USA 48:202 (1992)), and adenine phosphoribosyltransferase (Lowy et al., Cell 22:817 (1980)) genes can be employed in tk-, hgp^rt- or ap^rt- cells, respectively.
30 Also, antimetabolite resistance can be used as the basis of selection for the following genes: dhfr, which confers resistance to methotrexate (Wigler et al., Natl. Acad. Sci. USA 77:357 (1980); O'Hare et al., Proc. Natl. Acad. Sci. USA 78:1527 (1981)); gpt,

which confers resistance to mycophenolic acid (Mulligan & Berg, Proc. Natl. Acad. Sci. USA 78:2072 (1981)); neo, which confers resistance to the aminoglycoside G-418 Clinical Pharmacy 12:488-505; Wu and Wu, Biotherapy 3:87-95 (1991); Tolstoshev, Ann. Rev. Pharmacol. Toxicol. 32:573-596 (1993); Mulligan, Science 5 260:926-932 (1993); and Morgan and Anderson, Ann. Rev. Biochem. 62:191-217 (1993); May, 1993, TIB TECH 11(5):155-215; and hygromycin (Santerre et al., Gene 30:147 (1984)). Methods commonly known in the art of recombinant DNA technology may be routinely applied to select the desired recombinant clone, and such methods are described, for example, in Ausubel et al. 10 (eds.), Current Protocols in Molecular Biology, John Wiley & Sons, NY (1993); Kriegler, Gene Transfer and Expression, A Laboratory Manual, Stockton Press, NY (1990); and in Chapters 12 and 13, Dracopoli et al. (eds), Current Protocols in Human Genetics, John Wiley & Sons, NY (1994); Colberre-Garapin et al., J. Mol. Biol. 150:1 (1981), which are incorporated by reference herein in their entireties.

15 The expression levels of an antibody molecule can be increased by vector amplification (for a review, see Bebbington and Hentschel, The use of vectors based on gene amplification for the expression of cloned genes in mammalian cells in DNA cloning, Vol.3. (Academic Press, New York, 1987)). When a marker in the vector system expressing antibody is amplifiable, increase in the level of inhibitor present in 20 culture of host cell will increase the number of copies of the marker gene. Since the amplified region is associated with the antibody gene, production of the antibody will also increase (Crouse et al., Mol. Cell. Biol. 3:257 (1983)).

The host cell may be co-transfected with two expression vectors of the invention, the first vector encoding a heavy chain derived polypeptide and the second 25 vector encoding a light chain derived polypeptide. The two vectors may contain identical selectable markers which enable equal expression of heavy and light chain polypeptides. Alternatively, a single vector may be used which encodes, and is capable of expressing, both heavy and light chain polypeptides. In such situations, the light chain should be placed before the heavy chain to avoid an excess of toxic 30 free heavy chain (Proudfoot, Nature 322:52 (1986); Kohler, Proc. Natl. Acad. Sci. USA 77:2197 (1980)). The coding sequences for the heavy and light chains may comprise cDNA or genomic DNA.

Once an antibody molecule of the invention has been produced by an animal, chemically synthesized, or recombinantly expressed, it may be purified by any method known in the art for purification of an immunoglobulin molecule, for example, by chromatography (e.g., ion exchange, affinity, particularly by affinity for
5 the specific antigen after Protein A, and sizing column chromatography), centrifugation, differential solubility, or by any other standard technique for the purification of proteins. In addition, the antibodies of the present invention or fragments thereof can be fused to heterologous polypeptide sequences described herein or otherwise known in the art, to facilitate purification.

10 The present invention encompasses antibodies recombinantly fused or chemically conjugated (including both covalently and non-covalently conjugations) to a polypeptide (or portion thereof, preferably at least 10, 20, 30, 40, 50, 60, 70, 80, 90 or 100 amino acids of the polypeptide) of the present invention to generate fusion proteins. The fusion does not necessarily need to be direct, but may occur through
15 linker sequences. The antibodies may be specific for antigens other than polypeptides (or portion thereof, preferably at least 10, 20, 30, 40, 50, 60, 70, 80, 90 or 100 amino acids of the polypeptide) of the present invention. For example, antibodies may be used to target the polypeptides of the present invention to particular cell types, either in vitro or in vivo, by fusing or conjugating the polypeptides of the present invention
20 to antibodies specific for particular cell surface receptors. Antibodies fused or conjugated to the polypeptides of the present invention may also be used in in vitro immunoassays and purification methods using methods known in the art. See e.g., Harbor et al., supra, and PCT publication WO 93/21232; EP 439,095; Naramura et al., Immunol. Lett. 39:91-99 (1994); U.S. Patent 5,474,981; Gillies et al., PNAS
25 89:1428-1432 (1992); Fell et al., J. Immunol. 146:2446-2452(1991), which are incorporated by reference in their entirety.

The present invention further includes compositions comprising the polypeptides of the present invention fused or conjugated to antibody domains other than the variable regions. For example, the polypeptides of the present invention may
30 be fused or conjugated to an antibody Fc region, or portion thereof. The antibody portion fused to a polypeptide of the present invention may comprise the constant region, hinge region, CH1 domain, CH2 domain, and CH3 domain or any

combination of whole domains or portions thereof. The polypeptides may also be fused or conjugated to the above antibody portions to form multimers. For example, Fc portions fused to the polypeptides of the present invention can form dimers through disulfide bonding between the Fc portions. Higher multimeric forms can be made by fusing the polypeptides to portions of IgA and IgM. Methods for fusing or conjugating the polypeptides of the present invention to antibody portions are known in the art. See, e.g., U.S. Patent Nos. 5,336,603; 5,622,929; 5,359,046; 5,349,053; 5,447,851; 5,112,946; EP 307,434; EP 367,166; PCT publications WO 96/04388; WO 91/06570; Ashkenazi et al., Proc. Natl. Acad. Sci. USA 88:10535-10539 (1991); Zheng et al., J. Immunol. 154:5590-5600 (1995); and Vil et al., Proc. Natl. Acad. Sci. USA 89:11337-11341(1992) (said references incorporated by reference in their entireties).

As discussed, supra, the polypeptides corresponding to a polypeptide, polypeptide fragment, or a variant of SEQ ID NO:Y may be fused or conjugated to the above antibody portions to increase the in vivo half life of the polypeptides or for use in immunoassays using methods known in the art. Further, the polypeptides corresponding to SEQ ID NO:Y may be fused or conjugated to the above antibody portions to facilitate purification. One reported example describes chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins. (EP 394,827; Traunecker et al., Nature 331:84-86 (1988). The polypeptides of the present invention fused or conjugated to an antibody having disulfide- linked dimeric structures (due to the IgG) may also be more efficient in binding and neutralizing other molecules, than the monomeric secreted protein or protein fragment alone. (Fountoulakis et al., J. Biochem. 270:3958-3964 (1995)). In many cases, the Fc part in a fusion protein is beneficial in therapy and diagnosis, and thus can result in, for example, improved pharmacokinetic properties. (EP A 232,262). Alternatively, deleting the Fc part after the fusion protein has been expressed, detected, and purified, would be desired. For example, the Fc portion may hinder therapy and diagnosis if the fusion protein is used as an antigen for immunizations. In drug discovery, for example, human proteins, such as hIL-5, have been fused with Fc portions for the purpose of high-throughput screening assays to

identify antagonists of hIL-5. (See, Bennett et al., J. Molecular Recognition 8:52-58 (1995); Johanson et al., J. Biol. Chem. 270:9459-9471 (1995).

Moreover, the antibodies or fragments thereof of the present invention can be fused to marker sequences, such as a peptide to facilitate purification. In preferred
5 embodiments, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311), among others, many of which are commercially available. As described in Gentz et al., Proc. Natl. Acad. Sci. USA 86:821-824 (1989), for instance, hexa-histidine provides for convenient purification of the fusion protein. Other peptide tags
10 useful for purification include, but are not limited to, the "HA" tag, which corresponds to an epitope derived from the influenza hemagglutinin protein (Wilson et al., Cell 37:767 (1984)) and the "flag" tag.

The present invention further encompasses antibodies or fragments thereof conjugated to a diagnostic or therapeutic agent. The antibodies can be used
15 diagnostically to, for example, monitor the development or progression of a tumor as part of a clinical testing procedure to, e.g., determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent
20 materials, radioactive materials, positron emitting metals using various positron emission tomographies, and nonradioactive paramagnetic metal ions. The detectable substance may be coupled or conjugated either directly to the antibody (or fragment thereof) or indirectly, through an intermediate (such as, for example, a linker known in the art) using techniques known in the art. See, for example, U.S. Patent No.
25 4,741,900 for metal ions which can be conjugated to antibodies for use as diagnostics according to the present invention. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, beta-galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone,
30 fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin,

and aequorin; and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{111}In or ^{99}Tc .

Further, an antibody or fragment thereof may be conjugated to a therapeutic moiety such as a cytotoxin, e.g., a cytostatic or cytotoxic agent, a therapeutic agent or a radioactive metal ion, e.g., alpha-emitters such as, for example, ^{213}Bi . A cytotoxin or cytotoxic agent includes any agent that is detrimental to cells. Examples include paclitaxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, teniposide, vincristine, vinblastine, colchicine, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof. Therapeutic agents include, but are not limited to, antimetabolites (e.g., methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (e.g., mechlorethamine, thioepa chlorambucil, melphalan, carmustine (BSNU) and lomustine (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cis-dichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (e.g., daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (e.g., dactinomycin (formerly actinomycin), bleomycin, mithramycin, and anthramycin (AMC)), and anti-mitotic agents (e.g., vincristine and vinblastine).

The conjugates of the invention can be used for modifying a given biological response; the therapeutic agent or drug moiety is not to be construed as limited to classical chemical therapeutic agents. For example, the drug moiety may be a protein or polypeptide possessing a desired biological activity. Such proteins may include, for example, a toxin such as abrin, ricin A, pseudomonas exotoxin, or diphtheria toxin; a protein such as tumor necrosis factor, α -interferon, β -interferon, nerve growth factor, platelet derived growth factor, tissue plasminogen activator, an apoptotic agent, e.g., TNF- α , TNF- β , AIM I (See, International Publication No. WO 97/33899), AIM II (See, International Publication No. WO 97/34911), Fas Ligand (Takahashi *et al.*, *Int. Immunol.*, 6:1567-1574 (1994)), VEGI (See, International Publication No. WO 99/23105), a thrombotic agent or an anti-angiogenic agent, e.g., angiostatin or endostatin; or, biological response modifiers such as, for example, lymphokines, interleukin-1 ("IL-1"), interleukin-2 ("IL-2"), interleukin-6 ("IL-6"),

granulocyte macrophage colony stimulating factor ("GM-CSF"), granulocyte colony stimulating factor ("G-CSF"), or other growth factors.

Antibodies may also be attached to solid supports, which are particularly useful for immunoassays or purification of the target antigen. Such solid supports
5 include, but are not limited to, glass, cellulose, polyacrylamide, nylon, polystyrene, polyvinyl chloride or polypropylene.

Techniques for conjugating such therapeutic moiety to antibodies are well known, see, e.g., Arnon et al., "Monoclonal Antibodies For Immunotargeting Of Drugs In Cancer Therapy", in Monoclonal Antibodies And Cancer Therapy, Reisfeld et al. (eds.), pp. 243-56 (Alan R. Liss, Inc. 1985); Hellstrom et al., "Antibodies For Drug Delivery", in Controlled Drug Delivery (2nd Ed.), Robinson et al. (eds.), pp. 623-53 (Marcel Dekker, Inc. 1987); Thorpe, "Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review", in Monoclonal Antibodies '84: Biological And Clinical Applications, Pinchera et al. (eds.), pp. 475-506 (1985); "Analysis, Results,
15 And Future Prospective Of The Therapeutic Use Of Radiolabeled Antibody In Cancer Therapy", in Monoclonal Antibodies For Cancer Detection And Therapy, Baldwin et al. (eds.), pp. 303-16 (Academic Press 1985), and Thorpe et al., "The Preparation And Cytotoxic Properties Of Antibody-Toxin Conjugates", Immunol. Rev. 62:119-58 (1982).

20 Alternatively, an antibody can be conjugated to a second antibody to form an antibody heteroconjugate as described by Segal in U.S. Patent No. 4,676,980, which is incorporated herein by reference in its entirety.

An antibody, with or without a therapeutic moiety conjugated to it, administered alone or in combination with cytotoxic factor(s) and/or cytokine(s) can
25 be used as a therapeutic.

Immunophenotyping

The antibodies of the invention may be utilized for immunophenotyping of cell lines and biological samples. The translation product of the gene of the present
30 invention may be useful as a cell specific marker, or more specifically as a cellular marker that is differentially expressed at various stages of differentiation and/or maturation of particular cell types. Monoclonal antibodies directed against a specific

epitope, or combination of epitopes, will allow for the screening of cellular populations expressing the marker. Various techniques can be utilized using monoclonal antibodies to screen for cellular populations expressing the marker(s), and include magnetic separation using antibody-coated magnetic beads, "panning" with antibody attached to a solid matrix (i.e., plate), and flow cytometry (See, e.g., U.S. Patent 5,985,660; and Morrison *et al.*, *Cell*, 96:737-49 (1999)).

These techniques allow for the screening of particular populations of cells, such as might be found with hematological malignancies (i.e. minimal residual disease (MRD) in acute leukemic patients) and "non-self" cells in transplantations to prevent Graft-versus-Host Disease (GVHD). Alternatively, these techniques allow for the screening of hematopoietic stem and progenitor cells capable of undergoing proliferation and/or differentiation, as might be found in human umbilical cord blood.

Assays For Antibody Binding

The antibodies of the invention may be assayed for immunospecific binding by any method known in the art. The immunoassays which can be used include but are not limited to competitive and non-competitive assay systems using techniques such as western blots, radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoprecipitation assays, precipitin reactions, gel diffusion precipitin reactions, immunodiffusion assays, agglutination assays, complement-fixation assays, immunoradiometric assays, fluorescent immunoassays, protein A immunoassays, to name but a few. Such assays are routine and well known in the art (see, e.g., Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York, which is incorporated by reference herein in its entirety). Exemplary immunoassays are described briefly below (but are not intended by way of limitation).

Immunoprecipitation protocols generally comprise lysing a population of cells in a lysis buffer such as RIPA buffer (1% NP-40 or Triton X- 100, 1% sodium deoxycholate, 0.1% SDS, 0.15 M NaCl, 0.01 M sodium phosphate at pH 7.2, 1% Trasylol) supplemented with protein phosphatase and/or protease inhibitors (e.g., EDTA, PMSF, aprotinin, sodium vanadate), adding the antibody of interest to the cell lysate, incubating for a period of time (e.g., 1-4 hours) at 4° C, adding protein A

and/or protein G sepharose beads to the cell lysate, incubating for about an hour or more at 4° C, washing the beads in lysis buffer and resuspending the beads in SDS/sample buffer. The ability of the antibody of interest to immunoprecipitate a particular antigen can be assessed by, e.g., western blot analysis. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the binding of the antibody to an antigen and decrease the background (e.g., pre-clearing the cell lysate with sepharose beads). For further discussion regarding immunoprecipitation protocols see, e.g., Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York at 10.16.1.

Western blot analysis generally comprises preparing protein samples, electrophoresis of the protein samples in a polyacrylamide gel (e.g., 8%- 20% SDS-PAGE depending on the molecular weight of the antigen), transferring the protein sample from the polyacrylamide gel to a membrane such as nitrocellulose, PVDF or nylon, blocking the membrane in blocking solution (e.g., PBS with 3% BSA or non-fat milk), washing the membrane in washing buffer (e.g., PBS-Tween 20), blocking the membrane with primary antibody (the antibody of interest) diluted in blocking buffer, washing the membrane in washing buffer, blocking the membrane with a secondary antibody (which recognizes the primary antibody, e.g., an anti-human antibody) conjugated to an enzymatic substrate (e.g., horseradish peroxidase or alkaline phosphatase) or radioactive molecule (e.g., ³²P or ¹²⁵I) diluted in blocking buffer, washing the membrane in wash buffer, and detecting the presence of the antigen. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the signal detected and to reduce the background noise. For further discussion regarding western blot protocols see, e.g., Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York at 10.8.1.

ELISAs comprise preparing antigen, coating the well of a 96 well microtiter plate with the antigen, adding the antibody of interest conjugated to a detectable compound such as an enzymatic substrate (e.g., horseradish peroxidase or alkaline phosphatase) to the well and incubating for a period of time, and detecting the presence of the antigen. In ELISAs the antibody of interest does not have to be conjugated to a detectable compound; instead, a second antibody (which recognizes

the antibody of interest) conjugated to a detectable compound may be added to the well. Further, instead of coating the well with the antigen, the antibody may be coated to the well. In this case, a second antibody conjugated to a detectable compound may be added following the addition of the antigen of interest to the
5 coated well. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the signal detected as well as other variations of ELISAs known in the art. For further discussion regarding ELISAs see, e.g., Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York at 11.2.1.

10 The binding affinity of an antibody to an antigen and the off-rate of an antibody-antigen interaction can be determined by competitive binding assays. One example of a competitive binding assay is a radioimmunoassay comprising the incubation of labeled antigen (e.g., ^3H or ^{125}I) with the antibody of interest in the presence of increasing amounts of unlabeled antigen, and the detection of the
15 antibody bound to the labeled antigen. The affinity of the antibody of interest for a particular antigen and the binding off-rates can be determined from the data by scatchard plot analysis. Competition with a second antibody can also be determined using radioimmunoassays. In this case, the antigen is incubated with antibody of interest conjugated to a labeled compound (e.g., ^3H or ^{125}I) in the presence of
20 increasing amounts of an unlabeled second antibody.

Therapeutic Uses

The present invention is further directed to antibody-based therapies which involve administering antibodies of the invention to an animal, preferably a mammal,
25 and most preferably a human, patient for treating one or more of the disclosed diseases, disorders, or conditions. Therapeutic compounds of the invention include, but are not limited to, antibodies of the invention (including fragments, analogs and derivatives thereof as described herein) and nucleic acids encoding antibodies of the invention (including fragments, analogs and derivatives thereof and anti-idiotypic
30 antibodies as described herein). The antibodies of the invention can be used to treat, inhibit or prevent diseases, disorders or conditions associated with aberrant expression and/or activity of a polypeptide of the invention, including, but not limited to, any

one or more of the diseases, disorders, or conditions described herein. The treatment and/or prevention of diseases, disorders, or conditions associated with aberrant expression and/or activity of a polypeptide of the invention includes, but is not limited to, alleviating symptoms associated with those diseases, disorders or
5 conditions. Antibodies of the invention may be provided in pharmaceutically acceptable compositions as known in the art or as described herein.

A summary of the ways in which the antibodies of the present invention may be used therapeutically includes binding polynucleotides or polypeptides of the present invention locally or systemically in the body or by direct cytotoxicity of the
10 antibody, e.g. as mediated by complement (CDC) or by effector cells (ADCC). Some of these approaches are described in more detail below. Armed with the teachings provided herein, one of ordinary skill in the art will know how to use the antibodies of the present invention for diagnostic, monitoring or therapeutic purposes without undue experimentation.

15 The antibodies of this invention may be advantageously utilized in combination with other monoclonal or chimeric antibodies, or with lymphokines or hematopoietic growth factors (such as, e.g., IL-2, IL-3 and IL-7), for example, which serve to increase the number or activity of effector cells which interact with the antibodies.

20 The antibodies of the invention may be administered alone or in combination with other types of treatments (e.g., radiation therapy, chemotherapy, hormonal therapy, immunotherapy and anti-tumor agents). Generally, administration of products of a species origin or species reactivity (in the case of antibodies) that is the same species as that of the patient is preferred. Thus, in a preferred embodiment,
25 human antibodies, fragments derivatives, analogs, or nucleic acids, are administered to a human patient for therapy or prophylaxis.

It is preferred to use high affinity and/or potent in vivo inhibiting and/or neutralizing antibodies against polypeptides or polynucleotides of the present invention, fragments or regions thereof, for both immunoassays directed to and
30 therapy of disorders related to polynucleotides or polypeptides, including fragments thereof, of the present invention. Such antibodies, fragments, or regions, will preferably have an affinity for polynucleotides or polypeptides of the invention,

including fragments thereof. Preferred binding affinities include those with a dissociation constant or K_d less than 5×10^{-2} M, 10^{-2} M, 5×10^{-3} M, 10^{-3} M, 5×10^{-4} M, 10^{-4} M, 5×10^{-5} M, 10^{-5} M, 5×10^{-6} M, 10^{-6} M, 5×10^{-7} M, 10^{-7} M, 5×10^{-8} M, 10^{-8} M, 5×10^{-9} M, 10^{-9} M, 5×10^{-10} M, 10^{-10} M, 5×10^{-11} M, 10^{-11} M, 5×10^{-12} M, 10^{-12} M, 5×10^{-13} M, 10^{-13} M, 5×10^{-14} M, 10^{-14} M, 5×10^{-15} M, and 10^{-15} M.

Gene Therapy

In a specific embodiment, nucleic acids comprising sequences encoding antibodies or functional derivatives thereof, are administered to treat, inhibit or prevent a disease or disorder associated with aberrant expression and/or activity of a polypeptide of the invention, by way of gene therapy. Gene therapy refers to therapy performed by the administration to a subject of an expressed or expressible nucleic acid. In this embodiment of the invention, the nucleic acids produce their encoded protein that mediates a therapeutic effect.

Any of the methods for gene therapy available in the art can be used according to the present invention. Exemplary methods are described below.

For general reviews of the methods of gene therapy, see Goldspiel et al., *Clinical Pharmacy* 12:488-505 (1993); Wu and Wu, *Biotherapy* 3:87-95 (1991); Tolstoshev, *Ann. Rev. Pharmacol. Toxicol.* 32:573-596 (1993); Mulligan, *Science* 260:926-932 (1993); and Morgan and Anderson, *Ann. Rev. Biochem.* 62:191-217 (1993); May, *TIBTECH* 11(5):155-215 (1993). Methods commonly known in the art of recombinant DNA technology which can be used are described in Ausubel et al. (eds.), *Current Protocols in Molecular Biology*, John Wiley & Sons, NY (1993); and Kriegler, *Gene Transfer and Expression, A Laboratory Manual*, Stockton Press, NY (1990).

In a preferred aspect, the compound comprises nucleic acid sequences encoding an antibody, said nucleic acid sequences being part of expression vectors that express the antibody or fragments or chimeric proteins or heavy or light chains thereof in a suitable host. In particular, such nucleic acid sequences have promoters operably linked to the antibody coding region, said promoter being inducible or constitutive, and, optionally, tissue-specific. In another particular embodiment, nucleic acid molecules are used in which the antibody coding sequences and any other

desired sequences are flanked by regions that promote homologous recombination at a desired site in the genome, thus providing for intrachromosomal expression of the antibody encoding nucleic acids (Koller and Smithies, Proc. Natl. Acad. Sci. USA 86:8932-8935 (1989); Zijlstra et al., Nature 342:435-438 (1989). In specific
5 embodiments, the expressed antibody molecule is a single chain antibody; alternatively, the nucleic acid sequences include sequences encoding both the heavy and light chains, or fragments thereof, of the antibody.

Delivery of the nucleic acids into a patient may be either direct, in which case the patient is directly exposed to the nucleic acid or nucleic acid- carrying vectors, or
10 indirect, in which case, cells are first transformed with the nucleic acids in vitro, then transplanted into the patient. These two approaches are known, respectively, as in vivo or ex vivo gene therapy.

In a specific embodiment, the nucleic acid sequences are directly administered in vivo, where it is expressed to produce the encoded product. This can be
15 accomplished by any of numerous methods known in the art, e.g., by constructing them as part of an appropriate nucleic acid expression vector and administering it so that they become intracellular, e.g., by infection using defective or attenuated retrovirals or other viral vectors (see U.S. Patent No. 4,980,286), or by direct injection of naked DNA, or by use of microparticle bombardment (e.g., a gene gun;
20 Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, encapsulation in liposomes, microparticles, or microcapsules, or by administering them in linkage to a peptide which is known to enter the nucleus, by administering it in linkage to a ligand subject to receptor-mediated endocytosis (see, e.g., Wu and Wu, J. Biol. Chem. 262:4429-4432 (1987)) (which can be used to target
25 cell types specifically expressing the receptors), etc. In another embodiment, nucleic acid-ligand complexes can be formed in which the ligand comprises a fusogenic viral peptide to disrupt endosomes, allowing the nucleic acid to avoid lysosomal degradation. In yet another embodiment, the nucleic acid can be targeted in vivo for cell specific uptake and expression, by targeting a specific receptor (see, e.g., PCT
30 Publications WO 92/06180; WO 92/22635; WO92/20316; WO93/14188, WO 93/20221). Alternatively, the nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression, by homologous recombination

(Koller and Smithies, Proc. Natl. Acad. Sci. USA 86:8932-8935 (1989); Zijlstra et al., Nature 342:435-438 (1989)).

In a specific embodiment, viral vectors that contains nucleic acid sequences encoding an antibody of the invention are used. For example, a retroviral vector can be used (see Miller et al., Meth. Enzymol. 217:581-599 (1993)). These retroviral vectors contain the components necessary for the correct packaging of the viral genome and integration into the host cell DNA. The nucleic acid sequences encoding the antibody to be used in gene therapy are cloned into one or more vectors, which facilitates delivery of the gene into a patient. More detail about retroviral vectors can be found in Boesen et al., Biotherapy 6:291-302 (1994), which describes the use of a retroviral vector to deliver the *mdr1* gene to hematopoietic stem cells in order to make the stem cells more resistant to chemotherapy. Other references illustrating the use of retroviral vectors in gene therapy are: Clowes et al., J. Clin. Invest. 93:644-651 (1994); Kiem et al., Blood 83:1467-1473 (1994); Salmons and Gunzberg, Human Gene Therapy 4:129-141 (1993); and Grossman and Wilson, Curr. Opin. in Genetics and Devel. 3:110-114 (1993).

Adenoviruses are other viral vectors that can be used in gene therapy. Adenoviruses are especially attractive vehicles for delivering genes to respiratory epithelia. Adenoviruses naturally infect respiratory epithelia where they cause a mild disease. Other targets for adenovirus-based delivery systems are liver, the central nervous system, endothelial cells, and muscle. Adenoviruses have the advantage of being capable of infecting non-dividing cells. Kozarsky and Wilson, Current Opinion in Genetics and Development 3:499-503 (1993) present a review of adenovirus-based gene therapy. Bout et al., Human Gene Therapy 5:3-10 (1994) demonstrated the use of adenovirus vectors to transfer genes to the respiratory epithelia of rhesus monkeys. Other instances of the use of adenoviruses in gene therapy can be found in Rosenfeld et al., Science 252:431-434 (1991); Rosenfeld et al., Cell 68:143-155 (1992); Mastrangeli et al., J. Clin. Invest. 91:225-234 (1993); PCT Publication WO94/12649; and Wang, et al., Gene Therapy 2:775-783 (1995). In a preferred embodiment, adenovirus vectors are used.

Adeno-associated virus (AAV) has also been proposed for use in gene therapy (Walsh et al., Proc. Soc. Exp. Biol. Med. 204:289-300 (1993); U.S. Patent No. 5,436,146).

Another approach to gene therapy involves transferring a gene to cells in
5 tissue culture by such methods as electroporation, lipofection, calcium phosphate mediated transfection, or viral infection. Usually, the method of transfer includes the transfer of a selectable marker to the cells. The cells are then placed under selection to isolate those cells that have taken up and are expressing the transferred gene. Those cells are then delivered to a patient.

10 In this embodiment, the nucleic acid is introduced into a cell prior to administration in vivo of the resulting recombinant cell. Such introduction can be carried out by any method known in the art, including but not limited to transfection, electroporation, microinjection, infection with a viral or bacteriophage vector containing the nucleic acid sequences, cell fusion, chromosome-mediated gene
15 transfer, microcell-mediated gene transfer, spheroplast fusion, etc. Numerous techniques are known in the art for the introduction of foreign genes into cells (see, e.g., Loeffler and Behr, Meth. Enzymol. 217:599-618 (1993); Cohen et al., Meth. Enzymol. 217:618-644 (1993); Cline, Pharmac. Ther. 29:69-92m (1985) and may be used in accordance with the present invention, provided that the necessary
20 developmental and physiological functions of the recipient cells are not disrupted. The technique should provide for the stable transfer of the nucleic acid to the cell, so that the nucleic acid is expressible by the cell and preferably heritable and expressible by its cell progeny.

The resulting recombinant cells can be delivered to a patient by various
25 methods known in the art. Recombinant blood cells (e.g., hematopoietic stem or progenitor cells) are preferably administered intravenously. The amount of cells envisioned for use depends on the desired effect, patient state, etc., and can be determined by one skilled in the art.

Cells into which a nucleic acid can be introduced for purposes of gene therapy
30 encompass any desired, available cell type, and include but are not limited to epithelial cells, endothelial cells, keratinocytes, fibroblasts, muscle cells, hepatocytes; blood cells such as Tlymphocytes, Blymphocytes, monocytes, macrophages,

neutrophils, eosinophils, megakaryocytes, granulocytes; various stem or progenitor cells, in particular hematopoietic stem or progenitor cells, e.g., as obtained from bone marrow, umbilical cord blood, peripheral blood, fetal liver, etc.

5 In a preferred embodiment, the cell used for gene therapy is autologous to the patient.

In an embodiment in which recombinant cells are used in gene therapy, nucleic acid sequences encoding an antibody are introduced into the cells such that they are expressible by the cells or their progeny, and the recombinant cells are then administered in vivo for therapeutic effect. In a specific embodiment, stem or
10 progenitor cells are used. Any stem and/or progenitor cells which can be isolated and maintained in vitro can potentially be used in accordance with this embodiment of the present invention (see e.g. PCT Publication WO 94/08598; Stemple and Anderson, Cell 71:973-985 (1992); Rheinwald, Meth. Cell Bio. 21A:229 (1980); and Pittelkow and Scott, Mayo Clinic Proc. 61:771 (1986)).

15 In a specific embodiment, the nucleic acid to be introduced for purposes of gene therapy comprises an inducible promoter operably linked to the coding region, such that expression of the nucleic acid is controllable by controlling the presence or absence of the appropriate inducer of transcription.

20 *Demonstration of Therapeutic or Prophylactic Activity*

The compounds or pharmaceutical compositions of the invention are preferably tested in vitro, and then in vivo for the desired therapeutic or prophylactic activity, prior to use in humans. For example, in vitro assays to demonstrate the therapeutic or prophylactic utility of a compound or pharmaceutical composition
25 include, the effect of a compound on a cell line or a patient tissue sample. The effect of the compound or composition on the cell line and/or tissue sample can be determined utilizing techniques known to those of skill in the art including, but not limited to, rosette formation assays and cell lysis assays. In accordance with the invention, in vitro assays which can be used to determine whether administration of a
30 specific compound is indicated, include in vitro cell culture assays in which a patient tissue sample is grown in culture, and exposed to or otherwise administered a compound, and the effect of such compound upon the tissue sample is observed.

Therapeutic/Prophylactic Administration and Composition

The invention provides methods of treatment, inhibition and prophylaxis by administration to a subject of an effective amount of a compound or pharmaceutical composition of the invention, preferably an antibody of the invention. In a preferred aspect, the compound is substantially purified (e.g., substantially free from substances that limit its effect or produce undesired side-effects). The subject is preferably an animal, including but not limited to animals such as cows, pigs, horses, chickens, cats, dogs, etc., and is preferably a mammal, and most preferably human.

Formulations and methods of administration that can be employed when the compound comprises a nucleic acid or an immunoglobulin are described above; additional appropriate formulations and routes of administration can be selected from among those described herein below.

Various delivery systems are known and can be used to administer a compound of the invention, e.g., encapsulation in liposomes, microparticles, microcapsules, recombinant cells capable of expressing the compound, receptor-mediated endocytosis (see, e.g., Wu and Wu, J. Biol. Chem. 262:4429-4432 (1987)), construction of a nucleic acid as part of a retroviral or other vector, etc. Methods of introduction include but are not limited to intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, and oral routes. The compounds or compositions may be administered by any convenient route, for example by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (e.g., oral mucosa, rectal and intestinal mucosa, etc.) and may be administered together with other biologically active agents. Administration can be systemic or local. In addition, it may be desirable to introduce the pharmaceutical compounds or compositions of the invention into the central nervous system by any suitable route, including intraventricular and intrathecal injection; intraventricular injection may be facilitated by an intraventricular catheter, for example, attached to a reservoir, such as an Ommaya reservoir. Pulmonary administration can also be employed, e.g., by use of an inhaler or nebulizer, and formulation with an aerosolizing agent.

In a specific embodiment, it may be desirable to administer the pharmaceutical compounds or compositions of the invention locally to the area in need of treatment;

this may be achieved by, for example, and not by way of limitation, local infusion during surgery, topical application, e.g., in conjunction with a wound dressing after surgery, by injection, by means of a catheter, by means of a suppository, or by means of an implant, said implant being of a porous, non-porous, or gelatinous material, including membranes, such as sialastic membranes, or fibers. Preferably, when
5 administering a protein, including an antibody, of the invention, care must be taken to use materials to which the protein does not absorb.

In another embodiment, the compound or composition can be delivered in a vesicle, in particular a liposome (see Langer, *Science* 249:1527-1533 (1990); Treat et al., in *Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 353- 365 (1989); Lopez-Berestein, *ibid.*, pp. 317-327; see generally *ibid.*)
10

In yet another embodiment, the compound or composition can be delivered in a controlled release system. In one embodiment, a pump may be used (see Langer, supra; Sefton, *CRC Crit. Ref. Biomed. Eng.* 14:201 (1987); Buchwald et al., *Surgery* 88:507 (1980); Saudek et al., *N. Engl. J. Med.* 321:574 (1989)). In another embodiment, polymeric materials can be used (see *Medical Applications of Controlled Release*, Langer and Wise (eds.), CRC Pres., Boca Raton, Florida (1974); *Controlled Drug Bioavailability, Drug Product Design and Performance*, Smolen and Ball (eds.), Wiley, New York (1984); Ranger and Peppas, J., *Macromol. Sci. Rev.* 20 *Macromol. Chem.* 23:61 (1983); see also Levy et al., *Science* 228:190 (1985); During et al., *Ann. Neurol.* 25:351 (1989); Howard et al., *J. Neurosurg.* 71:105 (1989)). In yet another embodiment, a controlled release system can be placed in proximity of the therapeutic target, i.e., the brain, thus requiring only a fraction of the systemic dose
25 (see, e.g., Goodson, in *Medical Applications of Controlled Release*, supra, vol. 2, pp. 115-138 (1984)).

Other controlled release systems are discussed in the review by Langer (*Science* 249:1527-1533 (1990)).

In a specific embodiment where the compound of the invention is a nucleic acid encoding a protein, the nucleic acid can be administered in vivo to promote
30 expression of its encoded protein, by constructing it as part of an appropriate nucleic acid expression vector and administering it so that it becomes intracellular, e.g., by

use of a retroviral vector (see U.S. Patent No. 4,980,286), or by direct injection, or by use of microparticle bombardment (e.g., a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, or by administering it in linkage to a homeobox-like peptide which is known to enter the nucleus (see e.g.,
5 Joliot et al., Proc. Natl. Acad. Sci. USA 88:1864-1868 (1991)), etc. Alternatively, a nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression, by homologous recombination.

The present invention also provides pharmaceutical compositions. Such compositions comprise a therapeutically effective amount of a compound, and a
10 pharmaceutically acceptable carrier. In a specific embodiment, the term "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly in humans. The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the therapeutic
15 is administered. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water is a preferred carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid
20 carriers, particularly for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering
25 agents. These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like. The composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides. Oral formulation can include standard carriers such as
30 pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, etc. Examples of suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E.W. Martin. Such compositions will contain a therapeutically effective amount of

the compound, preferably in purified form, together with a suitable amount of carrier so as to provide the form for proper administration to the patient. The formulation should suit the mode of administration.

In a preferred embodiment, the composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous administration to human beings. Typically, compositions for intravenous administration are solutions in sterile isotonic aqueous buffer. Where necessary, the composition may also include a solubilizing agent and a local anesthetic such as lignocaine to ease pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

The compounds of the invention can be formulated as neutral or salt forms. Pharmaceutically acceptable salts include those formed with anions such as those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, etc., and those formed with cations such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, etc.

The amount of the compound of the invention which will be effective in the treatment, inhibition and prevention of a disease or disorder associated with aberrant expression and/or activity of a polypeptide of the invention can be determined by standard clinical techniques. In addition, in vitro assays may optionally be employed to help identify optimal dosage ranges. The precise dose to be employed in the formulation will also depend on the route of administration, and the seriousness of the disease or disorder, and should be decided according to the judgment of the practitioner and each patient's circumstances. Effective doses may be extrapolated from dose-response curves derived from in vitro or animal model test systems.

For antibodies, the dosage administered to a patient is typically 0.1 mg/kg to 100 mg/kg of the patient's body weight. Preferably, the dosage administered to a patient is between 0.1 mg/kg and 20 mg/kg of the patient's body weight, more preferably 1 mg/kg to 10 mg/kg of the patient's body weight. Generally, human
5 antibodies have a longer half-life within the human body than antibodies from other species due to the immune response to the foreign polypeptides. Thus, lower dosages of human antibodies and less frequent administration is often possible. Further, the dosage and frequency of administration of antibodies of the invention may be reduced by enhancing uptake and tissue penetration (e.g., into the brain) of the
10 antibodies by modifications such as, for example, lipidation.

The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture,
15 use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

Diagnosis and Imaging

Labeled antibodies, and derivatives and analogs thereof, which specifically
20 bind to a polypeptide of interest can be used for diagnostic purposes to detect, diagnose, or monitor diseases, disorders, and/or conditions associated with the aberrant expression and/or activity of a polypeptide of the invention. The invention provides for the detection of aberrant expression of a polypeptide of interest, comprising (a) assaying the expression of the polypeptide of interest in cells or body
25 fluid of an individual using one or more antibodies specific to the polypeptide interest and (b) comparing the level of gene expression with a standard gene expression level, whereby an increase or decrease in the assayed polypeptide gene expression level compared to the standard expression level is indicative of aberrant expression.

The invention provides a diagnostic assay for diagnosing a disorder,
30 comprising (a) assaying the expression of the polypeptide of interest in cells or body fluid of an individual using one or more antibodies specific to the polypeptide interest and (b) comparing the level of gene expression with a standard gene expression level,

whereby an increase or decrease in the assayed polypeptide gene expression level compared to the standard expression level is indicative of a particular disorder. With respect to cancer, the presence of a relatively high amount of transcript in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Antibodies of the invention can be used to assay protein levels in a biological sample using classical immunohistological methods known to those of skill in the art (e.g., see Jalkanen, et al., J. Cell. Biol. 101:976-985 (1985); Jalkanen, et al., J. Cell Biol. 105:3087-3096 (1987)). Other antibody-based methods useful for detecting protein gene expression include immunoassays, such as the enzyme linked immunosorbent assay (ELISA) and the radioimmunoassay (RIA). Suitable antibody assay labels are known in the art and include enzyme labels, such as, glucose oxidase; radioisotopes, such as iodine (^{125}I , ^{121}I), carbon (^{14}C), sulfur (^{35}S), tritium (^3H), indium (^{112}In), and technetium (^{99}Tc); luminescent labels, such as luminol; and fluorescent labels, such as fluorescein and rhodamine, and biotin.

One aspect of the invention is the detection and diagnosis of a disease or disorder associated with aberrant expression of a polypeptide of interest in an animal, preferably a mammal and most preferably a human. In one embodiment, diagnosis comprises: a) administering (for example, parenterally, subcutaneously, or intraperitoneally) to a subject an effective amount of a labeled molecule which specifically binds to the polypeptide of interest; b) waiting for a time interval following the administering for permitting the labeled molecule to preferentially concentrate at sites in the subject where the polypeptide is expressed (and for unbound labeled molecule to be cleared to background level); c) determining background level; and d) detecting the labeled molecule in the subject, such that detection of labeled molecule above the background level indicates that the subject has a particular disease or disorder associated with aberrant expression of the polypeptide of interest. Background level can be determined by various methods

including, comparing the amount of labeled molecule detected to a standard value previously determined for a particular system.

It will be understood in the art that the size of the subject and the imaging system used will determine the quantity of imaging moiety needed to produce
5 diagnostic images. In the case of a radioisotope moiety, for a human subject, the quantity of radioactivity injected will normally range from about 5 to 20 millicuries of ^{99m}Tc. The labeled antibody or antibody fragment will then preferentially accumulate at the location of cells which contain the specific protein. In vivo tumor imaging is described in S.W. Burchiel et al., "Immunopharmacokinetics of
10 Radiolabeled Antibodies and Their Fragments." (Chapter 13 in Tumor Imaging: The Radiochemical Detection of Cancer, S.W. Burchiel and B. A. Rhodes, eds., Masson Publishing Inc. (1982).

Depending on several variables, including the type of label used and the mode of administration, the time interval following the administration for permitting the
15 labeled molecule to preferentially concentrate at sites in the subject and for unbound labeled molecule to be cleared to background level is 6 to 48 hours or 6 to 24 hours or 6 to 12 hours. In another embodiment the time interval following administration is 5 to 20 days or 5 to 10 days.

In an embodiment, monitoring of the disease or disorder is carried out by
20 repeating the method for diagnosing the disease or disease, for example, one month after initial diagnosis, six months after initial diagnosis, one year after initial diagnosis, etc.

Presence of the labeled molecule can be detected in the patient using methods known in the art for in vivo scanning. These methods depend upon the type of label
25 used. Skilled artisans will be able to determine the appropriate method for detecting a particular label. Methods and devices that may be used in the diagnostic methods of the invention include, but are not limited to, computed tomography (CT), whole body scan such as position emission tomography (PET), magnetic resonance imaging (MRI), and sonography.

30 In a specific embodiment, the molecule is labeled with a radioisotope and is detected in the patient using a radiation responsive surgical instrument (Thurston et al., U.S. Patent No. 5,441,050). In another embodiment, the molecule is labeled with

a fluorescent compound and is detected in the patient using a fluorescence responsive scanning instrument. In another embodiment, the molecule is labeled with a positron emitting metal and is detected in the patient using positron emission-tomography. In yet another embodiment, the molecule is labeled with a paramagnetic label and is
5 detected in a patient using magnetic resonance imaging (MRI).

Kits

The present invention provides kits that can be used in the above methods. In one embodiment, a kit comprises an antibody of the invention, preferably a purified
10 antibody, in one or more containers. In a specific embodiment, the kits of the present invention contain a substantially isolated polypeptide comprising an epitope which is specifically immunoreactive with an antibody included in the kit. Preferably, the kits of the present invention further comprise a control antibody which does not react with the polypeptide of interest. In another specific embodiment, the kits of the present
15 invention contain a means for detecting the binding of an antibody to a polypeptide of interest (e.g., the antibody may be conjugated to a detectable substrate such as a fluorescent compound, an enzymatic substrate, a radioactive compound or a luminescent compound, or a second antibody which recognizes the first antibody may be conjugated to a detectable substrate).

20 In another specific embodiment of the present invention, the kit is a diagnostic kit for use in screening serum containing antibodies specific against proliferative and/or cancerous polynucleotides and polypeptides. Such a kit may include a control antibody that does not react with the polypeptide of interest. Such a kit may include a substantially isolated polypeptide antigen comprising an epitope which is specifically
25 immunoreactive with at least one anti-polypeptide antigen antibody. Further, such a kit includes means for detecting the binding of said antibody to the antigen (e.g., the antibody may be conjugated to a fluorescent compound such as fluorescein or rhodamine which can be detected by flow cytometry). In specific embodiments, the kit may include a recombinantly produced or chemically synthesized polypeptide
30 antigen. The polypeptide antigen of the kit may also be attached to a solid support.

In a more specific embodiment the detecting means of the above-described kit includes a solid support to which said polypeptide antigen is attached. Such a kit may

also include a non-attached reporter-labeled anti-human antibody. In this embodiment, binding of the antibody to the polypeptide antigen can be detected by binding of the said reporter-labeled antibody.

In an additional embodiment, the invention includes a diagnostic kit for use in
5 screening serum containing antigens of the polypeptide of the invention. The diagnostic kit includes a substantially isolated antibody specifically immunoreactive with polypeptide or polynucleotide antigens, and means for detecting the binding of the polynucleotide or polypeptide antigen to the antibody. In one embodiment, the antibody is attached to a solid support. In a specific embodiment, the antibody may be
10 a monoclonal antibody. The detecting means of the kit may include a second, labeled monoclonal antibody. Alternatively, or in addition, the detecting means may include a labeled, competing antigen.

In one diagnostic configuration, test serum is reacted with a solid phase reagent having a surface-bound antigen obtained by the methods of the present
15 invention. After binding with specific antigen antibody to the reagent and removing unbound serum components by washing, the reagent is reacted with reporter-labeled anti-human antibody to bind reporter to the reagent in proportion to the amount of bound anti-antigen antibody on the solid support. The reagent is again washed to remove unbound labeled antibody, and the amount of reporter associated with the
20 reagent is determined. Typically, the reporter is an enzyme which is detected by incubating the solid phase in the presence of a suitable fluorometric, luminescent or colorimetric substrate (Sigma, St. Louis, MO).

The solid surface reagent in the above assay is prepared by known techniques for attaching protein material to solid support material, such as polymeric beads, dip
25 sticks, 96-well plate or filter material. These attachment methods generally include non-specific adsorption of the protein to the support or covalent attachment of the protein, typically through a free amine group, to a chemically reactive group on the solid support, such as an activated carboxyl, hydroxyl, or aldehyde group. Alternatively, streptavidin coated plates can be used in conjunction with biotinylated
30 antigen(s).

Thus, the invention provides an assay system or kit for carrying out this diagnostic method. The kit generally includes a support with surface-bound

recombinant antigens, and a reporter-labeled anti-human antibody for detecting surface-bound anti-antigen antibody.

Fusion Proteins

5 Any polypeptide of the present invention can be used to generate fusion proteins. For example, the polypeptide of the present invention, when fused to a second protein, can be used as an antigenic tag. Antibodies raised against the polypeptide of the present invention can be used to indirectly detect the second protein by binding to the polypeptide. Moreover, because secreted proteins target
10 cellular locations based on trafficking signals, the polypeptides of the present invention can be used as targeting molecules once fused to other proteins.

 Examples of domains that can be fused to polypeptides of the present invention include not only heterologous signal sequences, but also other heterologous functional regions. The fusion does not necessarily need to be direct, but may occur
15 through linker sequences.

 Moreover, fusion proteins may also be engineered to improve characteristics of the polypeptide of the present invention. For instance, a region of additional amino acids, particularly charged amino acids, may be added to the N-terminus of the polypeptide to improve stability and persistence during purification from the host cell
20 or subsequent handling and storage. Also, peptide moieties may be added to the polypeptide to facilitate purification. Such regions may be removed prior to final preparation of the polypeptide. The addition of peptide moieties to facilitate handling of polypeptides are familiar and routine techniques in the art.

 Moreover, polypeptides of the present invention, including fragments, and
25 specifically epitopes, can be combined with parts of the constant domain of immunoglobulins (IgA, IgE, IgG, IgM) or portions thereof (CH1, CH2, CH3, and any combination thereof, including both entire domains and portions thereof), resulting in chimeric polypeptides. These fusion proteins facilitate purification and show an increased half-life in vivo. One reported example describes chimeric proteins
30 consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins. (EP A 394,827; Traunecker et al., Nature 331:84-86 (1988).)

Fusion proteins having disulfide-linked dimeric structures (due to the IgG) can also be more efficient in binding and neutralizing other molecules, than the monomeric secreted protein or protein fragment alone. (Fountoulakis et al., J. Biochem. 270:3958-3964 (1995).) Polynucleotides comprising or alternatively consisting of
5 nucleic acids which encode these fusion proteins are also encompassed by the invention.

Similarly, EP-A-O 464 533 (Canadian counterpart 2045869) discloses fusion proteins comprising various portions of constant region of immunoglobulin molecules together with another human protein or part thereof. In many cases, the Fc part in a
10 fusion protein is beneficial in therapy and diagnosis, and thus can result in, for example, improved pharmacokinetic properties. (EP-A 0232 262.) Alternatively, deleting the Fc part after the fusion protein has been expressed, detected, and purified, would be desired. For example, the Fc portion may hinder therapy and diagnosis if the fusion protein is used as an antigen for immunizations. In drug discovery, for
15 example, human proteins, such as hIL-5, have been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists of hIL-5. (See, D. Bennett et al., J. Molecular Recognition 8:52-58 (1995); K. Johanson et al., J. Biol. Chem. 270:9459-9471 (1995).)

Moreover, the polypeptides of the present invention can be fused to marker
20 sequences, such as a peptide which facilitates purification of the fused polypeptide. In preferred embodiments, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311), among others, many of which are commercially available. As described in Gentz et al., Proc. Natl. Acad. Sci. USA 86:821-824 (1989), for
25 instance, hexa-histidine provides for convenient purification of the fusion protein. Another peptide tag useful for purification, the "HA" tag, corresponds to an epitope derived from the influenza hemagglutinin protein. (Wilson et al., Cell 37:767 (1984).)

Thus, any of these above fusions can be engineered using the polynucleotides
30 or the polypeptides of the present invention.

Vectors, Host Cells, and Protein Production

The present invention also relates to vectors containing the polynucleotide of the present invention, host cells, and the production of polypeptides by recombinant techniques. The vector may be, for example, a phage, plasmid, viral, or retroviral vector. Retroviral vectors may be replication competent or replication defective. In the latter case, viral propagation generally will occur only in complementing host cells.

The polynucleotides may be joined to a vector containing a selectable marker for propagation in a host. Generally, a plasmid vector is introduced in a precipitate, such as a calcium phosphate precipitate, or in a complex with a charged lipid. If the vector is a virus, it may be packaged in vitro using an appropriate packaging cell line and then transduced into host cells.

The polynucleotide insert should be operatively linked to an appropriate promoter, such as the phage lambda PL promoter, the *E. coli* lac, trp, phoA and tac promoters, the SV40 early and late promoters and promoters of retroviral LTRs, to name a few. Other suitable promoters will be known to the skilled artisan. The expression constructs will further contain sites for transcription initiation, termination, and, in the transcribed region, a ribosome binding site for translation. The coding portion of the transcripts expressed by the constructs will preferably include a translation initiating codon at the beginning and a termination codon (UAA, UGA or UAG) appropriately positioned at the end of the polypeptide to be translated.

As indicated, the expression vectors will preferably include at least one selectable marker. Such markers include dihydrofolate reductase, G418 or neomycin resistance for eukaryotic cell culture and tetracycline, kanamycin or ampicillin resistance genes for culturing in *E. coli* and other bacteria. Representative examples of appropriate hosts include, but are not limited to, bacterial cells, such as *E. coli*, *Streptomyces* and *Salmonella typhimurium* cells; fungal cells, such as yeast cells (e.g., *Saccharomyces cerevisiae* or *Pichia pastoris* (ATCC Accession No. 201178)); insect cells such as *Drosophila* S2 and *Spodoptera Sf9* cells; animal cells such as CHO, COS, 293, and Bowes melanoma cells; and plant cells. Appropriate culture mediums and conditions for the above-described host cells are known in the art.

Among vectors preferred for use in bacteria include pQE70, pQE60 and pQE-9, available from QIAGEN, Inc.; pBluescript vectors, Phagescript vectors, pNH8A,

pNH16a, pNH18A, pNH46A, available from Stratagene Cloning Systems, Inc.; and ptrc99a, pKK223-3, pKK233-3, pDR540, pRIT5 available from Pharmacia Biotech, Inc. Among preferred eukaryotic vectors are pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; and pSVK3, pBPV, pMSG and pSVL available
5 from Pharmacia. Preferred expression vectors for use in yeast systems include, but are not limited to pYES2, pYD1, pTEF1/Zeo, pYES2/GS, pPICZ, pGAPZ, pGAPZalph, pPIC9, pPIC3.5, pHIL-D2, pHIL-S1, pPIC3.5K, pPIC9K, and PAO815 (all available from Invitrogen, Carlsbad, CA). Other suitable vectors will be readily apparent to the skilled artisan.

10 Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection, or other methods. Such methods are described in many standard laboratory manuals, such as Davis et al., Basic Methods In Molecular Biology (1986). It is specifically contemplated that the
15 polypeptides of the present invention may in fact be expressed by a host cell lacking a recombinant vector.

A polypeptide of this invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography,
20 phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography ("HPLC") is employed for purification.

Polypeptides of the present invention, and preferably the secreted form, can
25 also be recovered from: products purified from natural sources, including bodily fluids, tissues and cells, whether directly isolated or cultured; products of chemical synthetic procedures; and products produced by recombinant techniques from a prokaryotic or eukaryotic host, including, for example, bacterial, yeast, higher plant, insect, and mammalian cells. Depending upon the host employed in a recombinant
30 production procedure, the polypeptides of the present invention may be glycosylated or may be non-glycosylated. In addition, polypeptides of the invention may also include an initial modified methionine residue, in some cases as a result of host-

mediated processes. Thus, it is well known in the art that the N-terminal methionine encoded by the translation initiation codon generally is removed with high efficiency from any protein after translation in all eukaryotic cells. While the N-terminal methionine on most proteins also is efficiently removed in most prokaryotes, for some
5 proteins, this prokaryotic removal process is inefficient, depending on the nature of the amino acid to which the N-terminal methionine is covalently linked.

In one embodiment, the yeast *Pichia pastoris* is used to express the polypeptide of the present invention in a eukaryotic system. *Pichia pastoris* is a methylotrophic yeast which can metabolize methanol as its sole carbon source. A
10 main step in the methanol metabolism pathway is the oxidation of methanol to formaldehyde using O₂. This reaction is catalyzed by the enzyme alcohol oxidase. In order to metabolize methanol as its sole carbon source, *Pichia pastoris* must generate high levels of alcohol oxidase due, in part, to the relatively low affinity of alcohol oxidase for O₂. Consequently, in a growth medium depending on methanol as a main
15 carbon source, the promoter region of one of the two alcohol oxidase genes (*AOX1*) is highly active. In the presence of methanol, alcohol oxidase produced from the *AOX1* gene comprises up to approximately 30% of the total soluble protein in *Pichia pastoris*. See, Ellis, S.B., *et al.*, *Mol. Cell. Biol.* 5:1111-21 (1985); Koutz, P.J., *et al.*, *Yeast* 5:167-77 (1989); Tschopp, J.F., *et al.*, *Nucl. Acids Res.* 15:3859-76 (1987).
20 Thus, a heterologous coding sequence, such as, for example, a polynucleotide of the present invention, under the transcriptional regulation of all or part of the *AOX1* regulatory sequence is expressed at exceptionally high levels in *Pichia* yeast grown in the presence of methanol.

In one example, the plasmid vector pPIC9K is used to express DNA encoding
25 a polypeptide of the invention, as set forth herein, in a *Pichea* yeast system essentially as described in "*Pichia* Protocols: Methods in Molecular Biology," D.R. Higgins and J. Cregg, eds. The Humana Press, Totowa, NJ, 1998. This expression vector allows expression and secretion of a protein of the invention by virtue of the strong *AOX1* promoter linked to the *Pichia pastoris* alkaline phosphatase (PHO) secretory signal
30 peptide (i.e., leader) located upstream of a multiple cloning site.

Many other yeast vectors could be used in place of pPIC9K, such as, pYES2, pYD1, pTEF1/Zeo, pYES2/GS, pPICZ, pGAPZ, pGAPZalpha, pPIC9, pPIC3.5, pHIL-D2, pHIL-S1, pPIC3.5K, and PAO815, as one skilled in the art would readily appreciate, as long as the proposed expression construct provides appropriately
5 located signals for transcription, translation, secretion (if desired), and the like, including an in-frame AUG as required.

In another embodiment, high-level expression of a heterologous coding sequence, such as, for example, a polynucleotide of the present invention, may be achieved by cloning the heterologous polynucleotide of the invention into an
10 expression vector such as, for example, pGAPZ or pGAPZalpha, and growing the yeast culture in the absence of methanol.

In addition to encompassing host cells containing the vector constructs discussed herein, the invention also encompasses primary, secondary, and immortalized host cells of vertebrate origin, particularly mammalian origin, that have
15 been engineered to delete or replace endogenous genetic material (e.g., coding sequence), and/or to include genetic material (e.g., heterologous polynucleotide sequences) that is operably associated with the polynucleotides of the invention, and which activates, alters, and/or amplifies endogenous polynucleotides. For example, techniques known in the art may be used to operably associate heterologous control
20 regions (e.g., promoter and/or enhancer) and endogenous polynucleotide sequences via homologous recombination, resulting in the formation of a new transcription unit (see, e.g., U.S. Patent No. 5,641,670, issued June 24, 1997; U.S. Patent No. 5,733,761, issued March 31, 1998; International Publication No. WO 96/29411, published September 26, 1996; International Publication No. WO 94/12650,
25 published August 4, 1994; Koller et al., Proc. Natl. Acad. Sci. USA 86:8932-8935 (1989); and Zijlstra et al., Nature 342:435-438 (1989), the disclosures of each of which are incorporated by reference in their entireties).

In addition, polypeptides of the invention can be chemically synthesized using techniques known in the art (e.g., see Creighton, 1983, Proteins: Structures and
30 Molecular Principles, W.H. Freeman & Co., N.Y., and Hunkapiller et al., *Nature*, 310:105-111 (1984)). For example, a polypeptide corresponding to a fragment of a

polypeptide sequence of the invention can be synthesized by use of a peptide synthesizer. Furthermore, if desired, nonclassical amino acids or chemical amino acid analogs can be introduced as a substitution or addition into the polypeptide sequence. Non-classical amino acids include, but are not limited to, to the D-isomers of the common amino acids, 2,4-diaminobutyric acid, α -amino isobutyric acid, 4-aminobutyric acid, Abu, 2-amino butyric acid, g-Abu, e-Ahx, 6-amino hexanoic acid, Aib, 2-amino isobutyric acid, 3-amino propionic acid, ornithine, norleucine, norvaline, hydroxyproline, sarcosine, citrulline, homocitrulline, cysteic acid, t-butylglycine, t-butylalanine, phenylglycine, cyclohexylalanine, b-alanine, fluoro-amino acids, designer amino acids such as b-methyl amino acids, Ca-methyl amino acids, Na-methyl amino acids, and amino acid analogs in general. Furthermore, the amino acid can be D (dextrorotary) or L (levorotary).

The invention encompasses polypeptides which are differentially modified during or after translation, e.g., by glycosylation, acetylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to an antibody molecule or other cellular ligand, etc. Any of numerous chemical modifications may be carried out by known techniques, including but not limited, to specific chemical cleavage by cyanogen bromide, trypsin, chymotrypsin, papain, V8 protease, NaBH_4 ; acetylation, formylation, oxidation, reduction; metabolic synthesis in the presence of tunicamycin; etc.

Additional post-translational modifications encompassed by the invention include, for example, e.g., N-linked or O-linked carbohydrate chains, processing of N-terminal or C-terminal ends), attachment of chemical moieties to the amino acid backbone, chemical modifications of N-linked or O-linked carbohydrate chains, and addition or deletion of an N-terminal methionine residue as a result of procaryotic host cell expression. The polypeptides may also be modified with a detectable label, such as an enzymatic, fluorescent, isotopic or affinity label to allow for detection and isolation of the protein.

Also provided by the invention are chemically modified derivatives of the polypeptides of the invention which may provide additional advantages such as increased solubility, stability and circulating time of the polypeptide, or decreased immunogenicity (see U.S. Patent NO: 4,179,337). The chemical moieties for

derivitization may be selected from water soluble polymers such as polyethylene glycol, ethylene glycol/propylene glycol copolymers, carboxymethylcellulose, dextran, polyvinyl alcohol and the like. The polypeptides may be modified at random positions within the molecule, or at predetermined positions within the molecule and
5 may include one, two, three or more attached chemical moieties.

The polymer may be of any molecular weight, and may be branched or unbranched. For polyethylene glycol, the preferred molecular weight is between about 1 kDa and about 100 kDa (the term "about" indicating that in preparations of polyethylene glycol, some molecules will weigh more, some less, than the stated
10 molecular weight) for ease in handling and manufacturing. Other sizes may be used, depending on the desired therapeutic profile (e.g., the duration of sustained release desired, the effects, if any on biological activity, the ease in handling, the degree or lack of antigenicity and other known effects of the polyethylene glycol to a therapeutic protein or analog). For example, the polyethylene glycol may have an
15 average molecular weight of about 200, 500, 1000, 1500, 2000, 2500, 3000, 3500, 4000, 4500, 5000, 5500, 6000, 6500, 7000, 7500, 8000, 8500, 9000, 9500, 10,000, 10,500, 11,000, 11,500, 12,000, 12,500, 13,000, 13,500, 14,000, 14,500, 15,000, 15,500, 16,000, 16,500, 17,000, 17,500, 18,000, 18,500, 19,000, 19,500, 20,000, 25,000, 30,000, 35,000, 40,000, 50,000, 55,000, 60,000, 65,000, 70,000, 75,000,
20 80,000, 85,000, 90,000, 95,000, or 100,000 kDa.

As noted above, the polyethylene glycol may have a branched structure. Branched polyethylene glycols are described, for example, in U.S. Patent No. 5,643,575; Morpurgo *et al.*, *Appl. Biochem. Biotechnol.* 56:59-72 (1996); Vorobjev *et al.*, *Nucleosides Nucleotides* 18:2745-2750 (1999); and Caliceti *et al.*, *Bioconjug.*
25 *Chem.* 10:638-646 (1999), the disclosures of each of which are incorporated herein by reference.

The polyethylene glycol molecules (or other chemical moieties) should be attached to the protein with consideration of effects on functional or antigenic domains of the protein. There are a number of attachment methods available to those
30 skilled in the art, e.g., EP 0 401 384, herein incorporated by reference (coupling PEG to G-CSF), see also Malik *et al.*, *Exp. Hematol.* 20:1028-1035 (1992) (reporting pegylation of GM-CSF using tresyl chloride). For example, polyethylene glycol may

be covalently bound through amino acid residues via a reactive group, such as, a free amino or carboxyl group. Reactive groups are those to which an activated polyethylene glycol molecule may be bound. The amino acid residues having a free amino group may include lysine residues and the N-terminal amino acid residues; those having a free carboxyl group may include aspartic acid residues glutamic acid residues and the C-terminal amino acid residue. Sulfhydryl groups may also be used as a reactive group for attaching the polyethylene glycol molecules. Preferred for therapeutic purposes is attachment at an amino group, such as attachment at the N-terminus or lysine group.

As suggested above, polyethylene glycol may be attached to proteins via linkage to any of a number of amino acid residues. For example, polyethylene glycol can be linked to a proteins via covalent bonds to lysine, histidine, aspartic acid, glutamic acid, or cysteine residues. One or more reaction chemistries may be employed to attach polyethylene glycol to specific amino acid residues (e.g., lysine, histidine, aspartic acid, glutamic acid, or cysteine) of the protein or to more than one type of amino acid residue (e.g., lysine, histidine, aspartic acid, glutamic acid, cysteine and combinations thereof) of the protein.

One may specifically desire proteins chemically modified at the N-terminus. Using polyethylene glycol as an illustration of the present composition, one may select from a variety of polyethylene glycol molecules (by molecular weight, branching, etc.), the proportion of polyethylene glycol molecules to protein (polypeptide) molecules in the reaction mix, the type of pegylation reaction to be performed, and the method of obtaining the selected N-terminally pegylated protein. The method of obtaining the N-terminally pegylated preparation (i.e., separating this moiety from other monopegylated moieties if necessary) may be by purification of the N-terminally pegylated material from a population of pegylated protein molecules. Selective proteins chemically modified at the N-terminus modification may be accomplished by reductive alkylation which exploits differential reactivity of different types of primary amino groups (lysine versus the N-terminal) available for derivatization in a particular protein. Under the appropriate reaction conditions, substantially selective derivatization of the protein at the N-terminus with a carbonyl group containing polymer is achieved.

As indicated above, pegylation of the proteins of the invention may be accomplished by any number of means. For example, polyethylene glycol may be attached to the protein either directly or by an intervening linker. Linkerless systems for attaching polyethylene glycol to proteins are described in Delgado *et al.*, *Crit. Rev. Thera. Drug Carrier Sys.* 9:249-304 (1992); Francis *et al.*, *Intern. J. of Hematol.* 68:1-18 (1998); U.S. Patent No. 4,002,531; U.S. Patent No. 5,349,052; WO 95/06058; and WO 98/32466, the disclosures of each of which are incorporated herein by reference.

One system for attaching polyethylene glycol directly to amino acid residues of proteins without an intervening linker employs tresylated MPEG, which is produced by the modification of monmethoxy polyethylene glycol (MPEG) using tresylchloride ($\text{ClSO}_2\text{CH}_2\text{CF}_3$). Upon reaction of protein with tresylated MPEG, polyethylene glycol is directly attached to amine groups of the protein. Thus, the invention includes protein-polyethylene glycol conjugates produced by reacting proteins of the invention with a polyethylene glycol molecule having a 2,2,2-trifluoroethane sulphonyl group.

Polyethylene glycol can also be attached to proteins using a number of different intervening linkers. For example, U.S. Patent No. 5,612,460, the entire disclosure of which is incorporated herein by reference, discloses urethane linkers for connecting polyethylene glycol to proteins. Protein-polyethylene glycol conjugates wherein the polyethylene glycol is attached to the protein by a linker can also be produced by reaction of proteins with compounds such as MPEG-succinimidylsuccinate, MPEG activated with 1,1'-carbonyldiimidazole, MPEG-2,4,5-trichloropenylcarbonate, MPEG-p-nitrophenolcarbonate, and various MPEG-succinate derivatives. A number additional polyethylene glycol derivatives and reaction chemistries for attaching polyethylene glycol to proteins are described in WO 98/32466, the entire disclosure of which is incorporated herein by reference. Pegylated protein products produced using the reaction chemistries set out herein are included within the scope of the invention.

The number of polyethylene glycol moieties attached to each protein of the invention (*i.e.*, the degree of substitution) may also vary. For example, the pegylated proteins of the invention may be linked, on average, to 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12,

15, 17, 20, or more polyethylene glycol molecules. Similarly, the average degree of substitution within ranges such as 1-3, 2-4, 3-5, 4-6, 5-7, 6-8, 7-9, 8-10, 9-11, 10-12, 11-13, 12-14, 13-15, 14-16, 15-17, 16-18, 17-19, or 18-20 polyethylene glycol moieties per protein molecule. Methods for determining the degree of substitution are discussed, for example, in Delgado *et al.*, *Crit. Rev. Thera. Drug Carrier Sys.* 9:249-304 (1992).

The polypeptides of the invention may be in monomers or multimers (i.e., dimers, trimers, tetramers and higher multimers). Accordingly, the present invention relates to monomers and multimers of the polypeptides of the invention, their preparation, and compositions (preferably, *Therapeutics*) containing them. In specific embodiments, the polypeptides of the invention are monomers, dimers, trimers or tetramers. In additional embodiments, the multimers of the invention are at least dimers, at least trimers, or at least tetramers.

Multimers encompassed by the invention may be homomers or heteromers. As used herein, the term homomer, refers to a multimer containing only polypeptides corresponding to the amino acid sequence of SEQ ID NO:Y or encoded by the cDNA contained in a deposited clone (including fragments, variants, splice variants, and fusion proteins, corresponding to these polypeptides as described herein). These homomers may contain polypeptides having identical or different amino acid sequences. In a specific embodiment, a homomer of the invention is a multimer containing only polypeptides having an identical amino acid sequence. In another specific embodiment, a homomer of the invention is a multimer containing polypeptides having different amino acid sequences. In specific embodiments, the multimer of the invention is a homodimer (e.g., containing polypeptides having identical or different amino acid sequences) or a homotrimer (e.g., containing polypeptides having identical and/or different amino acid sequences). In additional embodiments, the homomeric multimer of the invention is at least a homodimer, at least a homotrimer, or at least a homotetramer.

As used herein, the term heteromer refers to a multimer containing one or more heterologous polypeptides (i.e., polypeptides of different proteins) in addition to the polypeptides of the invention. In a specific embodiment, the multimer of the invention is a heterodimer, a heterotrimer, or a heterotetramer. In additional

embodiments, the heteromeric multimer of the invention is at least a heterodimer, at least a heterotrimer, or at least a heterotetramer.

Multimers of the invention may be the result of hydrophobic, hydrophilic, ionic and/or covalent associations and/or may be indirectly linked, by for example, liposome formation. Thus, in one embodiment, multimers of the invention, such as, for example, homodimers or homotrimers, are formed when polypeptides of the invention contact one another in solution. In another embodiment, heteromultimers of the invention, such as, for example, heterotrimers or heterotetramers, are formed when polypeptides of the invention contact antibodies to the polypeptides of the invention (including antibodies to the heterologous polypeptide sequence in a fusion protein of the invention) in solution. In other embodiments, multimers of the invention are formed by covalent associations with and/or between the polypeptides of the invention. Such covalent associations may involve one or more amino acid residues contained in the polypeptide sequence (e.g., that recited in the sequence listing, or contained in the polypeptide encoded by a deposited clone). In one instance, the covalent associations are cross-linking between cysteine residues located within the polypeptide sequences which interact in the native (i.e., naturally occurring) polypeptide. In another instance, the covalent associations are the consequence of chemical or recombinant manipulation. Alternatively, such covalent associations may involve one or more amino acid residues contained in the heterologous polypeptide sequence in a fusion protein of the invention.

In one example, covalent associations are between the heterologous sequence contained in a fusion protein of the invention (see, e.g., US Patent Number 5,478,925). In a specific example, the covalent associations are between the heterologous sequence contained in an Fc fusion protein of the invention (as described herein). In another specific example, covalent associations of fusion proteins of the invention are between heterologous polypeptide sequence from another protein that is capable of forming covalently associated multimers, such as for example, osteopontin (see, e.g., International Publication NO: WO 98/49305, the contents of which are herein incorporated by reference in its entirety). In another embodiment, two or more polypeptides of the invention are joined through peptide linkers. Examples include those peptide linkers described in U.S. Pat. No. 5,073,627

(hereby incorporated by reference). Proteins comprising multiple polypeptides of the invention separated by peptide linkers may be produced using conventional recombinant DNA technology.

Another method for preparing multimer polypeptides of the invention involves
5 use of polypeptides of the invention fused to a leucine zipper or isoleucine zipper polypeptide sequence. Leucine zipper and isoleucine zipper domains are polypeptides that promote multimerization of the proteins in which they are found. Leucine zippers were originally identified in several DNA-binding proteins (Landschulz et al., Science 240:1759, (1988)), and have since been found in a variety of different
10 proteins. Among the known leucine zippers are naturally occurring peptides and derivatives thereof that dimerize or trimerize. Examples of leucine zipper domains suitable for producing soluble multimeric proteins of the invention are those described in PCT application WO 94/10308, hereby incorporated by reference. Recombinant fusion proteins comprising a polypeptide of the invention fused to a polypeptide
15 sequence that dimerizes or trimerizes in solution are expressed in suitable host cells, and the resulting soluble multimeric fusion protein is recovered from the culture supernatant using techniques known in the art.

Trimeric polypeptides of the invention may offer the advantage of enhanced biological activity. Preferred leucine zipper moieties and isoleucine moieties are
20 those that preferentially form trimers. One example is a leucine zipper derived from lung surfactant protein D (SPD), as described in Hoppe et al. (FEBS Letters 344:191, (1994)) and in U.S. patent application Ser. No. 08/446,922, hereby incorporated by reference. Other peptides derived from naturally occurring trimeric proteins may be employed in preparing trimeric polypeptides of the invention.

25 In another example, proteins of the invention are associated by interactions between Flag® polypeptide sequence contained in fusion proteins of the invention containing Flag® polypeptide sequence. In a further embodiment, associations proteins of the invention are associated by interactions between heterologous polypeptide sequence contained in Flag® fusion proteins of the invention and anti-
30 Flag® antibody.

The multimers of the invention may be generated using chemical techniques known in the art. For example, polypeptides desired to be contained in the multimers

of the invention may be chemically cross-linked using linker molecules and linker molecule length optimization techniques known in the art (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). Additionally, multimers of the invention may be generated using techniques known in the art to form one or more inter-molecule cross-links between the cysteine residues located within the sequence of the polypeptides desired to be contained in the multimer (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). Further, polypeptides of the invention may be routinely modified by the addition of cysteine or biotin to the C terminus or N-terminus of the polypeptide and techniques known in the art may be applied to generate multimers containing one or more of these modified polypeptides (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). Additionally, techniques known in the art may be applied to generate liposomes containing the polypeptide components desired to be contained in the multimer of the invention (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety).

Alternatively, multimers of the invention may be generated using genetic engineering techniques known in the art. In one embodiment, polypeptides contained in multimers of the invention are produced recombinantly using fusion protein technology described herein or otherwise known in the art (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). In a specific embodiment, polynucleotides coding for a homodimer of the invention are generated by ligating a polynucleotide sequence encoding a polypeptide of the invention to a sequence encoding a linker polypeptide and then further to a synthetic polynucleotide encoding the translated product of the polypeptide in the reverse orientation from the original C-terminus to the N-terminus (lacking the leader sequence) (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). In another embodiment, recombinant techniques described herein or otherwise known in the art are applied to generate recombinant polypeptides of the invention which contain a transmembrane domain (or hydrophobic or signal peptide) and which can be incorporated by membrane reconstitution techniques into

liposomes (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety).

Uses of the Polynucleotides

5 Each of the polynucleotides identified herein can be used in numerous ways as reagents. The following description should be considered exemplary and utilizes known techniques.

 The polynucleotides of the present invention are useful for chromosome identification. There exists an ongoing need to identify new chromosome markers,
10 since few chromosome marking reagents, based on actual sequence data (repeat polymorphisms), are presently available. Each polynucleotide of the present invention can be used as a chromosome marker.

 Briefly, sequences can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp) from the sequences shown in SEQ ID NO:X. Primers can be
15 selected using computer analysis so that primers do not span more than one predicted exon in the genomic DNA. These primers are then used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to the SEQ ID NO:X will yield an amplified fragment.

20 Similarly, somatic hybrids provide a rapid method of PCR mapping the polynucleotides to particular chromosomes. Three or more clones can be assigned per day using a single thermal cycler. Moreover, sublocalization of the polynucleotides can be achieved with panels of specific chromosome fragments. Other gene mapping strategies that can be used include in situ hybridization, prescreening with labeled
25 flow-sorted chromosomes, preselection by hybridization to construct chromosome specific-cDNA libraries and computer mapping techniques (See, e.g., Shuler, Trends Biotechnol 16:456-459 (1998) which is hereby incorporated by reference in its entirety)..

 Precise chromosomal location of the polynucleotides can also be achieved
30 using fluorescence in situ hybridization (FISH) of a metaphase chromosomal spread. This technique uses polynucleotides as short as 500 or 600 bases; however, polynucleotides 2,000-4,000 bp are preferred. For a review of this technique, see

Verma et al., "Human Chromosomes: a Manual of Basic Techniques," Pergamon Press, New York (1988).

For chromosome mapping, the polynucleotides can be used individually (to mark a single chromosome or a single site on that chromosome) or in panels (for marking multiple sites and/or multiple chromosomes).

The polynucleotides of the present invention would likewise be useful for radiation hybrid mapping, HAPPY mapping, and long range restriction mapping. For a review of these techniques and others known in the art, see, e.g., Dear, "Genome Mapping: A Practical Approach," IRL Press at Oxford University Press, London (1997); Aydin, J. Mol. Med. 77:691-694 (1999); Hacia et al., Mol. Psychiatry 3:483-492 (1998); Herrick et al., Chromosome Res. 7:409-423 (1999); Hamilton et al., Methods Cell Biol. 62:265-280 (2000); and/or Ott, J. Hered. 90:68-70 (1999) each of which is hereby incorporated by reference in its entirety.

Once a polynucleotide has been mapped to a precise chromosomal location, the physical position of the polynucleotide can be used in linkage analysis. Linkage analysis establishes coinheritance between a chromosomal location and presentation of a particular disease. (Disease mapping data are found, for example, in V. McKusick, Mendelian Inheritance in Man (available on line through Johns Hopkins University Welch Medical Library) .) Assuming 1 megabase mapping resolution and one gene per 20 kb, a cDNA precisely localized to a chromosomal region associated with the disease could be one of 50-500 potential causative genes.

Thus, once coinheritance is established, differences in the polynucleotide and the corresponding gene between affected and unaffected individuals can be examined. First, visible structural alterations in the chromosomes, such as deletions or translocations, are examined in chromosome spreads or by PCR. If no structural alterations exist, the presence of point mutations are ascertained. Mutations observed in some or all affected individuals, but not in normal individuals, indicates that the mutation may cause the disease. However, complete sequencing of the polypeptide and the corresponding gene from several normal individuals is required to distinguish the mutation from a polymorphism. If a new polymorphism is identified, this polymorphic polypeptide can be used for further linkage analysis.

Furthermore, increased or decreased expression of the gene in affected individuals as compared to unaffected individuals can be assessed using polynucleotides of the present invention. Any of these alterations (altered expression, chromosomal rearrangement, or mutation) can be used as a diagnostic or prognostic marker.

Thus, the invention also provides a diagnostic method useful during diagnosis of a disorder, involving measuring the expression level of polynucleotides of the present invention in cells or body fluid from an individual and comparing the measured gene expression level with a standard level of polynucleotide expression level, whereby an increase or decrease in the gene expression level compared to the standard is indicative of a disorder.

In still another embodiment, the invention includes a kit for analyzing samples for the presence of proliferative and/or cancerous polynucleotides derived from a test subject. In a general embodiment, the kit includes at least one polynucleotide probe containing a nucleotide sequence that will specifically hybridize with a polynucleotide of the present invention and a suitable container. In a specific embodiment, the kit includes two polynucleotide probes defining an internal region of the polynucleotide of the present invention, where each probe has one strand containing a 31'mer-end internal to the region. In a further embodiment, the probes may be useful as primers for polymerase chain reaction amplification.

Where a diagnosis of a disorder, has already been made according to conventional methods, the present invention is useful as a prognostic indicator, whereby patients exhibiting enhanced or depressed polynucleotide of the present invention expression will experience a worse clinical outcome relative to patients expressing the gene at a level nearer the standard level.

By "measuring the expression level of polynucleotide of the present invention" is intended qualitatively or quantitatively measuring or estimating the level of the polypeptide of the present invention or the level of the mRNA encoding the polypeptide in a first biological sample either directly (e.g., by determining or estimating absolute protein level or mRNA level) or relatively (e.g., by comparing to the polypeptide level or mRNA level in a second biological sample). Preferably, the polypeptide level or mRNA level in the first biological sample is measured or

estimated and compared to a standard polypeptide level or mRNA level, the standard being taken from a second biological sample obtained from an individual not having the disorder or being determined by averaging levels from a population of individuals not having a disorder. As will be appreciated in the art, once a standard polypeptide
5 level or mRNA level is known, it can be used repeatedly as a standard for comparison.

By "biological sample" is intended any biological sample obtained from an individual, body fluid, cell line, tissue culture, or other source which contains the polypeptide of the present invention or mRNA. As indicated, biological samples
10 include body fluids (such as semen, lymph, sera, plasma, urine, synovial fluid and spinal fluid) which contain the polypeptide of the present invention, and other tissue sources found to express the polypeptide of the present invention. Methods for obtaining tissue biopsies and body fluids from mammals are well known in the art. Where the biological sample is to include mRNA, a tissue biopsy is the preferred
15 source.

The method(s) provided above may preferably be applied in a diagnostic method and/or kits in which polynucleotides and/or polypeptides are attached to a solid support. In one exemplary method, the support may be a "gene chip" or a "biological chip" as described in US Patents 5,837,832, 5,874,219, and 5,856,174.

20 Further, such a gene chip with polynucleotides of the present invention attached may be used to identify polymorphisms between the polynucleotide sequences, with polynucleotides isolated from a test subject. The knowledge of such polymorphisms (i.e. their location, as well as, their existence) would be beneficial in identifying disease loci for many disorders, including cancerous diseases and conditions. Such a
25 method is described in US Patents 5,858,659 and 5,856,104. The US Patents referenced supra are hereby incorporated by reference in their entirety herein.

The present invention encompasses polynucleotides of the present invention that are chemically synthesized, or reproduced as peptide nucleic acids (PNA), or according to other methods known in the art. The use of PNAs would serve as the
30 preferred form if the polynucleotides are incorporated onto a solid support, or gene chip. For the purposes of the present invention, a peptide nucleic acid (PNA) is a polyamide type of DNA analog and the monomeric units for adenine, guanine,

thymine and cytosine are available commercially (Perceptive Biosystems). Certain components of DNA, such as phosphorus, phosphorus oxides, or deoxyribose derivatives, are not present in PNAs. As disclosed by P. E. Nielsen, M. Egholm, R. H. Berg and O. Buchardt, *Science* 254, 1497 (1991); and M. Egholm, O. Buchardt, L.Christensen, C. Behrens, S. M. Freier, D. A. Driver, R. H. Berg, S. K. Kim, B. Norden, and P. E. Nielsen, *Nature* 365, 666 (1993), PNAs bind specifically and tightly to complementary DNA strands and are not degraded by nucleases. In fact, PNA binds more strongly to DNA than DNA itself does. This is probably because there is no electrostatic repulsion between the two strands, and also the polyamide backbone is more flexible. Because of this, PNA/DNA duplexes bind under a wider range of stringency conditions than DNA/DNA duplexes, making it easier to perform multiplex hybridization. Smaller probes can be used than with DNA due to the strong binding. In addition, it is more likely that single base mismatches can be determined with PNA/DNA hybridization because a single mismatch in a PNA/DNA 15-mer lowers the melting point ($T_{sub.m}$) by 8°-20° C, vs. 4°-16° C for the DNA/DNA 15-mer duplex. Also, the absence of charge groups in PNA means that hybridization can be done at low ionic strengths and reduce possible interference by salt during the analysis.

The present invention is useful for detecting cancer in mammals. In particular the invention is useful during diagnosis of pathological cell proliferative neoplasias which include, but are not limited to: acute myelogenous leukemias including acute monocytic leukemia, acute myeloblastic leukemia, acute promyelocytic leukemia, acute myelomonocytic leukemia, acute erythroleukemia, acute megakaryocytic leukemia, and acute undifferentiated leukemia, etc.; and chronic myelogenous leukemias including chronic myelomonocytic leukemia, chronic granulocytic leukemia, etc. Preferred mammals include monkeys, apes, cats, dogs, cows, pigs, horses, rabbits and humans. Particularly preferred are humans.

Pathological cell proliferative diseases, disorders, and/or conditions are often associated with inappropriate activation of proto-oncogenes. (Gelman, E. P. et al., "The Etiology of Acute Leukemia: Molecular Genetics and Viral Oncology," in *Neoplastic Diseases of the Blood*, Vol 1., Wiernik, P. H. et al. eds., 161-182 (1985)). Neoplasias are now believed to result from the qualitative alteration of a normal

cellular gene product, or from the quantitative modification of gene expression by insertion into the chromosome of a viral sequence, by chromosomal translocation of a gene to a more actively transcribed region, or by some other mechanism. (Gelmann et al., supra) It is likely that mutated or altered expression of specific genes is involved in the pathogenesis of some leukemias, among other tissues and cell types. (Gelmann et al., supra) Indeed, the human counterparts of the oncogenes involved in some animal neoplasias have been amplified or translocated in some cases of human leukemia and carcinoma. (Gelmann et al., supra)

For example, c-myc expression is highly amplified in the non-lymphocytic leukemia cell line HL-60. When HL-60 cells are chemically induced to stop proliferation, the level of c-myc is found to be downregulated. (International Publication Number WO 91/15580) However, it has been shown that exposure of HL-60 cells to a DNA construct that is complementary to the 5' end of c-myc or c-myb blocks translation of the corresponding mRNAs which downregulates expression of the c-myc or c-myb proteins and causes arrest of cell proliferation and differentiation of the treated cells. (International Publication Number WO 91/15580; Wickstrom et al., Proc. Natl. Acad. Sci. 85:1028 (1988); Anfossi et al., Proc. Natl. Acad. Sci. 86:3379 (1989)). However, the skilled artisan would appreciate the present invention's usefulness would not be limited to treatment of proliferative diseases, disorders, and/or conditions of hematopoietic cells and tissues, in light of the numerous cells and cell types of varying origins which are known to exhibit proliferative phenotypes.

In addition to the foregoing, a polynucleotide can be used to control gene expression through triple helix formation or antisense DNA or RNA. Antisense techniques are discussed, for example, in Okano, J. Neurochem. 56: 560 (1991); "Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988). Triple helix formation is discussed in, for instance Lee et al., Nucleic Acids Research 6: 3073 (1979); Cooney et al., Science 241: 456 (1988); and Dervan et al., Science 251: 1360 (1991). Both methods rely on binding of the polynucleotide to a complementary DNA or RNA. For these techniques, preferred polynucleotides are usually oligonucleotides 20 to 40 bases in length and complementary to either the region of the gene involved in transcription (triple helix - see Lee et al., Nucl. Acids Res. 6:3073 (1979); Cooney et al., Science 241:456

(1988); and Dervan et al., Science 251:1360 (1991)) or to the mRNA itself (antisense - Okano, J. Neurochem. 56:560 (1991); Oligodeoxy-nucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988).) Triple helix formation optimally results in a shut-off of RNA transcription from DNA, while
5 antisense RNA hybridization blocks translation of an mRNA molecule into polypeptide. Both techniques are effective in model systems, and the information disclosed herein can be used to design antisense or triple helix polynucleotides in an effort to treat or prevent disease.

Polynucleotides of the present invention are also useful in gene therapy. One
10 goal of gene therapy is to insert a normal gene into an organism having a defective gene, in an effort to correct the genetic defect. The polynucleotides disclosed in the present invention offer a means of targeting such genetic defects in a highly accurate manner. Another goal is to insert a new gene that was not present in the host genome, thereby producing a new trait in the host cell.

15 The polynucleotides are also useful for identifying individuals from minute biological samples. The United States military, for example, is considering the use of restriction fragment length polymorphism (RFLP) for identification of its personnel. In this technique, an individual's genomic DNA is digested with one or more restriction enzymes, and probed on a Southern blot to yield unique bands for
20 identifying personnel. This method does not suffer from the current limitations of "Dog Tags" which can be lost, switched, or stolen, making positive identification difficult. The polynucleotides of the present invention can be used as additional DNA markers for RFLP.

The polynucleotides of the present invention can also be used as an alternative
25 to RFLP, by determining the actual base-by-base DNA sequence of selected portions of an individual's genome. These sequences can be used to prepare PCR primers for amplifying and isolating such selected DNA, which can then be sequenced. Using this technique, individuals can be identified because each individual will have a unique set of DNA sequences. Once an unique ID database is established for an
30 individual, positive identification of that individual, living or dead, can be made from extremely small tissue samples.

Forensic biology also benefits from using DNA-based identification techniques as disclosed herein. DNA sequences taken from very small biological samples such as tissues, e.g., hair or skin, or body fluids, e.g., blood, saliva, semen, synovial fluid, amniotic fluid, breast milk, lymph, pulmonary sputum or surfactant, urine, fecal matter, etc., can be amplified using PCR. In one prior art technique, gene sequences amplified from polymorphic loci, such as DQa class II HLA gene, are used in forensic biology to identify individuals. (Erich, H., PCR Technology, Freeman and Co. (1992).) Once these specific polymorphic loci are amplified, they are digested with one or more restriction enzymes, yielding an identifying set of bands on a Southern blot probed with DNA corresponding to the DQa class II HLA gene. Similarly, polynucleotides of the present invention can be used as polymorphic markers for forensic purposes.

There is also a need for reagents capable of identifying the source of a particular tissue. Such need arises, for example, in forensics when presented with tissue of unknown origin. Appropriate reagents can comprise, for example, DNA probes or primers specific to particular tissue prepared from the sequences of the present invention. Panels of such reagents can identify tissue by species and/or by organ type. In a similar fashion, these reagents can be used to screen tissue cultures for contamination.

In the very least, the polynucleotides of the present invention can be used as molecular weight markers on Southern gels, as diagnostic probes for the presence of a specific mRNA in a particular cell type, as a probe to "subtract-out" known sequences in the process of discovering novel polynucleotides, for selecting and making oligomers for attachment to a "gene chip" or other support, to raise anti-DNA antibodies using DNA immunization techniques, and as an antigen to elicit an immune response.

Uses of the Polypeptides

Each of the polypeptides identified herein can be used in numerous ways. The following description should be considered exemplary and utilizes known techniques.

A polypeptide of the present invention can be used to assay protein levels in a biological sample using antibody-based techniques. For example, protein expression

in tissues can be studied with classical immunohistological methods. (Jalkanen, M., et al., J. Cell. Biol. 101:976-985 (1985); Jalkanen, M., et al., J. Cell . Biol. 105:3087-3096 (1987).) Other antibody-based methods useful for detecting protein gene expression include immunoassays, such as the enzyme linked immunosorbent assay (ELISA) and the radioimmunoassay (RIA). Suitable antibody assay labels are known in the art and include enzyme labels, such as, glucose oxidase, and radioisotopes, such as iodine (^{125}I , ^{121}I), carbon (^{14}C), sulfur (^{35}S), tritium (^3H), indium (^{112}In), and technetium ($^{99\text{m}}\text{Tc}$), and fluorescent labels, such as fluorescein and rhodamine, and biotin.

In addition to assaying secreted protein levels in a biological sample, proteins can also be detected in vivo by imaging. Antibody labels or markers for in vivo imaging of protein include those detectable by X-radiography, NMR or ESR. For X-radiography, suitable labels include radioisotopes such as barium or cesium, which emit detectable radiation but are not overtly harmful to the subject. Suitable markers for NMR and ESR include those with a detectable characteristic spin, such as deuterium, which may be incorporated into the antibody by labeling of nutrients for the relevant hybridoma.

A protein-specific antibody or antibody fragment which has been labeled with an appropriate detectable imaging moiety, such as a radioisotope (for example, ^{131}I , ^{112}In , $^{99\text{m}}\text{Tc}$), a radio-opaque substance, or a material detectable by nuclear magnetic resonance, is introduced (for example, parenterally, subcutaneously, or intraperitoneally) into the mammal. It will be understood in the art that the size of the subject and the imaging system used will determine the quantity of imaging moiety needed to produce diagnostic images. In the case of a radioisotope moiety, for a human subject, the quantity of radioactivity injected will normally range from about 5 to 20 millicuries of $^{99\text{m}}\text{Tc}$. The labeled antibody or antibody fragment will then preferentially accumulate at the location of cells which contain the specific protein. In vivo tumor imaging is described in S.W. Burchiel et al., "Immunopharmacokinetics of Radiolabeled Antibodies and Their Fragments." (Chapter 13 in Tumor Imaging: The Radiochemical Detection of Cancer, S.W. Burchiel and B. A. Rhodes, eds., Masson Publishing Inc. (1982).)

Thus, the invention provides a diagnostic method of a disorder, which involves (a) assaying the expression of a polypeptide of the present invention in cells or body fluid of an individual; (b) comparing the level of gene expression with a standard gene expression level, whereby an increase or decrease in the assayed polypeptide gene expression level compared to the standard expression level is indicative of a disorder. With respect to cancer, the presence of a relatively high amount of transcript in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Moreover, polypeptides of the present invention can be used to treat, prevent, and/or diagnose disease. For example, patients can be administered a polypeptide of the present invention in an effort to replace absent or decreased levels of the polypeptide (e.g., insulin), to supplement absent or decreased levels of a different polypeptide (e.g., hemoglobin S for hemoglobin B, SOD, catalase, DNA repair proteins), to inhibit the activity of a polypeptide (e.g., an oncogene or tumor suppressor), to activate the activity of a polypeptide (e.g., by binding to a receptor), to reduce the activity of a membrane bound receptor by competing with it for free ligand (e.g., soluble TNF receptors used in reducing inflammation), or to bring about a desired response (e.g., blood vessel growth inhibition, enhancement of the immune response to proliferative cells or tissues).

Similarly, antibodies directed to a polypeptide of the present invention can also be used to treat, prevent, and/or diagnose disease. For example, administration of an antibody directed to a polypeptide of the present invention can bind and reduce overproduction of the polypeptide. Similarly, administration of an antibody can activate the polypeptide, such as by binding to a polypeptide bound to a membrane (receptor).

At the very least, the polypeptides of the present invention can be used as molecular weight markers on SDS-PAGE gels or on molecular sieve gel filtration columns using methods well known to those of skill in the art. Polypeptides can also

be used to raise antibodies, which in turn are used to measure protein expression from a recombinant cell, as a way of assessing transformation of the host cell. Moreover, the polypeptides of the present invention can be used to test the following biological activities.

5

Gene Therapy Methods

Another aspect of the present invention is to gene therapy methods for treating or preventing disorders, diseases and conditions. The gene therapy methods relate to the introduction of nucleic acid (DNA, RNA and antisense DNA or RNA) sequences into an animal to achieve expression of a polypeptide of the present invention. This method requires a polynucleotide which codes for a polypeptide of the invention that operatively linked to a promoter and any other genetic elements necessary for the expression of the polypeptide by the target tissue. Such gene therapy and delivery techniques are known in the art, see, for example, WO90/11092, which is herein incorporated by reference.

Thus, for example, cells from a patient may be engineered with a polynucleotide (DNA or RNA) comprising a promoter operably linked to a polynucleotide of the invention *ex vivo*, with the engineered cells then being provided to a patient to be treated with the polypeptide. Such methods are well-known in the art. For example, see Belldgrun et al., J. Natl. Cancer Inst., 85:207-216 (1993); Ferrantini et al., Cancer Research, 53:107-1112 (1993); Ferrantini et al., J. Immunology 153: 4604-4615 (1994); Kaido, T., et al., Int. J. Cancer 60: 221-229 (1995); Ogura et al., Cancer Research 50: 5102-5106 (1990); Santodonato, et al., Human Gene Therapy 7:1-10 (1996); Santodonato, et al., Gene Therapy 4:1246-1255 (1997); and Zhang, et al., Cancer Gene Therapy 3: 31-38 (1996)), which are herein incorporated by reference. In one embodiment, the cells which are engineered are arterial cells. The arterial cells may be reintroduced into the patient through direct injection to the artery, the tissues surrounding the artery, or through catheter injection.

As discussed in more detail below, the polynucleotide constructs can be delivered by any method that delivers injectable materials to the cells of an animal, such as, injection into the interstitial space of tissues (heart, muscle, skin, lung, liver,

30

and the like). The polynucleotide constructs may be delivered in a pharmaceutically acceptable liquid or aqueous carrier.

In one embodiment, the polynucleotide of the invention is delivered as a naked polynucleotide. The term "naked" polynucleotide, DNA or RNA refers to sequences
5 that are free from any delivery vehicle that acts to assist, promote or facilitate entry into the cell, including viral sequences, viral particles, liposome formulations, lipofectin or precipitating agents and the like. However, the polynucleotides of the invention can also be delivered in liposome formulations and lipofectin formulations and the like can be prepared by methods well known to those skilled in the art. Such
10 methods are described, for example, in U.S. Patent Nos. 5,593,972, 5,589,466, and 5,580,859, which are herein incorporated by reference.

The polynucleotide vector constructs of the invention used in the gene therapy method are preferably constructs that will not integrate into the host genome nor will they contain sequences that allow for replication. Appropriate vectors
15 include pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; pSVK3, pBPV, pMSG and pSVL available from Pharmacia; and pEF1/V5, pcDNA3.1, and pRc/CMV2 available from Invitrogen. Other suitable vectors will be readily apparent to the skilled artisan.

Any strong promoter known to those skilled in the art can be used for driving
20 the expression of polynucleotide sequence of the invention. Suitable promoters include adenoviral promoters, such as the adenoviral major late promoter; or heterologous promoters, such as the cytomegalovirus (CMV) promoter; the respiratory syncytial virus (RSV) promoter; inducible promoters, such as the MMT promoter, the metallothionein promoter; heat shock promoters; the albumin promoter;
25 the ApoAI promoter; human globin promoters; viral thymidine kinase promoters, such as the Herpes Simplex thymidine kinase promoter; retroviral LTRs; the b-actin promoter; and human growth hormone promoters. The promoter also may be the native promoter for the polynucleotides of the invention.

Unlike other gene therapy techniques, one major advantage of introducing
30 naked nucleic acid sequences into target cells is the transitory nature of the polynucleotide synthesis in the cells. Studies have shown that non-replicating DNA

sequences can be introduced into cells to provide production of the desired polypeptide for periods of up to six months.

The polynucleotide construct of the invention can be delivered to the interstitial space of tissues within the an animal, including of muscle, skin, brain, lung, liver, spleen, bone marrow, thymus, heart, lymph, blood, bone, cartilage, pancreas, kidney, gall bladder, stomach, intestine, testis, ovary, uterus, rectum, nervous system, eye, gland, and connective tissue. Interstitial space of the tissues comprises the intercellular, fluid, mucopolysaccharide matrix among the reticular fibers of organ tissues, elastic fibers in the walls of vessels or chambers, collagen fibers of fibrous tissues, or that same matrix within connective tissue ensheathing muscle cells or in the lacunae of bone. It is similarly the space occupied by the plasma of the circulation and the lymph fluid of the lymphatic channels. Delivery to the interstitial space of muscle tissue is preferred for the reasons discussed below. They may be conveniently delivered by injection into the tissues comprising these cells. They are preferably delivered to and expressed in persistent, non-dividing cells which are differentiated, although delivery and expression may be achieved in non-differentiated or less completely differentiated cells, such as, for example, stem cells of blood or skin fibroblasts. *In vivo* muscle cells are particularly competent in their ability to take up and express polynucleotides.

For the naked *nucleic acid* sequence injection, an effective dosage amount of DNA or RNA will be in the range of from about 0.05 mg/kg body weight to about 50 mg/kg body weight. Preferably the dosage will be from about 0.005 mg/kg to about 20 mg/kg and more preferably from about 0.05 mg/kg to about 5 mg/kg. Of course, as the artisan of ordinary skill will appreciate, this dosage will vary according to the tissue site of injection. The appropriate and effective dosage of nucleic acid sequence can readily be determined by those of ordinary skill in the art and may depend on the condition being treated and the route of administration.

The preferred route of administration is by the parenteral route of injection into the interstitial space of tissues. However, other parenteral routes may also be used, such as, inhalation of an aerosol formulation particularly for delivery to lungs or bronchial tissues, throat or mucous membranes of the nose. In addition, naked DNA constructs can be delivered to arteries during angioplasty by the catheter used in the procedure.

The naked polynucleotides are delivered by any method known in the art, including, but not limited to, direct needle injection at the delivery site, intravenous injection, topical administration, catheter infusion, and so-called "gene guns". These delivery methods are known in the art:

5 The constructs may also be delivered with delivery vehicles such as viral sequences, viral particles, liposome formulations, lipofectin, precipitating agents, etc. Such methods of delivery are known in the art.

10 In certain embodiments, the polynucleotide constructs of the invention are complexed in a liposome preparation. Liposomal preparations for use in the instant invention include cationic (positively charged), anionic (negatively charged) and neutral preparations. However, cationic liposomes are particularly preferred because a tight charge complex can be formed between the cationic liposome and the polyanionic nucleic acid. Cationic liposomes have been shown to mediate intracellular delivery of plasmid DNA (Felgner et al., Proc. Natl. Acad. Sci. USA ,
15 84:7413-7416 (1987), which is herein incorporated by reference); mRNA (Malone et al., Proc. Natl. Acad. Sci. USA , 86:6077-6081 (1989), which is herein incorporated by reference); and purified transcription factors (Debs et al., J. Biol. Chem., 265:10189-10192 (1990), which is herein incorporated by reference), in functional form.

20 Cationic liposomes are readily available. For example, N[1-2,3-dioleoyloxy)propyl]-N,N,N-triethylammonium (DOTMA) liposomes are particularly useful and are available under the trademark Lipofectin, from GIBCO BRL, Grand Island, N.Y. (See, also, Felgner et al., Proc. Natl Acad. Sci. USA , 84:7413-7416 (1987), which is herein incorporated by reference). Other commercially
25 available liposomes include transfectace (DDAB/DOPE) and DOTAP/DOPE (Boehringer).

Other cationic liposomes can be prepared from readily available materials using techniques well known in the art. See, e.g. PCT Publication NO: WO 90/11092 (which is herein incorporated by reference) for a description of the synthesis of
30 DOTAP (1,2-bis(oleoyloxy)-3-(trimethylammonio)propane) liposomes. Preparation of DOTMA liposomes is explained in the literature, see, e.g., Felgner et al., Proc.

Natl. Acad. Sci. USA, 84:7413-7417, which is herein incorporated by reference.

Similar methods can be used to prepare liposomes from other cationic lipid materials.

Similarly, anionic and neutral liposomes are readily available, such as from Avanti Polar Lipids (Birmingham, Ala.), or can be easily prepared using readily available materials. Such materials include phosphatidyl, choline, cholesterol, phosphatidyl ethanolamine, dioleoylphosphatidyl choline (DOPC), dioleoylphosphatidyl glycerol (DOPG), dioleoylphosphatidyl ethanolamine (DOPE), among others. These materials can also be mixed with the DOTMA and DOTAP starting materials in appropriate ratios. Methods for making liposomes using these materials are well known in the art.

For example, commercially dioleoylphosphatidyl choline (DOPC), dioleoylphosphatidyl glycerol (DOPG), and dioleoylphosphatidyl ethanolamine (DOPE) can be used in various combinations to make conventional liposomes, with or without the addition of cholesterol. Thus, for example, DOPG/DOPC vesicles can be prepared by drying 50 mg each of DOPG and DOPC under a stream of nitrogen gas into a sonication vial. The sample is placed under a vacuum pump overnight and is hydrated the following day with deionized water. The sample is then sonicated for 2 hours in a capped vial, using a Heat Systems model 350 sonicator equipped with an inverted cup (bath type) probe at the maximum setting while the bath is circulated at 15EC. Alternatively, negatively charged vesicles can be prepared without sonication to produce multilamellar vesicles or by extrusion through nucleopore membranes to produce unilamellar vesicles of discrete size. Other methods are known and available to those of skill in the art.

The liposomes can comprise multilamellar vesicles (MLVs), small unilamellar vesicles (SUVs), or large unilamellar vesicles (LUVs), with SUVs being preferred. The various liposome-nucleic acid complexes are prepared using methods well known in the art. See, e.g., Straubinger et al., *Methods of Immunology*, 101:512-527 (1983), which is herein incorporated by reference. For example, MLVs containing nucleic acid can be prepared by depositing a thin film of phospholipid on the walls of a glass tube and subsequently hydrating with a solution of the material to be encapsulated. SUVs are prepared by extended sonication of MLVs to produce a homogeneous population of unilamellar liposomes. The material to be entrapped is added to a

suspension of preformed MLVs and then sonicated. When using liposomes containing cationic lipids, the dried lipid film is resuspended in an appropriate solution such as sterile water or an isotonic buffer solution such as 10 mM Tris/NaCl, sonicated, and then the preformed liposomes are mixed directly with the DNA. The liposome and DNA form a very stable complex due to binding of the positively charged liposomes to the cationic DNA. SUVs find use with small nucleic acid fragments. LUVs are prepared by a number of methods, well known in the art. Commonly used methods include Ca^{2+} -EDTA chelation (Papahadjopoulos et al., Biochim. Biophys. Acta, 394:483 (1975); Wilson et al., Cell, 17:77 (1979)); ether injection (Deamer et al., Biochim. Biophys. Acta, 443:629 (1976); Ostro et al., Biochem. Biophys. Res. Commun., 76:836 (1977); Fraley et al., Proc. Natl. Acad. Sci. USA, 76:3348 (1979)); detergent dialysis (Enoch et al., Proc. Natl. Acad. Sci. USA, 76:145 (1979)); and reverse-phase evaporation (REV) (Fraley et al., J. Biol. Chem., 255:10431 (1980); Szoka et al., Proc. Natl. Acad. Sci. USA, 75:145 (1978); Schaefer-Ridder et al., Science, 215:166 (1982)), which are herein incorporated by reference.

Generally, the ratio of DNA to liposomes will be from about 10:1 to about 1:10. Preferably, the ration will be from about 5:1 to about 1:5. More preferably, the ration will be about 3:1 to about 1:3. Still more preferably, the ratio will be about 1:1.

U.S. Patent NO: 5,676,954 (which is herein incorporated by reference) reports on the injection of genetic material, complexed with cationic liposomes carriers, into mice. U.S. Patent Nos. 4,897,355, 4,946,787, 5,049,386, 5,459,127, 5,589,466, 5,693,622, 5,580,859, 5,703,055, and international publication NO: WO 94/9469 (which are herein incorporated by reference) provide cationic lipids for use in transfecting DNA into cells and mammals. U.S. Patent Nos. 5,589,466, 5,693,622, 5,580,859, 5,703,055, and international publication NO: WO 94/9469 (which are herein incorporated by reference) provide methods for delivering DNA-cationic lipid complexes to mammals.

In certain embodiments, cells are engineered, *ex vivo* or *in vivo*, using a retroviral particle containing RNA which comprises a sequence encoding polypeptides of the invention. Retroviruses from which the retroviral plasmid vectors may be derived include, but are not limited to, Moloney Murine Leukemia Virus, spleen necrosis virus, Rous sarcoma Virus, Harvey Sarcoma Virus, avian leukosis

virus, gibbon ape leukemia virus, human immunodeficiency virus, Myeloproliferative Sarcoma Virus, and mammary tumor virus.

The retroviral plasmid vector is employed to transduce packaging cell lines to form producer cell lines. Examples of packaging cells which may be transfected include, but are not limited to, the PE501, PA317, R-2, R-AM, PA12, T19-14X, VT-19-17-H2, RCRE, RCRIP, GP+E-86, GP+envAm12, and DAN cell lines as described in Miller, Human Gene Therapy, 1:5-14 (1990), which is incorporated herein by reference in its entirety. The vector may transduce the packaging cells through any means known in the art. Such means include, but are not limited to, electroporation, the use of liposomes, and CaPO₄ precipitation. In one alternative, the retroviral plasmid vector may be encapsulated into a liposome, or coupled to a lipid, and then administered to a host.

The producer cell line generates infectious retroviral vector particles which include polynucleotide encoding polypeptides of the invention. Such retroviral vector particles then may be employed, to transduce eukaryotic cells, either *in vitro* or *in vivo*. The transduced eukaryotic cells will express polypeptides of the invention.

In certain other embodiments, cells are engineered, *ex vivo* or *in vivo*, with polynucleotides of the invention contained in an adenovirus vector. Adenovirus can be manipulated such that it encodes and expresses polypeptides of the invention, and at the same time is inactivated in terms of its ability to replicate in a normal lytic viral life cycle. Adenovirus expression is achieved without integration of the viral DNA into the host cell chromosome, thereby alleviating concerns about insertional mutagenesis. Furthermore, adenoviruses have been used as live enteric vaccines for many years with an excellent safety profile (Schwartz et al., Am. Rev. Respir. Dis., 109:233-238 (1974)). Finally, adenovirus mediated gene transfer has been demonstrated in a number of instances including transfer of alpha-1-antitrypsin and CFTR to the lungs of cotton rats (Rosenfeld et al., Science, 252:431-434 (1991); Rosenfeld et al., Cell, 68:143-155 (1992)). Furthermore, extensive studies to attempt to establish adenovirus as a causative agent in human cancer were uniformly negative (Green et al. Proc. Natl. Acad. Sci. USA, 76:6606 (1979)).

Suitable adenoviral vectors useful in the present invention are described, for example, in Kozarsky and Wilson, Curr. Opin. Genet. Devel., 3:499-503 (1993);

Rosenfeld et al., Cell , 68:143-155 (1992); Engelhardt et al., Human Genet. Ther., 4:759-769 (1993); Yang et al., Nature Genet., 7:362-369 (1994); Wilson et al., Nature , 365:691-692 (1993); and U.S. Patent NO: 5,652,224, which are herein incorporated by reference. For example, the adenovirus vector Ad2 is useful and can
5 be grown in human 293 cells. These cells contain the E1 region of adenovirus and constitutively express Ela and Elb, which complement the defective adenoviruses by providing the products of the genes deleted from the vector. In addition to Ad2, other varieties of adenovirus (e.g., Ad3, Ad5, and Ad7) are also useful in the present invention.

10 Preferably, the adenoviruses used in the present invention are replication deficient. Replication deficient adenoviruses require the aid of a helper virus and/or packaging cell line to form infectious particles. The resulting virus is capable of infecting cells and can express a polynucleotide of interest which is operably linked to a promoter, but cannot replicate in most cells. Replication deficient adenoviruses
15 may be deleted in one or more of all or a portion of the following genes: Ela, Elb, E3, E4, E2a, or L1 through L5.

In certain other embodiments, the cells are engineered, *ex vivo* or *in vivo*, using an adeno-associated virus (AAV). AAVs are naturally occurring defective viruses that require helper viruses to produce infectious particles (Muzyczka, Curr.
20 Topics in Microbiol. Immunol., 158:97 (1992)). It is also one of the few viruses that may integrate its DNA into non-dividing cells. Vectors containing as little as 300 base pairs of AAV can be packaged and can integrate, but space for exogenous DNA is limited to about 4.5 kb. Methods for producing and using such AAVs are known in the art. See, for example, U.S. Patent Nos. 5,139,941, 5,173,414, 5,354,678,
25 5,436,146, 5,474,935, 5,478,745, and 5,589,377.

For example, an appropriate AAV vector for use in the present invention will include all the sequences necessary for DNA replication, encapsidation, and host-cell integration. The polynucleotide construct containing polynucleotides of the invention is inserted into the AAV vector using standard cloning methods, such as those found
30 in Sambrook et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Press (1989). The recombinant AAV vector is then transfected into packaging cells which are infected with a helper virus, using any standard technique, including

lipofection, electroporation, calcium phosphate precipitation, etc. Appropriate helper viruses include adenoviruses, cytomegaloviruses, vaccinia viruses, or herpes viruses. Once the packaging cells are transfected and infected, they will produce infectious AAV viral particles which contain the polynucleotide construct of the invention.

- 5 These viral particles are then used to transduce eukaryotic cells, either *ex vivo* or *in vivo*. The transduced cells will contain the polynucleotide construct integrated into its genome, and will express the desired gene product.

Another method of gene therapy involves operably associating heterologous control regions and endogenous polynucleotide sequences (e.g. encoding the
10 polypeptide sequence of interest) via homologous recombination (see, e.g., U.S. Patent NO: 5,641,670, issued June 24, 1997; International Publication NO: WO 96/29411, published September 26, 1996; International Publication NO: WO 94/12650, published August 4, 1994; Koller et al., Proc. Natl. Acad. Sci. USA, 86:8932-8935 (1989); and Zijlstra et al., Nature, 342:435-438 (1989). This method
15 involves the activation of a gene which is present in the target cells, but which is not normally expressed in the cells, or is expressed at a lower level than desired.

Polynucleotide constructs are made, using standard techniques known in the art, which contain the promoter with targeting sequences flanking the promoter. Suitable promoters are described herein. The targeting sequence is sufficiently
20 complementary to an endogenous sequence to permit homologous recombination of the promoter-targeting sequence with the endogenous sequence. The targeting sequence will be sufficiently near the 5' end of the desired endogenous polynucleotide sequence so the promoter will be operably linked to the endogenous sequence upon homologous recombination.

25 The promoter and the targeting sequences can be amplified using PCR. Preferably, the amplified promoter contains distinct restriction enzyme sites on the 5' and 3' ends. Preferably, the 3' end of the first targeting sequence contains the same restriction enzyme site as the 5' end of the amplified promoter and the 5' end of the second targeting sequence contains the same restriction site as the 3' end of the
30 amplified promoter. The amplified promoter and targeting sequences are digested and ligated together.

The promoter-targeting sequence construct is delivered to the cells, either as naked polynucleotide, or in conjunction with transfection-facilitating agents, such as liposomes, viral sequences, viral particles, whole viruses, lipofection, precipitating agents, etc., described in more detail above. The P promoter-targeting sequence can
5 be delivered by any method, included direct needle injection, intravenous injection, topical administration, catheter infusion, particle accelerators, etc. The methods are described in more detail below.

The promoter-targeting sequence construct is taken up by cells. Homologous recombination between the construct and the endogenous sequence takes place, such
10 that an endogenous sequence is placed under the control of the promoter. The promoter then drives the expression of the endogenous sequence.

The polynucleotides encoding polypeptides of the present invention may be administered along with other polynucleotides encoding other angiogenic proteins. Angiogenic proteins include, but are not limited to, acidic and basic fibroblast growth
15 factors, VEGF-1, VEGF-2 (VEGF-C), VEGF-3 (VEGF-B), epidermal growth factor alpha and beta, platelet-derived endothelial cell growth factor, platelet-derived growth factor, tumor necrosis factor alpha, hepatocyte growth factor, insulin like growth factor, colony stimulating factor, macrophage colony stimulating factor, granulocyte/macrophage colony stimulating factor, and nitric oxide synthase.

20 Preferably, the polynucleotide encoding a polypeptide of the invention contains a secretory signal sequence that facilitates secretion of the protein. Typically, the signal sequence is positioned in the coding region of the polynucleotide to be expressed towards or at the 5' end of the coding region. The signal sequence may be homologous or heterologous to the polynucleotide of interest and may be
25 homologous or heterologous to the cells to be transfected. Additionally, the signal sequence may be chemically synthesized using methods known in the art.

Any mode of administration of any of the above-described polynucleotides constructs can be used so long as the mode results in the expression of one or more molecules in an amount sufficient to provide a therapeutic effect. This includes direct
30 needle injection, systemic injection, catheter infusion, biolistic injectors, particle accelerators (i.e., "gene guns"), gelfoam sponge depots, other commercially available depot materials, osmotic pumps (e.g., Alza minipumps), oral or suppository solid

(tablet or pill) pharmaceutical formulations, and decanting or topical applications during surgery. For example, direct injection of naked calcium phosphate-precipitated plasmid into rat liver and rat spleen or a protein-coated plasmid into the portal vein has resulted in gene expression of the foreign gene in the
5 rat livers. (Kaneda et al., Science, 243:375 (1989)).

A preferred method of local administration is by direct injection. Preferably, a recombinant molecule of the present invention complexed with a delivery vehicle is administered by direct injection into or locally within the area of arteries. Administration of a composition locally within the area of arteries refers to injecting
10 the composition centimeters and preferably, millimeters within arteries.

Another method of local administration is to contact a polynucleotide construct of the present invention in or around a surgical wound. For example, a patient can undergo surgery and the polynucleotide construct can be coated on the surface of tissue inside the wound or the construct can be injected into areas of tissue
15 inside the wound.

Therapeutic compositions useful in systemic administration, include recombinant molecules of the present invention complexed to a targeted delivery vehicle of the present invention. Suitable delivery vehicles for use with systemic administration comprise liposomes comprising ligands for targeting the vehicle to a
20 particular site.

Preferred methods of systemic administration, include intravenous injection, aerosol, oral and percutaneous (topical) delivery. Intravenous injections can be performed using methods standard in the art. Aerosol delivery can also be performed using methods standard in the art (see, for example, Stribling et al., Proc. Natl. Acad.
25 Sci. USA, 189:11277-11281 (1992), which is incorporated herein by reference). Oral delivery can be performed by complexing a polynucleotide construct of the present invention to a carrier capable of withstanding degradation by digestive enzymes in the gut of an animal. Examples of such carriers, include plastic capsules or tablets, such as those known in the art. Topical delivery can be performed by mixing a
30 polynucleotide construct of the present invention with a lipophilic reagent (e.g., DMSO) that is capable of passing into the skin.

Determining an effective amount of substance to be delivered can depend upon a number of factors including, for example, the chemical structure and biological activity of the substance, the age and weight of the animal, the precise condition requiring treatment and its severity, and the route of administration. The frequency of treatments depends upon a number of factors, such as the amount of polynucleotide constructs administered per dose, as well as the health and history of the subject. The precise amount, number of doses, and timing of doses will be determined by the attending physician or veterinarian. Therapeutic compositions of the present invention can be administered to any animal, preferably to mammals and birds. Preferred mammals include humans, dogs, cats, mice, rats, rabbits, sheep, cattle, horses and pigs, with humans being particularly

Biological Activities

The polynucleotides or polypeptides, or agonists or antagonists of the present invention can be used in assays to test for one or more biological activities. If these polynucleotides and polypeptides do exhibit activity in a particular assay, it is likely that these molecules may be involved in the diseases associated with the biological activity. Thus, the polynucleotides or polypeptides, or agonists or antagonists could be used to treat the associated disease.

Immune Activity

Polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in treating, preventing, and/or diagnosing diseases, disorders, and/or conditions of the immune system, by, for example, activating or inhibiting the proliferation, differentiation, or mobilization (chemotaxis) of immune cells. Immune cells develop through a process called hematopoiesis, producing myeloid (platelets, red blood cells, neutrophils, and macrophages) and lymphoid (B and T lymphocytes) cells from pluripotent stem cells. The etiology of these immune diseases, disorders, and/or conditions may be genetic, somatic, such as cancer and some autoimmune diseases, acquired (e.g., by chemotherapy or toxins), or infectious. Moreover, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of

the present invention can be used as a marker or detector of a particular immune system disease or disorder.

Polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in treating, preventing, and/or diagnosing diseases, disorders, and/or conditions of hematopoietic cells. Polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention could be used to increase differentiation and proliferation of hematopoietic cells, including the pluripotent stem cells, in an effort to treat or prevent those diseases, disorders, and/or conditions associated with a decrease in certain (or many) types hematopoietic cells. Examples of immunologic deficiency syndromes include, but are not limited to: blood protein diseases, disorders, and/or conditions (e.g., agammaglobulinemia, dysgammaglobulinemia), ataxia telangiectasia, common variable immunodeficiency, Digeorge Syndrome, HIV infection, HTLV-BLV infection, leukocyte adhesion deficiency syndrome, lymphopenia, phagocyte bactericidal dysfunction, severe combined immunodeficiency (SCIDs), Wiskott-Aldrich Disorder, anemia, thrombocytopenia, or hemoglobinuria.

Moreover, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention could also be used to modulate hemostatic (the stopping of bleeding) or thrombolytic activity (clot formation). For example, by increasing hemostatic or thrombolytic activity, polynucleotides or polypeptides, and/or agonists or antagonists of the present invention could be used to treat or prevent blood coagulation diseases, disorders, and/or conditions (e.g., afibrinogenemia, factor deficiencies), blood platelet diseases, disorders, and/or conditions (e.g., thrombocytopenia), or wounds resulting from trauma, surgery, or other causes. Alternatively, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention that can decrease hemostatic or thrombolytic activity could be used to inhibit or dissolve clotting. These molecules could be important in the treatment or prevention of heart attacks (infarction), strokes, or scarring.

The polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in treating, preventing, and/or diagnosing autoimmune disorders. Many autoimmune disorders result from inappropriate

recognition of self as foreign material by immune cells. This inappropriate recognition results in an immune response leading to the destruction of the host tissue. Therefore, the administration of polynucleotides and polypeptides of the invention that can inhibit an immune response, particularly the proliferation, differentiation, or
5 chemotaxis of T-cells, may be an effective therapy in preventing autoimmune disorders.

Autoimmune diseases or disorders that may be treated, prevented, and/or diagnosed by polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention include, but are not limited to, one or more of the following:
10 autoimmune hemolytic anemia, autoimmune neonatal thrombocytopenia, idiopathic thrombocytopenia purpura, autoimmunocytopenia, hemolytic anemia, antiphospholipid syndrome, dermatitis, allergic encephalomyelitis, myocarditis, relapsing polychondritis, rheumatic heart disease, glomerulonephritis (e.g. IgA nephropathy), Multiple Sclerosis, Neuritis, Uveitis Ophthalmia,
15 Polyendocrinopathies, Purpura (e.g., Henloch-Scoenlein purpura), Reiter's Disease, Stiff-Man Syndrome, Autoimmune Pulmonary Inflammation, Autism, Guillain-Barre Syndrome, insulin dependent diabetes mellitus, and autoimmune inflammatory eye, autoimmune thyroiditis, hypothyroidism (i.e., Hashimoto's thyroiditis, systemic lupus erythematosus, Goodpasture's syndrome, Pemphigus, Receptor autoimmunities such
20 as, for example, (a) Graves' Disease, (b) Myasthenia Gravis, and (c) insulin resistance, autoimmune hemolytic anemia, autoimmune thrombocytopenic purpura, rheumatoid arthritis, scleroderma with anti-collagen antibodies, mixed connective tissue disease, polymyositis/dermatomyositis, pernicious anemia, idiopathic Addison's disease, infertility, glomerulonephritis such as primary glomerulonephritis and IgA
25 nephropathy, bullous pemphigoid, Sjogren's syndrome, diabetes mellitus, and adrenergic drug resistance (including adrenergic drug resistance with asthma or cystic fibrosis), chronic active hepatitis, primary biliary cirrhosis, other endocrine gland failure, vitiligo, vasculitis, post-MI, cardiomyopathy syndrome, urticaria, atopic dermatitis, asthma, inflammatory myopathies, and other inflammatory, granulomatous,
30 degenerative, and atrophic disorders.

Additional autoimmune disorders (that are probable) that may be treated, prevented, and/or diagnosed with the compositions of the invention include, but are

not limited to, rheumatoid arthritis (often characterized, e.g., by immune complexes in joints), scleroderma with anti-collagen antibodies (often characterized, e.g., by nucleolar and other nuclear antibodies), mixed connective tissue disease (often characterized, e.g., by antibodies to extractable nuclear antigens (e.g.,
5 ribonucleoprotein)), polymyositis (often characterized, e.g., by nonhistone ANA), pernicious anemia (often characterized, e.g., by antiparietal cell, microsomes, and intrinsic factor antibodies), idiopathic Addison's disease (often characterized, e.g., by humoral and cell-mediated adrenal cytotoxicity, infertility (often characterized, e.g., by antispermatozoal antibodies), glomerulonephritis (often characterized, e.g., by
10 glomerular basement membrane antibodies or immune complexes), bullous pemphigoid (often characterized, e.g., by IgG and complement in basement membrane), Sjogren's syndrome (often characterized, e.g., by multiple tissue antibodies, and/or a specific nonhistone ANA (SS-B)), diabetes mellitus (often characterized, e.g., by cell-mediated and humoral islet cell antibodies), and adrenergic
15 drug resistance (including adrenergic drug resistance with asthma or cystic fibrosis) (often characterized, e.g., by beta-adrenergic receptor antibodies).

Additional autoimmune disorders (that are possible) that may be treated, prevented, and/or diagnosed with the compositions of the invention include, but are not limited to, chronic active hepatitis (often characterized, e.g., by smooth muscle
20 antibodies), primary biliary cirrhosis (often characterized, e.g., by mitochondrial antibodies), other endocrine gland failure (often characterized, e.g., by specific tissue antibodies in some cases), vitiligo (often characterized, e.g., by melanocyte antibodies), vasculitis (often characterized, e.g., by Ig and complement in vessel walls and/or low serum complement), post-MI (often characterized, e.g., by myocardial
25 antibodies), cardiomyopathy syndrome (often characterized, e.g., by myocardial antibodies), urticaria (often characterized, e.g., by IgG and IgM antibodies to IgE), atopic dermatitis (often characterized, e.g., by IgG and IgM antibodies to IgE), asthma (often characterized, e.g., by IgG and IgM antibodies to IgE), and many other inflammatory, granulomatous, degenerative, and atrophic disorders.

30 In a preferred embodiment, the autoimmune diseases and disorders and/or conditions associated with the diseases and disorders recited above are treated,

prevented, and/or diagnosed using for example, antagonists or agonists, polypeptides or polynucleotides, or antibodies of the present invention.

In a preferred embodiment polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention could be used as an agent to boost
5 immunoresponsiveness among B cell and/or T cell immunodeficient individuals.

B cell immunodeficiencies that may be ameliorated or treated by administering the polypeptides or polynucleotides of the invention, and/or agonists thereof, include, but are not limited to, severe combined immunodeficiency (SCID)-X linked, SCID-autosomal, adenosine deaminase deficiency (ADA deficiency), X-
10 linked agammaglobulinemia (XLA), Bruton's disease, congenital agammaglobulinemia, X-linked infantile agammaglobulinemia, acquired agammaglobulinemia, adult onset agammaglobulinemia, late-onset agammaglobulinemia, dysgammaglobulinemia, hypogammaglobulinemia, transient hypogammaglobulinemia of infancy, unspecified hypogammaglobulinemia,
15 agammaglobulinemia, common variable immunodeficiency (CVI) (acquired), Wiskott-Aldrich Syndrome (WAS), X-linked immunodeficiency with hyper IgM, non X-linked immunodeficiency with hyper IgM, selective IgA deficiency, IgG subclass deficiency (with or without IgA deficiency), antibody deficiency with normal or elevated Igs, immunodeficiency with thymoma, Ig heavy chain deletions, kappa
20 chain deficiency, B cell lymphoproliferative disorder (BLPD), selective IgM immunodeficiency, recessive agammaglobulinemia (Swiss type), reticular dysgenesis, neonatal neutropenia, severe congenital leukopenia, thymic aplasia or dysplasia with immunodeficiency, ataxia-telangiectasia, short limbed dwarfism, X-linked lymphoproliferative syndrome (XLP), Nezelof
25 syndrome-combined immunodeficiency with Igs, purine nucleoside phosphorylase deficiency (PNP), MHC Class II deficiency (Bare Lymphocyte Syndrome) and severe combined immunodeficiency.

T cell deficiencies that may be ameliorated or treated by administering the polypeptides or polynucleotides of the invention, and/or agonists thereof include, but
30 are not limited to, for example, DiGeorge anomaly, thymic hypoplasia, third and fourth pharyngeal pouch syndrome, 22q11.2 deletion, chronic mucocutaneous candidiasis, natural killer cell deficiency (NK), idiopathic CD4+ T-lymphocytopenia,

immunodeficiency with predominant T cell defect (unspecified), and unspecified immunodeficiency of cell mediated immunity. In specific embodiments, DiGeorge anomaly or conditions associated with DiGeorge anomaly are ameliorated or treated by, for example, administering the polypeptides or polynucleotides of the invention, or antagonists or agonists thereof.

Other immunodeficiencies that may be ameliorated or treated by administering polypeptides or polynucleotides of the invention, and/or agonists thereof, include, but are not limited to, severe combined immunodeficiency (SCID; e.g., X-linked SCID, autosomal SCID, and adenosine deaminase deficiency), ataxia-telangiectasia, Wiskott-Aldrich syndrome, short-limber dwarfism, X-linked lymphoproliferative syndrome (XLP), Nezelof syndrome (e.g., purine nucleoside phosphorylase deficiency), MHC Class II deficiency. In specific embodiments, ataxia-telangiectasia or conditions associated with ataxia-telangiectasia are ameliorated or treated by administering the polypeptides or polynucleotides of the invention, and/or agonists thereof.

In a specific preferred embodiment, rheumatoid arthritis is treated, prevented, and/or diagnosed using polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention. In another specific preferred embodiment, systemic lupus erythematosus is treated, prevented, and/or diagnosed using polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention. In another specific preferred embodiment, idiopathic thrombocytopenia purpura is treated, prevented, and/or diagnosed using polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention. In another specific preferred embodiment IgA nephropathy is treated, prevented, and/or diagnosed using polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention. In a preferred embodiment, the autoimmune diseases and disorders and/or conditions associated with the diseases and disorders recited above are treated, prevented, and/or diagnosed using antibodies against the protein of the invention.

Similarly, allergic reactions and conditions, such as asthma (particularly allergic asthma) or other respiratory problems, may also be treated, prevented, and/or diagnosed using polypeptides, antibodies, or polynucleotides of the invention, and/or

agonists or antagonists thereof. Moreover, these molecules can be used to treat, prevent, and/or diagnose anaphylaxis, hypersensitivity to an antigenic molecule, or blood group incompatibility.

Moreover, inflammatory conditions may also be treated, diagnosed, and/or prevented with polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention. Such inflammatory conditions include, but are not limited to, for example, respiratory disorders (such as, e.g., asthma and allergy); gastrointestinal disorders (such as, e.g., inflammatory bowel disease); cancers (such as, e.g., gastric, ovarian, lung, bladder, liver, and breast); CNS disorders (such as, e.g., multiple sclerosis, blood-brain barrier permeability, ischemic brain injury and/or stroke, traumatic brain injury, neurodegenerative disorders (such as, e.g., Parkinson's disease and Alzheimer's disease), AIDS-related dementia, and prion disease); cardiovascular disorders (such as, e.g., atherosclerosis, myocarditis, cardiovascular disease, and cardiopulmonary bypass complications); as well as many additional diseases, conditions, and disorders that are characterized by inflammation (such as, e.g., chronic hepatitis (B and C), rheumatoid arthritis, gout, trauma, septic shock, pancreatitis, sarcoidosis, dermatitis, renal ischemia-reperfusion injury, Grave's disease, systemic lupus erythematosus, diabetes mellitus (i.e., type 1 diabetes), and allogenic transplant rejection).

In specific embodiments, polypeptides, antibodies, or polynucleotides of the invention, and/or agonists or antagonists thereof, are useful to treat, diagnose, and/or prevent transplantation rejections, graft-versus-host disease, autoimmune and inflammatory diseases (e.g., immune complex-induced vasculitis, glomerulonephritis, hemolytic anemia, myasthenia gravis, type II collagen-induced arthritis, experimental allergic and hyperacute xenograft rejection, rheumatoid arthritis, and systemic lupus erythematosus (SLE)). Organ rejection occurs by host immune cell destruction of the transplanted tissue through an immune response. Similarly, an immune response is also involved in GVHD, but, in this case, the foreign transplanted immune cells destroy the host tissues. Polypeptides, antibodies, or polynucleotides of the invention, and/or agonists or antagonists thereof, that inhibit an immune response, particularly the activation, proliferation, differentiation, or chemotaxis of T-cells, may be an effective therapy in preventing organ rejection or GVHD.

Similarly, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may also be used to modulate and/or diagnose inflammation. For example, since polypeptides, antibodies, or polynucleotides of the invention, and/or agonists or antagonists of the invention may inhibit the activation, proliferation and/or differentiation of cells involved in an inflammatory response, these molecules can be used to treat, diagnose, or prognose, inflammatory conditions, both chronic and acute conditions, including, but not limited to, inflammation associated with infection (e.g., septic shock, sepsis, or systemic inflammatory response syndrome (SIRS)), ischemia-reperfusion injury, endotoxin lethality, arthritis, complement-mediated hyperacute rejection, nephritis, cytokine or chemokine induced lung injury, inflammatory bowel disease, Crohn's disease, and resulting from over production of cytokines (e.g., TNF or IL-1.).

Polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the invention can be used to treat, detect, and/or prevent infectious agents. For example, by increasing the immune response, particularly increasing the proliferation activation and/or differentiation of B and/or T cells, infectious diseases may be treated, detected, and/or prevented. The immune response may be increased by either enhancing an existing immune response, or by initiating a new immune response. Alternatively, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may also directly inhibit the infectious agent (refer to section of application listing infectious agents, etc), without necessarily eliciting an immune response.

Additional preferred embodiments of the invention include, but are not limited to, the use of polypeptides, antibodies, polynucleotides and/or agonists or antagonists in the following applications:

Administration to an animal (e.g., mouse, rat, rabbit, hamster, guinea pig, pigs, micro-pig, chicken, camel, goat, horse, cow, sheep, dog, cat, non-human primate, and human, most preferably human) to boost the immune system to produce increased quantities of one or more antibodies (e.g., IgG, IgA, IgM, and IgE), to induce higher affinity antibody production (e.g., IgG, IgA, IgM, and IgE), and/or to increase an immune response.

Administration to an animal (including, but not limited to, those listed above, and also including transgenic animals) incapable of producing functional endogenous antibody molecules or having an otherwise compromised endogenous immune system, but which is capable of producing human immunoglobulin molecules by means of a reconstituted or partially reconstituted immune system from another animal (see, e.g., published PCT Application Nos. WO98/24893, WO/9634096, WO/9633735, and WO/9110741.

A vaccine adjuvant that enhances immune responsiveness to specific antigen.

An adjuvant to enhance tumor-specific immune responses.

10 An adjuvant to enhance anti-viral immune responses. Anti-viral immune responses that may be enhanced using the compositions of the invention as an adjuvant, include virus and virus associated diseases or symptoms described herein or otherwise known in the art. In specific embodiments, the compositions of the invention are used as an adjuvant to enhance an immune response to a virus, disease, or symptom selected from the group consisting of: AIDS, meningitis, Dengue, EBV, and hepatitis (e.g., hepatitis B). In another specific embodiment, the compositions of the invention are used as an adjuvant to enhance an immune response to a virus, disease, or symptom selected from the group consisting of: HIV/AIDS, Respiratory syncytial virus, Dengue, Rotavirus, Japanese B encephalitis, Influenza A and B, Parainfluenza, Measles, Cytomegalovirus, Rabies, Junin, Chikungunya, Rift Valley fever, Herpes simplex, and yellow fever.

25 An adjuvant to enhance anti-bacterial or anti-fungal immune responses. Anti-bacterial or anti-fungal immune responses that may be enhanced using the compositions of the invention as an adjuvant, include bacteria or fungus and bacteria or fungus associated diseases or symptoms described herein or otherwise known in the art. In specific embodiments, the compositions of the invention are used as an adjuvant to enhance an immune response to a bacteria or fungus, disease, or symptom selected from the group consisting of: tetanus, Diphtheria, botulism, and meningitis type B. In another specific embodiment, the compositions of the invention are used as an adjuvant to enhance an immune response to a bacteria or fungus, disease, or symptom selected from the group consisting of: *Vibrio cholerae*, *Mycobacterium leprae*, *Salmonella typhi*, *Salmonella paratyphi*, *Meissneria meningitidis*,

Streptococcus pneumoniae, Group B streptococcus, *Shigella spp.*, Enterotoxigenic *Escherichia coli*, Enterohemorrhagic *E. coli*, *Borrelia burgdorferi*, and Plasmodium (malaria).

5 An adjuvant to enhance anti-parasitic immune responses. Anti-parasitic immune responses that may be enhanced using the compositions of the invention as an adjuvant, include parasite and parasite associated diseases or symptoms described herein or otherwise known in the art. In specific embodiments, the compositions of the invention are used as an adjuvant to enhance an immune response to a parasite. In another specific embodiment, the compositions of the invention are used as an
10 adjuvant to enhance an immune response to Plasmodium (malaria).

As a stimulator of B cell responsiveness to pathogens.

As an activator of T cells.

As an agent that elevates the immune status of an individual prior to their receipt of immunosuppressive therapies.

15 As an agent to induce higher affinity antibodies.

As an agent to increase serum immunoglobulin concentrations.

As an agent to accelerate recovery of immunocompromised individuals.

As an agent to boost immunoresponsiveness among aged populations.

As an immune system enhancer prior to, during, or after bone marrow
20 transplant and/or other transplants (e.g., allogeneic or xenogeneic organ transplantation). With respect to transplantation, compositions of the invention may be administered prior to, concomitant with, and/or after transplantation. In a specific embodiment, compositions of the invention are administered after transplantation, prior to the beginning of recovery of T-cell populations. In another specific
25 embodiment, compositions of the invention are first administered after transplantation after the beginning of recovery of T cell populations, but prior to full recovery of B cell populations.

As an agent to boost immunoresponsiveness among individuals having an acquired loss of B cell function. Conditions resulting in an acquired loss of B cell
30 function that may be ameliorated or treated by administering the polypeptides, antibodies, polynucleotides and/or agonists or antagonists thereof, include, but are

not limited to, HIV Infection, AIDS, bone marrow transplant, and B cell chronic lymphocytic leukemia (CLL).

As an agent to boost immunoresponsiveness among individuals having a temporary immune deficiency. Conditions resulting in a temporary immune deficiency that may be ameliorated or treated by administering the polypeptides, antibodies, polynucleotides and/or agonists or antagonists thereof, include, but are not limited to, recovery from viral infections (e.g., influenza), conditions associated with malnutrition, recovery from infectious mononucleosis, or conditions associated with stress, recovery from measles, recovery from blood transfusion, recovery from surgery.

As a regulator of antigen presentation by monocytes, dendritic cells, and/or B-cells. In one embodiment, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention enhance antigen presentation or antagonizes antigen presentation in vitro or in vivo. Moreover, in related embodiments, said enhancement or antagonization of antigen presentation may be useful as an anti-tumor treatment or to modulate the immune system.

As an agent to direct an individuals immune system towards development of a humoral response (i.e. TH2) as opposed to a TH1 cellular response.

As a means to induce tumor proliferation and thus make it more susceptible to anti-neoplastic agents. For example, multiple myeloma is a slowly dividing disease and is thus refractory to virtually all anti-neoplastic regimens. If these cells were forced to proliferate more rapidly their susceptibility profile would likely change.

As a stimulator of B cell production in pathologies such as AIDS, chronic lymphocyte disorder and/or Common Variable Immunodeficiency.

As a therapy for generation and/or regeneration of lymphoid tissues following surgery, trauma or genetic defect..

As a gene-based therapy for genetically inherited disorders resulting in immuno-incompetence such as observed among SCID patients.

As an antigen for the generation of antibodies to inhibit or enhance immune mediated responses against polypeptides of the invention.

As a means of activating T cells.

As a means of activating monocytes/macrophages to defend against parasitic diseases that effect monocytes such as Leshmania.

As pretreatment of bone marrow samples prior to transplant. Such treatment would increase B cell representation and thus accelerate recover.

5 As a means of regulating secreted cytokines that are elicited by polypeptides of the invention.

Additionally, polypeptides or polynucleotides of the invention, and/or agonists thereof, may be used to treat or prevent IgE-mediated allergic reactions. Such allergic reactions include, but are not limited to, asthma, rhinitis, and eczema.

10 All of the above described applications as they may apply to veterinary medicine.

Antagonists of the invention include, for example, binding and/or inhibitory antibodies, antisense nucleic acids, or ribozymes. These would be expected to reverse many of the activities of the ligand described above as well as find clinical or
15 practical application as:

A means of blocking various aspects of immune responses to foreign agents or self. Examples include autoimmune disorders such as lupus, and arthritis, as well as immunoresponsiveness to skin allergies, inflammation, bowel disease, injury and pathogens.

20 A therapy for preventing the B cell proliferation and Ig secretion associated with autoimmune diseases such as idiopathic thrombocytopenic purpura, systemic lupus erythramatosus and MS.

An inhibitor of B and/or T cell migration in endothelial cells. This activity disrupts tissue architecture or cognate responses and is useful, for example in
25 disrupting immune responses, and blocking sepsis.

An inhibitor of graft versus host disease or transplant rejection.

A therapy for B cell and/or T cell malignancies such as ALL, Hodgkins disease, non-Hodgkins lymphoma, Chronic lymphocyte leukemia, plasmacytomas, multiple myeloma, Burkitt's lymphoma, and EBV-transformed diseases.

30 A therapy for chronic hypergammaglobulinemeia evident in such diseases as monoclonal gammopathy of undetermined significance (MGUS), Waldenstrom's disease, related idiopathic monoclonal gammopathies, and plasmacytomas.

A therapy for decreasing cellular proliferation of Large B-cell Lymphomas.

A means of decreasing the involvement of B cells and Ig associated with Chronic Myelogenous Leukemia.

An immunosuppressive agent(s).

5 Polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be used to modulate IgE concentrations in vitro or in vivo.

In another embodiment, administration of polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the invention, may be used to treat or prevent IgE-mediated allergic reactions including, but not limited to, asthma,
10 rhinitis, and eczema.

The agonists and antagonists may be employed in a composition with a pharmaceutically acceptable carrier, e.g., as described herein.

The agonists or antagonists may be employed for instance to inhibit polypeptide chemotaxis and activation of macrophages and their precursors, and of
15 neutrophils, basophils, B lymphocytes and some T-cell subsets, e.g., activated and CD8 cytotoxic T cells and natural killer cells, in certain auto-immune and chronic inflammatory and infective diseases. Examples of autoimmune diseases are described herein and include multiple sclerosis, and insulin-dependent diabetes. The antagonists or agonists may also be employed to treat infectious diseases including
20 silicosis, sarcoidosis, idiopathic pulmonary fibrosis by, for example, preventing the recruitment and activation of mononuclear phagocytes. They may also be employed to treat idiopathic hyper-eosinophilic syndrome by, for example, preventing eosinophil production and migration. The antagonists or agonists may also be employed for treating atherosclerosis, for example, by preventing monocyte
25 infiltration in the artery wall.

Antibodies against polypeptides of the invention may be employed to treat ARDS.

Agonists and/or antagonists of the invention also have uses in stimulating wound and tissue repair, stimulating angiogenesis, stimulating the repair of vascular
30 or lymphatic diseases or disorders. Additionally, agonists and antagonists of the invention may be used to stimulate the regeneration of mucosal surfaces.

In a specific embodiment, polynucleotides or polypeptides, and/or agonists thereof are used to treat or prevent a disorder characterized by primary or acquired immunodeficiency, deficient serum immunoglobulin production, recurrent infections, and/or immune system dysfunction. Moreover, polynucleotides or polypeptides, and/or agonists thereof may be used to treat or prevent infections of the joints, bones, skin, and/or parotid glands, blood-borne infections (e.g., sepsis, meningitis, septic arthritis, and/or osteomyelitis), autoimmune diseases (e.g., those disclosed herein), inflammatory disorders, and malignancies, and/or any disease or disorder or condition associated with these infections, diseases, disorders and/or malignancies) including, but not limited to, CVID, other primary immune deficiencies, HIV disease, CLL, recurrent bronchitis, sinusitis, otitis media, conjunctivitis, pneumonia, hepatitis, meningitis, herpes zoster (e.g., severe herpes zoster), and/or pneumocystis carinii.

In another embodiment, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention are used to treat, and/or diagnose an individual having common variable immunodeficiency disease ("CVID"; also known as "acquired agammaglobulinemia" and "acquired hypogammaglobulinemia") or a subset of this disease.

In a specific embodiment, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be used to treat, diagnose, and/or prevent (1) cancers or neoplasms and (2) autoimmune cell or tissue-related cancers or neoplasms. In a preferred embodiment, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention conjugated to a toxin or a radioactive isotope, as described herein, may be used to treat, diagnose, and/or prevent acute myelogenous leukemia. In a further preferred embodiment, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention conjugated to a toxin or a radioactive isotope, as described herein, may be used to treat, diagnose, and/or prevent, chronic myelogenous leukemia, multiple myeloma, non-Hodgkins lymphoma, and/or Hodgkins disease.

In another specific embodiment, polynucleotides or polypeptides, and/or agonists or antagonists of the invention may be used to treat, diagnose, prognose, and/or prevent selective IgA deficiency, myeloperoxidase deficiency, C2 deficiency, ataxia-telangiectasia, DiGeorge anomaly, common variable immunodeficiency (CVI),

X-linked agammaglobulinemia, severe combined immunodeficiency (SCID), chronic granulomatous disease (CGD), and Wiskott-Aldrich syndrome.

Examples of autoimmune disorders that can be treated or detected are described above and also include, but are not limited to: Addison's Disease, hemolytic anemia, antiphospholipid syndrome, rheumatoid arthritis, dermatitis, allergic encephalomyelitis, glomerulonephritis, Goodpasture's Syndrome, Graves' Disease, Multiple Sclerosis, Myasthenia Gravis, Neuritis, Ophthalmia, Bullous Pemphigoid, Pemphigus, Polyendocrinopathies, Purpura, Reiter's Disease, Stiff-Man Syndrome, Autoimmune Thyroiditis, Systemic Lupus Erythematosus, Autoimmune Pulmonary Inflammation, Guillain-Barre Syndrome, insulin dependent diabetes mellitis, and autoimmune inflammatory eye disease.

In a preferred embodiment, the autoimmune diseases and disorders and/or conditions associated with the diseases and disorders recited above are treated, prognosed, prevented, and/or diagnosed using antibodies against the polypeptide of the invention.

As an agent to boost immunoresponsiveness among B cell immunodeficient individuals, such as, for example, an individual who has undergone a partial or complete splenectomy.

Additionally, polynucleotides, polypeptides, and/or antagonists of the invention may affect apoptosis, and therefore, would be useful in treating a number of diseases associated with increased cell survival or the inhibition of apoptosis. For example, diseases associated with increased cell survival or the inhibition of apoptosis that could be treated or detected by polynucleotides, polypeptides, and/or antagonists of the invention, include cancers (such as follicular lymphomas, carcinomas with p53 mutations, and hormone-dependent tumors, including, but not limited to colon cancer, cardiac tumors, pancreatic cancer, melanoma, retinoblastoma, glioblastoma, lung cancer, intestinal cancer, testicular cancer, stomach cancer, neuroblastoma, myxoma, myoma, lymphoma, endothelioma, osteoblastoma, osteoclastoma, osteosarcoma, chondrosarcoma, adenoma, breast cancer, prostate cancer, Kaposi's sarcoma and ovarian cancer); autoimmune disorders (such as, multiple sclerosis, Sjogren's syndrome, Hashimoto's thyroiditis, biliary cirrhosis, Behcet's disease, Crohn's disease, polymyositis, systemic lupus erythematosus and immune-related

glomerulonephritis and rheumatoid arthritis) and viral infections (such as herpes viruses, pox viruses and adenoviruses), inflammation, graft v. host disease, acute graft rejection, and chronic graft rejection. In preferred embodiments, polynucleotides, polypeptides, and/or antagonists of the invention are used to inhibit growth, progression, and/or metastasis of cancers, in particular those listed above.

Additional diseases or conditions associated with increased cell survival that could be treated or detected by polynucleotides, polypeptides, and/or antagonists of the invention, include, but are not limited to, progression, and/or metastases of malignancies and related disorders such as leukemia (including acute leukemias (e.g., acute lymphocytic leukemia, acute myelocytic leukemia (including myeloblastic, promyelocytic, myelomonocytic, monocytic, and erythroleukemia)) and chronic leukemias (e.g., chronic myelocytic (granulocytic) leukemia and chronic lymphocytic leukemia)), polycythemia vera, lymphomas (e.g., Hodgkin's disease and non-Hodgkin's disease), multiple myeloma, Waldenstrom's macroglobulinemia, heavy chain disease, and solid tumors including, but not limited to, sarcomas and carcinomas such as fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilm's tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, menangioma, melanoma, neuroblastoma, and retinoblastoma.

Diseases associated with increased apoptosis that could be treated or detected by polynucleotides, polypeptides, and/or antagonists of the invention, include AIDS; neurodegenerative disorders (such as Alzheimer's disease, Parkinson's disease, Amyotrophic lateral sclerosis, Retinitis pigmentosa, Cerebellar degeneration and

brain tumor or prior associated disease); autoimmune disorders (such as, multiple sclerosis, Sjogren's syndrome, Hashimoto's thyroiditis, biliary cirrhosis, Behcet's disease, Crohn's disease, polymyositis, systemic lupus erythematosus and immune-related glomerulonephritis and rheumatoid arthritis) myelodysplastic syndromes (such as aplastic anemia), graft v. host disease, ischemic injury (such as that caused by myocardial infarction, stroke and reperfusion injury), liver injury (e.g., hepatitis related liver injury, ischemia/reperfusion injury, cholestasis (bile duct injury) and liver cancer); toxin-induced liver disease (such as that caused by alcohol), septic shock, cachexia and anorexia.

Hyperproliferative diseases and/or disorders that could be detected and/or treated by polynucleotides, polypeptides, and/or antagonists of the invention, include, but are not limited to neoplasms located in the: liver, abdomen, bone, breast, digestive system, pancreas, peritoneum, endocrine glands (adrenal, parathyroid, pituitary, testicles, ovary, thymus, thyroid), eye, head and neck, nervous (central and peripheral), lymphatic system, pelvic, skin, soft tissue, spleen, thoracic, and urogenital.

Similarly, other hyperproliferative disorders can also be treated or detected by polynucleotides, polypeptides, and/or antagonists of the invention. Examples of such hyperproliferative disorders include, but are not limited to:

hypergammaglobulinemia, lymphoproliferative disorders, paraproteinemias, purpura, sarcoidosis, Sezary Syndrome, Waldenström's Macroglobulinemia, Gaucher's Disease, histiocytosis, and any other hyperproliferative disease, besides neoplasia, located in an organ system listed above.

Hyperproliferative Disorders

A polynucleotides or polypeptides, or agonists or antagonists of the invention can be used to treat, prevent, and/or diagnose hyperproliferative diseases, disorders, and/or conditions, including neoplasms. A polynucleotides or polypeptides, or agonists or antagonists of the present invention may inhibit the proliferation of the disorder through direct or indirect interactions. Alternatively, a polynucleotides or polypeptides, or agonists or antagonists of the present invention may proliferate other cells which can inhibit the hyperproliferative disorder.

For example, by increasing an immune response, particularly increasing antigenic qualities of the hyperproliferative disorder or by proliferating, differentiating, or mobilizing T-cells, hyperproliferative diseases, disorders, and/or conditions can be treated, prevented, and/or diagnosed. This immune response may
5 be increased by either enhancing an existing immune response, or by initiating a new immune response. Alternatively, decreasing an immune response may also be a method of treating, preventing, and/or diagnosing hyperproliferative diseases, disorders, and/or conditions, such as a chemotherapeutic agent.

Examples of hyperproliferative diseases, disorders, and/or conditions that can
10 be treated, prevented, and/or diagnosed by polynucleotides or polypeptides, or agonists or antagonists of the present invention include, but are not limited to neoplasms located in the: colon, abdomen, bone, breast, digestive system, liver, pancreas, peritoneum, endocrine glands (adrenal, parathyroid, pituitary, testicles, ovary, thymus, thyroid), eye, head and neck, nervous (central and peripheral),
15 lymphatic system, pelvic, skin, soft tissue, spleen, thoracic, and urogenital.

Similarly, other hyperproliferative diseases, disorders, and/or conditions can also be treated, prevented, and/or diagnosed by a polynucleotides or polypeptides, or agonists or antagonists of the present invention. Examples of such hyperproliferative diseases, disorders, and/or conditions include, but are not limited to:
20 hypergammaglobulinemia, lymphoproliferative diseases, disorders, and/or conditions, paraproteinemias, purpura, sarcoidosis, Sezary Syndrome, Waldenstrom's Macroglobulinemia, Gaucher's Disease, histiocytosis, and any other hyperproliferative disease, besides neoplasia, located in an organ system listed above.

One preferred embodiment utilizes polynucleotides of the present invention to
25 inhibit aberrant cellular division, by gene therapy using the present invention, and/or protein fusions or fragments thereof.

Thus, the present invention provides a method for treating or preventing cell proliferative diseases, disorders, and/or conditions by inserting into an abnormally proliferating cell a polynucleotide of the present invention, wherein said
30 polynucleotide represses said expression.

Another embodiment of the present invention provides a method of treating or preventing cell-proliferative diseases, disorders, and/or conditions in individuals

comprising administration of one or more active gene copies of the present invention to an abnormally proliferating cell or cells. In a preferred embodiment, polynucleotides of the present invention is a DNA construct comprising a recombinant expression vector effective in expressing a DNA sequence encoding said polynucleotides. In another preferred embodiment of the present invention, the DNA construct encoding the polynucleotides of the present invention is inserted into cells to be treated utilizing a retrovirus, or more preferably an adenoviral vector (See G J. Nabel, et. al., PNAS 1999 96: 324-326, which is hereby incorporated by reference). In a most preferred embodiment, the viral vector is defective and will not transform non-proliferating cells, only proliferating cells. Moreover, in a preferred embodiment, the polynucleotides of the present invention inserted into proliferating cells either alone, or in combination with or fused to other polynucleotides, can then be modulated via an external stimulus (i.e. magnetic, specific small molecule, chemical, or drug administration, etc.), which acts upon the promoter upstream of said polynucleotides to induce expression of the encoded protein product. As such the beneficial therapeutic affect of the present invention may be expressly modulated (i.e. to increase, decrease, or inhibit expression of the present invention) based upon said external stimulus.

Polynucleotides of the present invention may be useful in repressing expression of oncogenic genes or antigens. By "repressing expression of the oncogenic genes " is intended the suppression of the transcription of the gene, the degradation of the gene transcript (pre-message RNA), the inhibition of splicing, the destruction of the messenger RNA, the prevention of the post-translational modifications of the protein, the destruction of the protein, or the inhibition of the normal function of the protein.

For local administration to abnormally proliferating cells, polynucleotides of the present invention may be administered by any method known to those of skill in the art including, but not limited to transfection, electroporation, microinjection of cells, or in vehicles such as liposomes, lipofectin, or as naked polynucleotides, or any other method described throughout the specification. The polynucleotide of the present invention may be delivered by known gene delivery systems such as, but not limited to, retroviral vectors (Gilboa, J. Virology 44:845 (1982); Hocke, Nature

320:275 (1986); Wilson, et al., Proc. Natl. Acad. Sci. U.S.A. 85:3014), vaccinia virus system (Chakrabarty et al., Mol. Cell Biol. 5:3403 (1985) or other efficient DNA delivery systems (Yates et al., Nature 313:812 (1985)) known to those skilled in the art. These references are exemplary only and are hereby incorporated by reference.

5 In order to specifically deliver or transfect cells which are abnormally proliferating and spare non-dividing cells, it is preferable to utilize a retrovirus, or adenoviral (as described in the art and elsewhere herein) delivery system known to those of skill in the art. Since host DNA replication is required for retroviral DNA to integrate and the retrovirus will be unable to self replicate due to the lack of the retrovirus genes
10 needed for its life cycle. Utilizing such a retroviral delivery system for polynucleotides of the present invention will target said gene and constructs to abnormally proliferating cells and will spare the non-dividing normal cells.

The polynucleotides of the present invention may be delivered directly to cell proliferative disorder/disease sites in internal organs, body cavities and the like by use
15 of imaging devices used to guide an injecting needle directly to the disease site. The polynucleotides of the present invention may also be administered to disease sites at the time of surgical intervention.

By "cell proliferative disease" is meant any human or animal disease or disorder, affecting any one or any combination of organs, cavities, or body parts,
20 which is characterized by single or multiple local abnormal proliferations of cells, groups of cells, or tissues, whether benign or malignant.

Any amount of the polynucleotides of the present invention may be administered as long as it has a biologically inhibiting effect on the proliferation of the treated cells. Moreover, it is possible to administer more than one of the
25 polynucleotide of the present invention simultaneously to the same site. By "biologically inhibiting" is meant partial or total growth inhibition as well as decreases in the rate of proliferation or growth of the cells. The biologically inhibitory dose may be determined by assessing the effects of the polynucleotides of the present invention on target malignant or abnormally proliferating cell growth in
30 tissue culture, tumor growth in animals and cell cultures, or any other method known to one of ordinary skill in the art.

The present invention is further directed to antibody-based therapies which involve administering of anti-polypeptides and anti-polynucleotide antibodies to a mammalian, preferably human, patient for treating, preventing, and/or diagnosing one or more of the described diseases, disorders, and/or conditions. Methods for
5 producing anti-polypeptides and anti-polynucleotide antibodies polyclonal and monoclonal antibodies are described in detail elsewhere herein. Such antibodies may be provided in pharmaceutically acceptable compositions as known in the art or as described herein.

A summary of the ways in which the antibodies of the present invention may
10 be used therapeutically includes binding polynucleotides or polypeptides of the present invention locally or systemically in the body or by direct cytotoxicity of the antibody, e.g. as mediated by complement (CDC) or by effector cells (ADCC). Some of these approaches are described in more detail below. Armed with the teachings provided herein, one of ordinary skill in the art will know how to use the antibodies of
15 the present invention for diagnostic, monitoring or therapeutic purposes without undue experimentation.

In particular, the antibodies, fragments and derivatives of the present invention are useful for treating, preventing, and/or diagnosing a subject having or developing cell proliferative and/or differentiation diseases, disorders, and/or conditions as
20 described herein. Such treatment comprises administering a single or multiple doses of the antibody, or a fragment, derivative, or a conjugate thereof.

The antibodies of this invention may be advantageously utilized in combination with other monoclonal or chimeric antibodies, or with lymphokines or hematopoietic growth factors, for example, which serve to increase the number or
25 activity of effector cells which interact with the antibodies.

It is preferred to use high affinity and/or potent in vivo inhibiting and/or neutralizing antibodies against polypeptides or polynucleotides of the present invention, fragments or regions thereof, for both immunoassays directed to and therapy of diseases, disorders, and/or conditions related to polynucleotides or
30 polypeptides, including fragments thereof, of the present invention. Such antibodies, fragments, or regions, will preferably have an affinity for polynucleotides or polypeptides, including fragments thereof. Preferred binding affinities include those

with a dissociation constant or K_d less than $5 \times 10^{-6}M$, $10^{-6}M$, $5 \times 10^{-7}M$, $10^{-7}M$, $5 \times 10^{-8}M$, $10^{-8}M$, $5 \times 10^{-9}M$, $10^{-9}M$, $5 \times 10^{-10}M$, $10^{-10}M$, $5 \times 10^{-11}M$, $10^{-11}M$, $5 \times 10^{-12}M$, $10^{-12}M$, $5 \times 10^{-13}M$, $10^{-13}M$, $5 \times 10^{-14}M$, $10^{-14}M$, $5 \times 10^{-15}M$, and $10^{-15}M$.

Moreover, polypeptides of the present invention are useful in inhibiting the angiogenesis of proliferative cells or tissues, either alone, as a protein fusion, or in combination with other polypeptides directly or indirectly, as described elsewhere herein. In a most preferred embodiment, said anti-angiogenesis effect may be achieved indirectly, for example, through the inhibition of hematopoietic, tumor-specific cells, such as tumor-associated macrophages (See Joseph IB, et al. J Natl Cancer Inst, 90(21):1648-53 (1998), which is hereby incorporated by reference). Antibodies directed to polypeptides or polynucleotides of the present invention may also result in inhibition of angiogenesis directly, or indirectly (See Witte L, et al., Cancer Metastasis Rev. 17(2):155-61 (1998), which is hereby incorporated by reference)).

Polypeptides, including protein fusions, of the present invention, or fragments thereof may be useful in inhibiting proliferative cells or tissues through the induction of apoptosis. Said polypeptides may act either directly, or indirectly to induce apoptosis of proliferative cells and tissues, for example in the activation of a death-domain receptor, such as tumor necrosis factor (TNF) receptor-1, CD95 (Fas/APO-1), TNF-receptor-related apoptosis-mediated protein (TRAMP) and TNF-related apoptosis-inducing ligand (TRAIL) receptor-1 and -2 (See Schulze-Osthoff K, et.al., Eur J Biochem 254(3):439-59 (1998), which is hereby incorporated by reference). Moreover, in another preferred embodiment of the present invention, said polypeptides may induce apoptosis through other mechanisms, such as in the activation of other proteins which will activate apoptosis, or through stimulating the expression of said proteins, either alone or in combination with small molecule drugs or adjuvants, such as apoptonin, galectins, thioredoxins, antiinflammatory proteins (See for example, Mutat Res 400(1-2):447-55 (1998), Med Hypotheses.50(5):423-33 (1998), Chem Biol Interact. Apr 24;111-112:23-34 (1998), J Mol Med.76(6):402-12 (1998), Int J Tissue React;20(1):3-15 (1998), which are all hereby incorporated by reference).

Polypeptides, including protein fusions to, or fragments thereof, of the present invention are useful in inhibiting the metastasis of proliferative cells or tissues. Inhibition may occur as a direct result of administering polypeptides, or antibodies directed to said polypeptides as described elsewhere herein, or indirectly, such as
5 activating the expression of proteins known to inhibit metastasis, for example alpha 4 integrins, (See, e.g., Curr Top Microbiol Immunol 1998;231:125-41, which is hereby incorporated by reference). Such therapeutic affects of the present invention may be achieved either alone, or in combination with small molecule drugs or adjuvants.

In another embodiment, the invention provides a method of delivering
10 compositions containing the polypeptides of the invention (e.g., compositions containing polypeptides or polypeptide antibodies associated with heterologous polypeptides, heterologous nucleic acids, toxins, or prodrugs) to targeted cells expressing the polypeptide of the present invention. Polypeptides or polypeptide antibodies of the invention may be associated with with heterologous polypeptides,
15 heterologous nucleic acids, toxins, or prodrugs via hydrophobic, hydrophilic, ionic and/or covalent interactions.

Polypeptides, protein fusions to, or fragments thereof, of the present invention are useful in enhancing the immunogenicity and/or antigenicity of proliferating cells or tissues, either directly, such as would occur if the polypeptides of the present
20 invention 'vaccinated' the immune response to respond to proliferative antigens and immunogens, or indirectly, such as in activating the expression of proteins known to enhance the immune response (e.g. chemokines), to said antigens and immunogens.

Cardiovascular Disorders

25 Polynucleotides or polypeptides, or agonists or antagonists of the invention may be used to treat, prevent, and/or diagnose cardiovascular diseases, disorders, and/or conditions, including peripheral artery disease, such as limb ischemia.

Cardiovascular diseases, disorders, and/or conditions include cardiovascular abnormalities, such as arterio-arterial fistula, arteriovenous fistula, cerebral
30 arteriovenous malformations, congenital heart defects, pulmonary atresia, and Scimitar Syndrome. Congenital heart defects include aortic coarctation, ..cor triatriatum, coronary vessel anomalies, crisscross heart, dextrocardia, patent ductus

arteriosus, Ebstein's anomaly, Eisenmenger complex, hypoplastic left heart syndrome, levocardia, tetralogy of fallot, transposition of great vessels, double outlet right ventricle, tricuspid atresia, persistent truncus arteriosus, and heart septal defects, such as aortopulmonary septal defect, endocardial cushion defects, Lutembacher's
5 Syndrome, trilog of Fallot, ventricular heart septal defects.

Cardiovascular diseases, disorders, and/or conditions also include heart disease, such as arrhythmias, carcinoid heart disease, high cardiac output, low cardiac output, cardiac tamponade, endocarditis (including bacterial), heart aneurysm, cardiac arrest, congestive heart failure, congestive cardiomyopathy, paroxysmal dyspnea,
10 cardiac edema, heart hypertrophy, congestive cardiomyopathy, left ventricular hypertrophy, right ventricular hypertrophy, post-infarction heart rupture, ventricular septal rupture, heart valve diseases, myocardial diseases, myocardial ischemia, pericardial effusion, pericarditis (including constrictive and tuberculous), pneumopericardium, postpericardiotomy syndrome, pulmonary heart disease,
15 rheumatic heart disease, ventricular dysfunction, hyperemia, cardiovascular pregnancy complications, Scimitar Syndrome, cardiovascular syphilis, and cardiovascular tuberculosis.

Arrhythmias include sinus arrhythmia, atrial fibrillation, atrial flutter, bradycardia, extrasystole, Adams-Stokes Syndrome, bundle-branch block, sinoatrial
20 block, long QT syndrome, parasystole, Lown-Ganong-Levine Syndrome, Mahaim-type pre-excitation syndrome, Wolff-Parkinson-White syndrome, sick sinus syndrome, tachycardias, and ventricular fibrillation. Tachycardias include paroxysmal tachycardia, supraventricular tachycardia, accelerated idioventricular rhythm, atrioventricular nodal reentry tachycardia, ectopic atrial tachycardia, ectopic
25 junctional tachycardia, sinoatrial nodal reentry tachycardia, sinus tachycardia, Torsades de Pointes, and ventricular tachycardia.

Heart valve disease include aortic valve insufficiency, aortic valve stenosis, hear murmurs, aortic valve prolapse, mitral valve prolapse, tricuspid valve prolapse, mitral valve insufficiency, mitral valve stenosis, pulmonary atresia, pulmonary valve
30 insufficiency, pulmonary valve stenosis, tricuspid atresia, tricuspid valve insufficiency, and tricuspid valve stenosis.

Myocardial diseases include alcoholic cardiomyopathy, congestive cardiomyopathy, hypertrophic cardiomyopathy, aortic subvalvular stenosis, pulmonary subvalvular stenosis, restrictive cardiomyopathy, Chagas cardiomyopathy, endocardial fibroelastosis, endomyocardial fibrosis, Kearns Syndrome, myocardial
5 reperfusion injury, and myocarditis.

Myocardial ischemias include coronary disease, such as angina pectoris, coronary aneurysm, coronary arteriosclerosis, coronary thrombosis, coronary vasospasm, myocardial infarction and myocardial stunning.

Cardiovascular diseases also include vascular diseases such as aneurysms,
10 angiodyplasia, angiomas, bacillary angiomas, Hippel-Lindau Disease, Klippel-Trenaunay-Weber Syndrome, Sturge-Weber Syndrome, angioneurotic edema, aortic diseases, Takayasu's Arteritis, aortitis, Leriche's Syndrome, arterial occlusive diseases, arteritis, enarteritis, polyarteritis nodosa, cerebrovascular diseases, disorders, and/or conditions, diabetic angiopathies, diabetic retinopathy, embolisms, thrombosis,
15 erythromelalgia, hemorrhoids, hepatic veno-occlusive disease, hypertension, hypotension, ischemia, peripheral vascular diseases, phlebitis, pulmonary veno-occlusive disease, Raynaud's disease, CREST syndrome, retinal vein occlusion, Scimitar syndrome, superior vena cava syndrome, telangiectasia, ataxia telangiectasia, hereditary hemorrhagic telangiectasia, varicocele, varicose veins, varicose ulcer,
20 vasculitis, and venous insufficiency.

Aneurysms include dissecting aneurysms, false aneurysms, infected aneurysms, ruptured aneurysms, aortic aneurysms, cerebral aneurysms, coronary aneurysms, heart aneurysms, and iliac aneurysms.

Arterial occlusive diseases include arteriosclerosis, intermittent claudication,
25 carotid stenosis, fibromuscular dysplasias, mesenteric vascular occlusion, Moyamoya disease, renal artery obstruction, retinal artery occlusion, and thromboangiitis obliterans.

Cerebrovascular diseases, disorders, and/or conditions include carotid artery diseases, cerebral amyloid angiopathy, cerebral aneurysm, cerebral anoxia, cerebral
30 arteriosclerosis, cerebral arteriovenous malformation, cerebral artery diseases, cerebral embolism and thrombosis, carotid artery thrombosis, sinus thrombosis, Wallenberg's syndrome, cerebral hemorrhage, epidural hematoma, subdural

hematoma, subarachnoid hemorrhage, cerebral infarction, cerebral ischemia (including transient), subclavian steal syndrome, periventricular leukomalacia, vascular headache, cluster headache, migraine, and vertebrobasilar insufficiency.

Embolisms include air embolisms, amniotic fluid embolisms, cholesterol
5 embolisms, blue toe syndrome, fat embolisms, pulmonary embolisms, and thromboembolisms. Thrombosis include coronary thrombosis, hepatic vein thrombosis, retinal vein occlusion, carotid artery thrombosis, sinus thrombosis, Wallenberg's syndrome, and thrombophlebitis.

Ischemia includes cerebral ischemia, ischemic colitis, compartment
10 syndromes, anterior compartment syndrome, myocardial ischemia, reperfusion injuries, and peripheral limb ischemia. Vasculitis includes aortitis, arteritis, Behcet's Syndrome, Churg-Strauss Syndrome, mucocutaneous lymph node syndrome, thromboangiitis obliterans, hypersensitivity vasculitis, Schoenlein-Henoch purpura, allergic cutaneous vasculitis, and Wegener's granulomatosis.

15 Polynucleotides or polypeptides, or agonists or antagonists of the invention, are especially effective for the treatment of critical limb ischemia and coronary disease.

Polypeptides may be administered using any method known in the art, including, but not limited to, direct needle injection at the delivery site, intravenous
20 injection, topical administration, catheter infusion, biolistic injectors, particle accelerators, gelfoam sponge depots, other commercially available depot materials, osmotic pumps, oral or suppositorial solid pharmaceutical formulations, decanting or topical applications during surgery, aerosol delivery. Such methods are known in the art. Polypeptides of the invention may be administered as part of a *Therapeutic*,
25 described in more detail below. Methods of delivering polynucleotides of the invention are described in more detail herein..

Anti-Angiogenesis Activity

The naturally occurring balance between endogenous stimulators and
30 inhibitors of angiogenesis is one in which inhibitory influences predominate. Rastinejad *et al.*, *Cell* 56:345-355 (1989). In those rare instances in which neovascularization occurs under normal physiological conditions, such as wound

healing, organ regeneration, embryonic development, and female reproductive processes, angiogenesis is stringently regulated and spatially and temporally delimited. Under conditions of pathological angiogenesis such as that characterizing solid tumor growth, these regulatory controls fail. Unregulated angiogenesis becomes pathologic and sustains progression of many neoplastic and non-neoplastic diseases. A number of serious diseases are dominated by abnormal neovascularization including solid tumor growth and metastases, arthritis, some types of eye diseases, disorders, and/or conditions, and psoriasis. See, e.g., reviews by Moses *et al.*, *Biotech.* 9:630-634 (1991); Folkman *et al.*, *N. Engl. J. Med.*, 333:1757-1763 (1995); Auerbach *et al.*, *J. Microvasc. Res.* 29:401-411 (1985); Folkman, *Advances in Cancer Research*, eds. Klein and Weinhouse, Academic Press, New York, pp. 175-203 (1985); Patz, *Am. J. Ophthalmol.* 94:715-743 (1982); and Folkman *et al.*, *Science* 221:719-725 (1983). In a number of pathological conditions, the process of angiogenesis contributes to the disease state. For example, significant data have accumulated which suggest that the growth of solid tumors is dependent on angiogenesis. Folkman and Klagsbrun, *Science* 235:442-447 (1987).

The present invention provides for treatment of diseases, disorders, and/or conditions associated with neovascularization by administration of the polynucleotides and/or polypeptides of the invention, as well as agonists or antagonists of the present invention. Malignant and metastatic conditions which can be treated with the polynucleotides and polypeptides, or agonists or antagonists of the invention include, but are not limited to, malignancies, solid tumors, and cancers described herein and otherwise known in the art (for a review of such disorders, see Fishman *et al.*, *Medicine*, 2d Ed., J. B. Lippincott Co., Philadelphia (1985)). Thus, the present invention provides a method of treating, preventing, and/or diagnosing an angiogenesis-related disease and/or disorder, comprising administering to an individual in need thereof a therapeutically effective amount of a polynucleotide, polypeptide, antagonist and/or agonist of the invention. For example, polynucleotides, polypeptides, antagonists and/or agonists may be utilized in a variety of additional methods in order to therapeutically treat or prevent a cancer or tumor. Cancers which may be treated, prevented, and/or diagnosed with polynucleotides, polypeptides, antagonists and/or agonists include, but are not limited to solid tumors,

including prostate, lung, breast, ovarian, stomach, pancreas, larynx, esophagus, testes, liver, parotid, biliary tract, colon, rectum, cervix, uterus, endometrium, kidney, bladder, thyroid cancer; primary tumors and metastases; melanomas; glioblastoma; Kaposi's sarcoma; leiomyosarcoma; non-small cell lung cancer; colorectal cancer; advanced malignancies; and blood born tumors such as leukemias. For example, polynucleotides, polypeptides, antagonists and/or agonists may be delivered topically, in order to treat or prevent cancers such as skin cancer, head and neck tumors, breast tumors, and Kaposi's sarcoma.

Within yet other aspects, polynucleotides, polypeptides, antagonists and/or agonists may be utilized to treat superficial forms of bladder cancer by, for example, intravesical administration. Polynucleotides, polypeptides, antagonists and/or agonists may be delivered directly into the tumor, or near the tumor site, via injection or a catheter. Of course, as the artisan of ordinary skill will appreciate, the appropriate mode of administration will vary according to the cancer to be treated. Other modes of delivery are discussed herein.

Polynucleotides, polypeptides, antagonists and/or agonists may be useful in treating, preventing, and/or diagnosing other diseases, disorders, and/or conditions, besides cancers, which involve angiogenesis. These diseases, disorders, and/or conditions include, but are not limited to: benign tumors, for example hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas; arteriosclerotic plaques; ocular angiogenic diseases, for example, diabetic retinopathy, retinopathy of prematurity, macular degeneration, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, rubeosis, retinoblastoma, uveitis and Pterygia (abnormal blood vessel growth) of the eye; rheumatoid arthritis; psoriasis; delayed wound healing; endometriosis; vasculogenesis; granulations; hypertrophic scars (keloids); nonunion fractures; scleroderma; trachoma; vascular adhesions; myocardial angiogenesis; coronary collaterals; cerebral collaterals; arteriovenous malformations; ischemic limb angiogenesis; Osler-Webber Syndrome; plaque neovascularization; telangiectasia; hemophilic joints; angiofibroma; fibromuscular dysplasia; wound granulation; Crohn's disease; and atherosclerosis.

For example, within one aspect of the present invention methods are provided for treating, preventing, and/or diagnosing hypertrophic scars and keloids, comprising

the step of administering a polynucleotide, polypeptide, antagonist and/or agonist of the invention to a hypertrophic scar or keloid.

Within one embodiment of the present invention polynucleotides, polypeptides, antagonists and/or agonists are directly injected into a hypertrophic scar or keloid, in order to prevent the progression of these lesions. This therapy is of particular value in the prophylactic treatment of conditions which are known to result in the development of hypertrophic scars and keloids (e.g., burns), and is preferably initiated after the proliferative phase has had time to progress (approximately 14 days after the initial injury), but before hypertrophic scar or keloid development. As noted above, the present invention also provides methods for treating, preventing, and/or diagnosing neovascular diseases of the eye, including for example, corneal neovascularization, neovascular glaucoma, proliferative diabetic retinopathy, retrolental fibroplasia and macular degeneration.

Moreover, Ocular diseases, disorders, and/or conditions associated with neovascularization which can be treated, prevented, and/or diagnosed with the polynucleotides and polypeptides of the present invention (including agonists and/or antagonists) include, but are not limited to: neovascular glaucoma, diabetic retinopathy, retinoblastoma, retrolental fibroplasia, uveitis, retinopathy of prematurity macular degeneration, corneal graft neovascularization, as well as other eye inflammatory diseases, ocular tumors and diseases associated with choroidal or iris neovascularization. See, e.g., reviews by Waltman *et al.*, *Am. J. Ophthalmol.* 85:704-710 (1978) and Gartner *et al.*, *Surv. Ophthalmol.* 22:291-312 (1978).

Thus, within one aspect of the present invention methods are provided for treating or preventing neovascular diseases of the eye such as corneal neovascularization (including corneal graft neovascularization), comprising the step of administering to a patient a therapeutically effective amount of a compound (as described above) to the cornea, such that the formation of blood vessels is inhibited. Briefly, the cornea is a tissue which normally lacks blood vessels. In certain pathological conditions however, capillaries may extend into the cornea from the pericorneal vascular plexus of the limbus. When the cornea becomes vascularized, it also becomes clouded, resulting in a decline in the patient's visual acuity. Visual loss may become complete if the cornea completely opacitates. A wide variety of

diseases, disorders, and/or conditions can result in corneal neovascularization, including for example, corneal infections (e.g., trachoma, herpes simplex keratitis, leishmaniasis and onchocerciasis), immunological processes (e.g., graft rejection and Stevens-Johnson's syndrome), alkali burns, trauma, inflammation (of any cause),
5 toxic and nutritional deficiency states, and as a complication of wearing contact lenses.

Within particularly preferred embodiments of the invention, may be prepared for topical administration in saline (combined with any of the preservatives and antimicrobial agents commonly used in ocular preparations), and administered in
10 eyedrop form. The solution or suspension may be prepared in its pure form and administered several times daily. Alternatively, anti-angiogenic compositions, prepared as described above, may also be administered directly to the cornea. Within preferred embodiments, the anti-angiogenic composition is prepared with a muco-adhesive polymer which binds to cornea. Within further embodiments, the anti-
15 angiogenic factors or anti-angiogenic compositions may be utilized as an adjunct to conventional steroid therapy. Topical therapy may also be useful prophylactically in corneal lesions which are known to have a high probability of inducing an angiogenic response (such as chemical burns). In these instances the treatment, likely in combination with steroids, may be instituted immediately to help prevent subsequent
20 complications.

Within other embodiments, the compounds described above may be injected directly into the corneal stroma by an ophthalmologist under microscopic guidance. The preferred site of injection may vary with the morphology of the individual lesion, but the goal of the administration would be to place the composition at the advancing
25 front of the vasculature (i.e., interspersed between the blood vessels and the normal cornea). In most cases this would involve perilimbic corneal injection to "protect" the cornea from the advancing blood vessels. This method may also be utilized shortly after a corneal insult in order to prophylactically prevent corneal neovascularization. In this situation the material could be injected in the perilimbic cornea interspersed
30 between the corneal lesion and its undesired potential limbic blood supply. Such methods may also be utilized in a similar fashion to prevent capillary invasion of transplanted corneas. In a sustained-release form injections might only be required 2-

3 times per year. A steroid could also be added to the injection solution to reduce inflammation resulting from the injection itself.

Within another aspect of the present invention, methods are provided for treating or preventing neovascular glaucoma, comprising the step of administering to
5 a patient a therapeutically effective amount of a polynucleotide, polypeptide, antagonist and/or agonist to the eye, such that the formation of blood vessels is inhibited. In one embodiment, the compound may be administered topically to the eye in order to treat or prevent early forms of neovascular glaucoma. Within other embodiments, the compound may be implanted by injection into the region of the
10 anterior chamber angle. Within other embodiments, the compound may also be placed in any location such that the compound is continuously released into the aqueous humor. Within another aspect of the present invention, methods are provided for treating or preventing proliferative diabetic retinopathy, comprising the step of administering to a patient a therapeutically effective amount of a polynucleotide,
15 polypeptide, antagonist and/or agonist to the eyes, such that the formation of blood vessels is inhibited.

Within particularly preferred embodiments of the invention, proliferative diabetic retinopathy may be treated by injection into the aqueous humor or the vitreous, in order to increase the local concentration of the polynucleotide,
20 polypeptide, antagonist and/or agonist in the retina. Preferably, this treatment should be initiated prior to the acquisition of severe disease requiring photocoagulation.

Within another aspect of the present invention, methods are provided for treating or preventing retrolental fibroplasia, comprising the step of administering to a patient a therapeutically effective amount of a polynucleotide, polypeptide, antagonist
25 and/or agonist to the eye, such that the formation of blood vessels is inhibited. The compound may be administered topically, via intravitreal injection and/or via intraocular implants.

Additionally, diseases, disorders, and/or conditions which can be treated, prevented, and/or diagnosed with the polynucleotides, polypeptides, agonists and/or
30 agonists include, but are not limited to, hemangioma, arthritis, psoriasis, angiofibroma, atherosclerotic plaques, delayed wound healing, granulations, hemophilic joints, hypertrophic scars, nonunion fractures, Osler-Weber syndrome,

pyogenic granuloma, scleroderma, trachoma, and vascular adhesions.

Moreover, diseases, disorders, and/or conditions and/or states, which can be treated, prevented, and/or diagnosed with the the polynucleotides, polypeptides, agonists and/or agonists include, but are not limited to, solid tumors, blood born
5 tumors such as leukemias, tumor metastasis, Kaposi's sarcoma, benign tumors, for example hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas, rheumatoid arthritis, psoriasis, ocular angiogenic diseases, for example, diabetic retinopathy, retinopathy of prematurity, macular degeneration, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, rubeosis, retinoblastoma, and
10 uvietis, delayed wound healing, endometriosis, vasculogenesis, granulations, hypertrophic scars (keloids), nonunion fractures, scleroderma, trachoma, vascular adhesions, myocardial angiogenesis, coronary collaterals, cerebral collaterals, arteriovenous malformations, ischemic limb angiogenesis, Osler-Webber Syndrome, plaque neovascularization, telangiectasia, hemophiliac joints, angiofibroma
15 fibromuscular dysplasia, wound granulation, Crohn's disease, atherosclerosis, birth control agent by preventing vascularization required for embryo implantation controlling menstruation, diseases that have angiogenesis as a pathologic consequence such as cat scratch disease (*Rochela minalia quintosa*), ulcers (*Helicobacter pylori*), Bartonellosis and bacillary angiomatosis.

20 In one aspect of the birth control method, an amount of the compound sufficient to block embryo implantation is administered before or after intercourse and fertilization have occurred, thus providing an effective method of birth control, possibly a "morning after" method. Polynucleotides, polypeptides, agonists and/or agonists may also be used in controlling menstruation or administered as either a
25 peritoneal lavage fluid or for peritoneal implantation in the treatment of endometriosis.

Polynucleotides, polypeptides, agonists and/or agonists of the present invention may be incorporated into surgical sutures in order to prevent stitch granulomas.

30 Polynucleotides, polypeptides, agonists and/or agonists may be utilized in a wide variety of surgical procedures. For example, within one aspect of the present invention a compositions (in the form of, for example, a spray or film) may be utilized

to coat or spray an area prior to removal of a tumor, in order to isolate normal surrounding tissues from malignant tissue, and/or to prevent the spread of disease to surrounding tissues. Within other aspects of the present invention, compositions (e.g., in the form of a spray) may be delivered via endoscopic procedures in order to coat
5 tumors, or inhibit angiogenesis in a desired locale. Within yet other aspects of the present invention, surgical meshes which have been coated with anti- angiogenic compositions of the present invention may be utilized in any procedure wherein a surgical mesh might be utilized. For example, within one embodiment of the invention a surgical mesh laden with an anti-angiogenic composition may be utilized
10 during abdominal cancer resection surgery (e.g., subsequent to colon resection) in order to provide support to the structure, and to release an amount of the anti-angiogenic factor.

Within further aspects of the present invention, methods are provided for treating tumor excision sites, comprising administering a polynucleotide, polypeptide,
15 agonist and/or agonist to the resection margins of a tumor subsequent to excision, such that the local recurrence of cancer and the formation of new blood vessels at the site is inhibited. Within one embodiment of the invention, the anti-angiogenic compound is administered directly to the tumor excision site (e.g., applied by swabbing, brushing or otherwise coating the resection margins of the tumor with the
20 anti-angiogenic compound). Alternatively, the anti-angiogenic compounds may be incorporated into known surgical pastes prior to administration. Within particularly preferred embodiments of the invention, the anti-angiogenic compounds are applied after hepatic resections for malignancy, and after neurosurgical operations.

Within one aspect of the present invention, polynucleotides, polypeptides,
25 agonists and/or agonists may be administered to the resection margin of a wide variety of tumors, including for example, breast, colon, brain and hepatic tumors. For example, within one embodiment of the invention, anti-angiogenic compounds may be administered to the site of a neurological tumor subsequent to excision, such that the formation of new blood vessels at the site are inhibited.

30 The polynucleotides, polypeptides, agonists and/or agonists of the present invention may also be administered along with other anti-angiogenic factors. Representative examples of other anti-angiogenic factors include: Anti-Invasive

Factor, retinoic acid and derivatives thereof, paclitaxel, Suramin, Tissue Inhibitor of Metalloproteinase-1, Tissue Inhibitor of Metalloproteinase-2, Plasminogen Activator Inhibitor-1, Plasminogen Activator Inhibitor-2, and various forms of the lighter "d group" transition metals.

- 5 Lighter "d group" transition metals include, for example, vanadium, molybdenum, tungsten, titanium, niobium, and tantalum species. Such transition metal species may form transition metal complexes. Suitable complexes of the above-mentioned transition metal species include oxo transition metal complexes.

10 Representative examples of vanadium complexes include oxo vanadium complexes such as vanadate and vanadyl complexes. Suitable vanadate complexes include metavanadate and orthovanadate complexes such as, for example, ammonium metavanadate, sodium metavanadate, and sodium orthovanadate. Suitable vanadyl complexes include, for example, vanadyl acetylacetonate and vanadyl sulfate including vanadyl sulfate hydrates such as vanadyl sulfate mono- and trihydrates.

- 15 Representative examples of tungsten and molybdenum complexes also include oxo complexes. Suitable oxo tungsten complexes include tungstate and tungsten oxide complexes. Suitable tungstate complexes include ammonium tungstate, calcium tungstate, sodium tungstate dihydrate, and tungstic acid. Suitable tungsten oxides include tungsten (IV) oxide and tungsten (VI) oxide. Suitable oxo
20 molybdenum complexes include molybdate, molybdenum oxide, and molybdenyl complexes. Suitable molybdate complexes include ammonium molybdate and its hydrates, sodium molybdate and its hydrates, and potassium molybdate and its hydrates. Suitable molybdenum oxides include molybdenum (VI) oxide, molybdenum (VI) oxide, and molybdic acid. Suitable molybdenyl complexes include, for example,
25 molybdenyl acetylacetonate. Other suitable tungsten and molybdenum complexes include hydroxo derivatives derived from, for example, glycerol, tartaric acid, and sugars.

 A wide variety of other anti-angiogenic factors may also be utilized within the context of the present invention. Representative examples include platelet factor 4;
30 protamine sulphate; sulphated chitin derivatives (prepared from queen crab shells), (Murata et al., Cancer Res. 51:22-26, 1991); Sulphated Polysaccharide Peptidoglycan Complex (SP- PG) (the function of this compound may be enhanced by the presence

of steroids such as estrogen, and tamoxifen citrate); Staurosporine; modulators of matrix metabolism, including for example, proline analogs, cishydroxyproline, d,L-3,4-dehydroproline, Thiaproline, alpha,alpha-dipyridyl, aminopropionitrile fumarate; 4-propyl-5-(4-pyridinyl)-2(3H)-oxazolone; Methotrexate; Mitoxantrone; Heparin; 5 Interferons; 2 Macroglobulin-serum; ChIMP-3 (Pavloff et al., J. Bio. Chem. 267:17321-17326, 1992); Chymostatin (Tomkinson et al., Biochem J. 286:475-480, 1992); Cyclodextrin Tetradasulfate; Eponemycin; Camptothecin; Fumagillin (Ingber et al., Nature 348:555-557, 1990); Gold Sodium Thiomalate ("GST"; Matsubara and Ziff, J. Clin. Invest. 79:1440-1446, 1987); anticollagenase-serum; 10 alpha2-antiplasmin (Holmes et al., J. Biol. Chem. 262(4):1659-1664, 1987); Bisantrene (National Cancer Institute); Lobenzarit disodium (N-(2)-carboxyphenyl-4-chloroanthronilic acid disodium or "CCA"; Takeuchi et al., Agents Actions 36:312-316, 1992); Thalidomide; Angostatic steroid; AGM-1470; carboxynaminolimidazole; and metalloproteinase inhibitors such as BB94.

15

Diseases at the Cellular Level

Diseases associated with increased cell survival or the inhibition of apoptosis that could be treated, prevented, and/or diagnosed by the polynucleotides or polypeptides and/or antagonists or agonists of the invention, include cancers (such as 20 follicular lymphomas, carcinomas with p53 mutations, and hormone-dependent tumors, including, but not limited to colon cancer, cardiac tumors, pancreatic cancer, melanoma, retinoblastoma, glioblastoma, lung cancer, intestinal cancer, testicular cancer, stomach cancer, neuroblastoma, myxoma, myoma, lymphoma, endothelioma, osteoblastoma, osteoclastoma, osteosarcoma, chondrosarcoma, adenoma, breast 25 cancer, prostate cancer, Kaposi's sarcoma and ovarian cancer); autoimmune diseases, disorders, and/or conditions (such as, multiple sclerosis, Sjogren's syndrome, Hashimoto's thyroiditis, biliary cirrhosis, Behcet's disease, Crohn's disease, polymyositis, systemic lupus erythematosus and immune-related glomerulonephritis and rheumatoid arthritis) and viral infections (such as herpes viruses, pox viruses and 30 adenoviruses), inflammation, graft v. host disease, acute graft rejection, and chronic graft rejection. In preferred embodiments, the polynucleotides or polypeptides, and/or

agonists or antagonists of the invention are used to inhibit growth, progression, and/or metasis of cancers, in particular those listed above.

Additional diseases or conditions associated with increased cell survival that could be treated, prevented or diagnosed by the polynucleotides or polypeptides, or agonists or antagonists of the invention, include, but are not limited to, progression, and/or metastases of malignancies and related disorders such as leukemia (including acute leukemias (e.g., acute lymphocytic leukemia, acute myelocytic leukemia (including myeloblastic, promyelocytic, myelomonocytic, monocytic, and erythroleukemia)) and chronic leukemias (e.g., chronic myelocytic (granulocytic) leukemia and chronic lymphocytic leukemia)), polycythemia vera, lymphomas (e.g., Hodgkin's disease and non-Hodgkin's disease), multiple myeloma, Waldenstrom's macroglobulinemia, heavy chain disease, and solid tumors including, but not limited to, sarcomas and carcinomas such as fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilm's tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, menangioma, melanoma, neuroblastoma, and retinoblastoma.

Diseases associated with increased apoptosis that could be treated, prevented, and/or diagnosed by the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, include AIDS; neurodegenerative diseases, disorders, and/or conditions (such as Alzheimer's disease, Parkinson's disease, Amyotrophic lateral sclerosis, Retinitis pigmentosa, Cerebellar degeneration and brain tumor or prior associated disease); autoimmune diseases, disorders, and/or conditions (such as, multiple sclerosis, Sjogren's syndrome, Hashimoto's thyroiditis, biliary cirrhosis,

Behcet's disease, Crohn's disease, polymyositis, systemic lupus erythematosus and immune-related glomerulonephritis and rheumatoid arthritis) myelodysplastic syndromes (such as aplastic anemia), graft v. host disease, ischemic injury (such as that caused by myocardial infarction, stroke and reperfusion injury), liver injury (e.g.,
5 hepatitis related liver injury, ischemia/reperfusion injury, cholestasis (bile duct injury) and liver cancer); toxin-induced liver disease (such as that caused by alcohol), septic shock, cachexia and anorexia.

Wound Healing and Epithelial Cell Proliferation

10 In accordance with yet a further aspect of the present invention, there is provided a process for utilizing the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, for therapeutic purposes, for example, to stimulate epithelial cell proliferation and basal keratinocytes for the purpose of wound healing, and to stimulate hair follicle production and healing of dermal wounds.
15 Polynucleotides or polypeptides, as well as agonists or antagonists of the invention, may be clinically useful in stimulating wound healing including surgical wounds, excisional wounds, deep wounds involving damage of the dermis and epidermis, eye tissue wounds, dental tissue wounds, oral cavity wounds, diabetic ulcers, dermal ulcers, cubitus ulcers, arterial ulcers, venous stasis ulcers, burns resulting from heat
20 exposure or chemicals, and other abnormal wound healing conditions such as uremia, malnutrition, vitamin deficiencies and complications associated with systemic treatment with steroids, radiation therapy and antineoplastic drugs and antimetabolites. Polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to promote dermal reestablishment subsequent to dermal
25 loss

The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to increase the adherence of skin grafts to a wound bed and to stimulate re-epithelialization from the wound bed. The following are a non-exhaustive list of grafts that polynucleotides or polypeptides, agonists or antagonists
30 of the invention, could be used to increase adherence to a wound bed: autografts, artificial skin, allografts, autodermic graft, autoepdermic grafts, avacular grafts, Blair-Brown grafts, bone graft, brephoplastic grafts, cutis graft, delayed graft, dermic graft,

epidermic graft, fascia graft, full thickness graft, heterologous graft, xenograft, homologous graft, hyperplastic graft, lamellar graft, mesh graft, mucosal graft, Ollier-Thiersch graft, omentum graft, patch graft, pedicle graft, penetrating graft, split skin graft, thick split graft. The polynucleotides or polypeptides, and/or agonists or
5 antagonists of the invention, can be used to promote skin strength and to improve the appearance of aged skin.

It is believed that the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, will also produce changes in hepatocyte proliferation, and epithelial cell proliferation in the lung, breast, pancreas, stomach, small intestine,
10 and large intestine. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could promote proliferation of epithelial cells such as sebocytes, hair follicles, hepatocytes, type II pneumocytes, mucin-producing goblet cells, and other epithelial cells and their progenitors contained within the skin, lung, liver, and gastrointestinal tract. The polynucleotides or polypeptides, and/or agonists
15 or antagonists of the invention, may promote proliferation of endothelial cells, keratinocytes, and basal keratinocytes.

The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could also be used to reduce the side effects of gut toxicity that result from radiation, chemotherapy treatments or viral infections. The polynucleotides or
20 polypeptides, and/or agonists or antagonists of the invention, may have a cytoprotective effect on the small intestine mucosa. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, may also stimulate healing of mucositis (mouth ulcers) that result from chemotherapy and viral infections.

25 The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could further be used in full regeneration of skin in full and partial thickness skin defects, including burns, (i.e., repopulation of hair follicles, sweat glands, and sebaceous glands), treatment of other skin defects such as psoriasis. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could
30 be used to treat epidermolysis bullosa, a defect in adherence of the epidermis to the underlying dermis which results in frequent, open and painful blisters by accelerating reepithelialization of these lesions. The polynucleotides or polypeptides, and/or

agonists or antagonists of the invention, could also be used to treat gastric and duodenal ulcers and help heal by scar formation of the mucosal lining and regeneration of glandular mucosa and duodenal mucosal lining more rapidly. Inflammatory bowel diseases, such as Crohn's disease and ulcerative colitis, are diseases which result in destruction of the mucosal surface of the small or large intestine, respectively. Thus, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to promote the resurfacing of the mucosal surface to aid more rapid healing and to prevent progression of inflammatory bowel disease. Treatment with the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, is expected to have a significant effect on the production of mucus throughout the gastrointestinal tract and could be used to protect the intestinal mucosa from injurious substances that are ingested or following surgery. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to treat diseases associated with the under expression of the polynucleotides of the invention.

Moreover, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to prevent and heal damage to the lungs due to various pathological states. A growth factor such as the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, which could stimulate proliferation and differentiation and promote the repair of alveoli and bronchiolar epithelium to prevent or treat acute or chronic lung damage. For example, emphysema, which results in the progressive loss of alveoli, and inhalation injuries, i.e., resulting from smoke inhalation and burns, that cause necrosis of the bronchiolar epithelium and alveoli could be effectively treated, prevented, and/or diagnosed using the polynucleotides or polypeptides, and/or agonists or antagonists of the invention. Also, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to stimulate the proliferation of and differentiation of type II pneumocytes, which may help treat or prevent disease such as hyaline membrane diseases, such as infant respiratory distress syndrome and bronchopulmonary dysplasia, in premature infants.

The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could stimulate the proliferation and differentiation of hepatocytes and,

thus, could be used to alleviate or treat liver diseases and pathologies such as fulminant liver failure caused by cirrhosis, liver damage caused by viral hepatitis and toxic substances (i.e., acetaminophen, carbon tetrachloride and other hepatotoxins known in the art).

5 In addition, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used treat or prevent the onset of diabetes mellitus. In patients with newly diagnosed Types I and II diabetes, where some islet cell function remains, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to maintain the islet function so as to alleviate, delay or
10 prevent permanent manifestation of the disease. Also, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used as an auxiliary in islet cell transplantation to improve or promote islet cell function.

15 Neurological Diseases

Nervous system diseases, disorders, and/or conditions, which can be treated, prevented, and/or diagnosed with the compositions of the invention (e.g., polypeptides, polynucleotides, and/or agonists or antagonists), include, but are not limited to, nervous system injuries, and diseases, disorders, and/or conditions which
20 result in either a disconnection of axons, a diminution or degeneration of neurons, or demyelination. Nervous system lesions which may be treated, prevented, and/or diagnosed in a patient (including human and non-human mammalian patients) according to the invention, include but are not limited to, the following lesions of either the central (including spinal cord, brain) or peripheral nervous systems: (1)
25 ischemic lesions, in which a lack of oxygen in a portion of the nervous system results in neuronal injury or death, including cerebral infarction or ischemia, or spinal cord infarction or ischemia; (2) traumatic lesions, including lesions caused by physical injury or associated with surgery, for example, lesions which sever a portion of the nervous system, or compression injuries; (3) malignant lesions, in which a portion of
30 the nervous system is destroyed or injured by malignant tissue which is either a nervous system associated malignancy or a malignancy derived from non-nervous system tissue; (4) infectious lesions, in which a portion of the nervous system is

destroyed or injured as a result of infection, for example, by an abscess or associated with infection by human immunodeficiency virus, herpes zoster, or herpes simplex virus or with Lyme disease, tuberculosis, syphilis; (5) degenerative lesions, in which a portion of the nervous system is destroyed or injured as a result of a degenerative process including but not limited to degeneration associated with Parkinson's disease, Alzheimer's disease, Huntington's chorea, or amyotrophic lateral sclerosis (ALS); (6) lesions associated with nutritional diseases, disorders, and/or conditions, in which a portion of the nervous system is destroyed or injured by a nutritional disorder or disorder of metabolism including but not limited to, vitamin B12 deficiency, folic acid deficiency, Wernicke disease, tobacco-alcohol amblyopia, Marchiafava-Bignami disease (primary degeneration of the corpus callosum), and alcoholic cerebellar degeneration; (7) neurological lesions associated with systemic diseases including, but not limited to, diabetes (diabetic neuropathy, Bell's palsy), systemic lupus erythematosus, carcinoma, or sarcoidosis; (8) lesions caused by toxic substances including alcohol, lead, or particular neurotoxins; and (9) demyelinated lesions in which a portion of the nervous system is destroyed or injured by a demyelinating disease including, but not limited to, multiple sclerosis, human immunodeficiency virus-associated myelopathy, transverse myelopathy or various etiologies, progressive multifocal leukoencephalopathy, and central pontine myelinolysis.

In a preferred embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to protect neural cells from the damaging effects of cerebral hypoxia. According to this embodiment, the compositions of the invention are used to treat, prevent, and/or diagnose neural cell injury associated with cerebral hypoxia. In one aspect of this embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to treat, prevent, and/or diagnose neural cell injury associated with cerebral ischemia. In another aspect of this embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to treat, prevent, and/or diagnose neural cell injury associated with cerebral infarction. In another aspect of this embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to treat, prevent, and/or diagnose or prevent neural cell injury associated with a stroke. In a further aspect of this embodiment, the polypeptides, polynucleotides, or agonists

or antagonists of the invention are used to treat, prevent, and/or diagnose neural cell injury associated with a heart attack.

The compositions of the invention which are useful for treating or preventing a nervous system disorder may be selected by testing for biological activity in promoting the survival or differentiation of neurons. For example, and not by way of limitation, compositions of the invention which elicit any of the following effects may be useful according to the invention: (1) increased survival time of neurons in culture; (2) increased sprouting of neurons in culture or *in vivo*; (3) increased production of a neuron-associated molecule in culture or *in vivo*, e.g., choline acetyltransferase or acetylcholinesterase with respect to motor neurons; or (4) decreased symptoms of neuron dysfunction *in vivo*. Such effects may be measured by any method known in the art. In preferred, non-limiting embodiments, increased survival of neurons may routinely be measured using a method set forth herein or otherwise known in the art, such as, for example, the method set forth in Arakawa et al. (J. Neurosci. 10:3507-3515 (1990)); increased sprouting of neurons may be detected by methods known in the art, such as, for example, the methods set forth in Pestronk et al. (Exp. Neurol. 70:65-82 (1980)) or Brown et al. (Ann. Rev. Neurosci. 4:17-42 (1981)); increased production of neuron-associated molecules may be measured by bioassay, enzymatic assay, antibody binding, Northern blot assay, etc., using techniques known in the art and depending on the molecule to be measured; and motor neuron dysfunction may be measured by assessing the physical manifestation of motor neuron disorder, e.g., weakness, motor neuron conduction velocity, or functional disability.

In specific embodiments, motor neuron diseases, disorders, and/or conditions that may be treated, prevented, and/or diagnosed according to the invention include, but are not limited to, diseases, disorders, and/or conditions such as infarction, infection, exposure to toxin, trauma, surgical damage, degenerative disease or malignancy that may affect motor neurons as well as other components of the nervous system, as well as diseases, disorders, and/or conditions that selectively affect neurons such as amyotrophic lateral sclerosis, and including, but not limited to, progressive spinal muscular atrophy, progressive bulbar palsy, primary lateral sclerosis, infantile and juvenile muscular atrophy, progressive bulbar paralysis of childhood (Fazio-

Londe syndrome), poliomyelitis and the post polio syndrome, and Hereditary
Motorsensory Neuropathy (Charcot-Marie-Tooth Disease).

Further, polypeptides or polynucleotides of the invention may play a role in
neuronal survival; synapse formation; conductance; neural differentiation, etc. Thus,
5 compositions of the invention (including polynucleotides, polypeptides, and agonists
or antagonists) may be used to diagnose and/or treat or prevent diseases or disorders
associated with these roles, including, but not limited to, learning and/or cognition
disorders. The compositions of the invention may also be useful in the treatment or
prevention of neurodegenerative disease states and/or behavioural disorders. Such
10 neurodegenerative disease states and/or behavioral disorders include, but are not
limited to, Alzheimers Disease, Parkinsons Disease, Huntingtons Disease, Tourette
Syndrome, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder,
panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors,
including disorders in feeding, sleep patterns, balance, and perception. In addition,
15 compositions of the invention may also play a role in the treatment, prevention and/or
detection of developmental disorders associated with the developing embryo, or
sexually-linked disorders.

Additionally, polypeptides, polynucleotides and/or agonists or antagonists of
the invention, may be useful in protecting neural cells from diseases, damage,
20 disorders, or injury, associated with cerebrovascular disorders including, but not
limited to, carotid artery diseases (e.g., carotid artery thrombosis, carotid stenosis, or
Moyamoya Disease), cerebral amyloid angiopathy, cerebral aneurysm, cerebral
anoxia, cerebral arteriosclerosis, cerebral arteriovenous malformations, cerebral artery
diseases, cerebral embolism and thrombosis (e.g., carotid artery thrombosis, sinus
25 thrombosis, or Wallenberg's Syndrome), cerebral hemorrhage (e.g., epidural or
subdural hematoma, or subarachnoid hemorrhage), cerebral infarction, cerebral
ischemia (e.g., transient cerebral ischemia, Subclavian Steal Syndrome, or
vertebrobasilar insufficiency), vascular dementia (e.g., multi-infarct), leukomalacia,
periventricular, and vascular headache (e.g., cluster headache or migraines).

30 In accordance with yet a further aspect of the present invention, there is
provided a process for utilizing polynucleotides or polypeptides, as well as agonists or
antagonists of the present invention, for therapeutic purposes, for example, to

stimulate neurological cell proliferation and/or differentiation. Therefore, polynucleotides, polypeptides, agonists and/or antagonists of the invention may be used to treat and/or detect neurologic diseases. Moreover, polynucleotides or polypeptides, or agonists or antagonists of the invention, can be used as a marker or
5 detector of a particular nervous system disease or disorder.

Examples of neurologic diseases which can be treated or detected with polynucleotides, polypeptides, agonists, and/or antagonists of the present invention include brain diseases, such as metabolic brain diseases which includes phenylketonuria such as maternal phenylketonuria, pyruvate carboxylase deficiency,
10 pyruvate dehydrogenase complex deficiency, Wernicke's Encephalopathy, brain edema, brain neoplasms such as cerebellar neoplasms which include infratentorial neoplasms, cerebral ventricle neoplasms such as choroid plexus neoplasms, hypothalamic neoplasms, supratentorial neoplasms, canavan disease, cerebellar diseases such as cerebellar ataxia which include spinocerebellar degeneration such as
15 ataxia telangiectasia, cerebellar dyssynergia, Friederich's Ataxia, Machado-Joseph Disease, olivopontocerebellar atrophy, cerebellar neoplasms such as infratentorial neoplasms, diffuse cerebral sclerosis such as encephalitis periaxialis, globoid cell leukodystrophy, metachromatic leukodystrophy and subacute sclerosing panencephalitis.

20 Additional neurologic diseases which can be treated or detected with polynucleotides, polypeptides, agonists, and/or antagonists of the present invention include cerebrovascular disorders (such as carotid artery diseases which include carotid artery thrombosis, carotid stenosis and Moyamoya Disease), cerebral amyloid angiopathy, cerebral aneurysm, cerebral anoxia, cerebral arteriosclerosis, cerebral
25 arteriovenous malformations, cerebral artery diseases, cerebral embolism and thrombosis such as carotid artery thrombosis, sinus thrombosis and Wallenberg's Syndrome, cerebral hemorrhage such as epidural hematoma, subdural hematoma and subarachnoid hemorrhage, cerebral infarction, cerebral ischemia such as transient cerebral ischemia, Subclavian Steal Syndrome and vertebrobasilar insufficiency,
30 vascular dementia such as multi-infarct dementia, periventricular leukomalacia, vascular headache such as cluster headache and migraine.

Additional neurologic diseases which can be treated or detected with polynucleotides, polypeptides, agonists, and/or antagonists of the present invention include dementia such as AIDS Dementia Complex, presenile dementia such as Alzheimer's Disease and Creutzfeldt-Jakob Syndrome, senile dementia such as
5 Alzheimer's Disease and progressive supranuclear palsy, vascular dementia such as multi-infarct dementia, encephalitis which include encephalitis periaxialis, viral encephalitis such as epidemic encephalitis, Japanese Encephalitis, St. Louis Encephalitis, tick-borne encephalitis and West Nile Fever, acute disseminated encephalomyelitis, meningoencephalitis such as uveomeningoencephalitic syndrome,
10 Postencephalitic Parkinson Disease and subacute sclerosing panencephalitis, encephalomalacia such as periventricular leukomalacia, epilepsy such as generalized epilepsy which includes infantile spasms, absence epilepsy, myoclonic epilepsy which includes MERRF Syndrome, tonic-clonic epilepsy, partial epilepsy such as complex partial epilepsy, frontal lobe epilepsy and temporal lobe epilepsy, post-traumatic
15 epilepsy, status epilepticus such as Epilepsia Partialis Continua, and Hallervorden-Spatz Syndrome.

Additional neurologic diseases which can be treated or detected with polynucleotides, polypeptides, agonists, and/or antagonists of the present invention include hydrocephalus such as Dandy-Walker Syndrome and normal pressure
20 hydrocephalus, hypothalamic diseases such as hypothalamic neoplasms, cerebral malaria, narcolepsy which includes cataplexy, bulbar poliomyelitis, cerebri pseudotumor, Rett Syndrome, Reye's Syndrome, thalamic diseases, cerebral toxoplasmosis, intracranial tuberculoma and Zellweger Syndrome, central nervous system infections such as AIDS Dementia Complex, Brain Abscess, subdural
25 empyema, encephalomyelitis such as Equine Encephalomyelitis, Venezuelan Equine Encephalomyelitis, Necrotizing Hemorrhagic Encephalomyelitis, Visna, and cerebral malaria.

Additional neurologic diseases which can be treated or detected with polynucleotides, polypeptides, agonists, and/or antagonists of the present invention
30 include meningitis such as arachnoiditis, aseptic meningitis such as viral meningitis which includes lymphocytic choriomeningitis, Bacterial meningitis which includes Haemophilus Meningitis, Listeria Meningitis, Meningococcal Meningitis such as

Waterhouse-Friderichsen Syndrome, Pneumococcal Meningitis and meningeal tuberculosis, fungal meningitis such as Cryptococcal Meningitis, subdural effusion, meningoencephalitis such as uvemeningoencephalitic syndrome, myelitis such as transverse myelitis, neurosyphilis such as tabes dorsalis, poliomyelitis which includes
5 bulbar poliomyelitis and postpoliomyelitis syndrome, prion diseases (such as Creutzfeldt-Jakob Syndrome, Bovine Spongiform Encephalopathy, Gerstmann-Straussler Syndrome, Kuru, Scrapie), and cerebral toxoplasmosis.

Additional neurologic diseases which can be treated or detected with polynucleotides, polypeptides, agonists, and/or antagonists of the present invention
10 include central nervous system neoplasms such as brain neoplasms that include cerebellar neoplasms such as infratentorial neoplasms, cerebral ventricle neoplasms such as choroid plexus neoplasms, hypothalamic neoplasms and supratentorial neoplasms, meningeal neoplasms, spinal cord neoplasms which include epidural neoplasms, demyelinating diseases such as Canavan Diseases, diffuse cerebral
15 sceloris which includes adrenoleukodystrophy, encephalitis periaxialis, globoid cell leukodystrophy, diffuse cerebral sclerosis such as metachromatic leukodystrophy, allergic encephalomyelitis, necrotizing hemorrhagic encephalomyelitis, progressive multifocal leukoencephalopathy, multiple sclerosis, central pontine myelinolysis, transverse myelitis, neuromyelitis optica, Scrapie, Swayback, Chronic Fatigue
20 Syndrome, Visna, High Pressure Nervous Syndrome, Meningism, spinal cord diseases such as amyotonia congenita, amyotrophic lateral sclerosis, spinal muscular atrophy such as Werdnig-Hoffmann Disease, spinal cord compression, spinal cord neoplasms such as epidural neoplasms, syringomyelia, Tabes Dorsalis, Stiff-Man Syndrome, mental retardation such as Angelman Syndrome, Cri-du-Chat Syndrome, De Lange's
25 Syndrome, Down Syndrome, Gangliosidoses such as gangliosidoses G(M1), Sandhoff Disease, Tay-Sachs Disease, Hartnup Disease, homocystinuria, Laurence-Moon-Biedl Syndrome, Lesch-Nyhan Syndrome, Maple Syrup Urine Disease, mucopolipidosis such as fucosidosis, neuronal ceroid-lipofuscinosis, oculocerebrorenal syndrome, phenylketonuria such as maternal phenylketonuria, Prader-Willi Syndrome, Rett
30 Syndrome, Rubinstein-Taybi Syndrome, Tuberous Sclerosis, WAGR Syndrome, nervous system abnormalities such as holoprosencephaly, neural tube defects such as anencephaly which includes hydrangencephaly, Arnold-Chairi Deformity,

encephalocele, meningocele, meningomyelocele, spinal dysraphism such as spina bifida cystica and spina bifida occulta.

Additional neurologic diseases which can be treated or detected with polynucleotides, polypeptides, agonists, and/or antagonists of the present invention

5 include hereditary motor and sensory neuropathies which include Charcot-Marie Disease, Hereditary optic atrophy, Refsum's Disease, hereditary spastic paraplegia, Werdnig-Hoffmann Disease, Hereditary Sensory and Autonomic Neuropathies such as Congenital Analgesia and Familial Dysautonomia, Neurologic manifestations (such as agnosia that include Gerstmann's Syndrome, Amnesia such as retrograde amnesia,

10 apraxia, neurogenic bladder, cataplexy, communicative disorders such as hearing disorders that includes deafness, partial hearing loss, loudness recruitment and tinnitus, language disorders such as aphasia which include agraphia, anomia, broca aphasia, and Wernicke Aphasia, Dyslexia such as Acquired Dyslexia, language development disorders, speech disorders such as aphasia which includes anomia,

15 broca aphasia and Wernicke Aphasia, articulation disorders, communicative disorders such as speech disorders which include dysarthria, echolalia, mutism and stuttering, voice disorders such as aphonia and hoarseness, decerebrate state, delirium, fasciculation, hallucinations, meningism, movement disorders such as angelman syndrome, ataxia, athetosis, chorea, dystonia, hypokinesia, muscle hypotonia,

20 myoclonus, tic, torticollis and tremor, muscle hypertonia such as muscle rigidity such as stiff-man syndrome, muscle spasticity, paralysis such as facial paralysis which includes Herpes Zoster Oticus, Gastroparesis, Hemiplegia, ophthalmoplegia such as diplopia, Duane's Syndrome, Horner's Syndrome, Chronic progressive external ophthalmoplegia such as Kearns Syndrome, Bulbar Paralysis, Tropical Spastic

25 Paraparesis, Paraplegia such as Brown-Sequard Syndrome, quadriplegia, respiratory paralysis and vocal cord paralysis, paresis, phantom limb, taste disorders such as ageusia and dysgeusia, vision disorders such as amblyopia, blindness, color vision defects, diplopia, hemianopsia, scotoma and subnormal vision, sleep disorders such as hypersomnia which includes Kleine-Levin Syndrome, insomnia, and somnambulism,

30 spasm such as trismus, unconsciousness such as coma, persistent vegetative state and syncope and vertigo, neuromuscular diseases such as amyotonia congenita, amyotrophic lateral sclerosis, Lambert-Eaton Myasthenic Syndrome, motor neuron

disease, muscular atrophy such as spinal muscular atrophy, Charcot-Marie Disease and Werdnig-Hoffmann Disease, Postpoliomyelitis Syndrome, Muscular Dystrophy, Myasthenia Gravis, Myotonia Atrophica, Myotonia Confenita, Nemaline Myopathy, Familial Periodic Paralysis, Multiplex Paramyoclonus, Tropical Spastic Paraparesis and Stiff-Man Syndrome, peripheral nervous system diseases such as acrodynia, amyloid neuropathies, autonomic nervous system diseases such as Adie's Syndrome, Barre-Lieou Syndrome, Familial Dysautonomia, Horner's Syndrome, Reflex Sympathetic Dystrophy and Shy-Drager Syndrome, Cranial Nerve Diseases such as Acoustic Nerve Diseases such as Acoustic Neuroma which includes Neurofibromatosis 2, Facial Nerve Diseases such as Facial Neuralgia, Melkersson-Rosenthal Syndrome, ocular motility disorders which includes amblyopia, nystagmus, oculomotor nerve paralysis, ophthalmoplegia such as Duane's Syndrome, Horner's Syndrome, Chronic Progressive External Ophthalmoplegia which includes Kearns Syndrome, Strabismus such as Esotropia and Exotropia, Oculomotor Nerve Paralysis, Optic Nerve Diseases such as Optic Atrophy which includes Hereditary Optic Atrophy, Optic Disk Drusen, Optic Neuritis such as Neuromyelitis Optica, Papilledema, Trigeminal Neuralgia, Vocal Cord Paralysis, Demyelinating Diseases such as Neuromyelitis Optica and Swayback, and Diabetic neuropathies such as diabetic foot.

Additional neurologic diseases which can be treated or detected with polynucleotides, polypeptides, agonists, and/or antagonists of the present invention include nerve compression syndromes such as carpal tunnel syndrome, tarsal tunnel syndrome, thoracic outlet syndrome such as cervical rib syndrome, ulnar nerve compression syndrome, neuralgia such as causalgia, cervico-brachial neuralgia, facial neuralgia and trigeminal neuralgia, neuritis such as experimental allergic neuritis, optic neuritis, polyneuritis, polyradiculoneuritis and radiculities such as polyradiculitis, hereditary motor and sensory neuropathies such as Charcot-Marie Disease, Hereditary Optic Atrophy, Refsum's Disease, Hereditary Spastic Paraplegia and Werdnig-Hoffmann Disease, Hereditary Sensory and Autonomic Neuropathies which include Congenital Analgesia and Familial Dysautonomia, POEMS Syndrome, Sciatica, Gustatory Sweating and Tetany).

Infectious Disease

A polypeptide or polynucleotide and/or agonist or antagonist of the present invention can be used to treat, prevent, and/or diagnose infectious agents. For example, by increasing the immune response, particularly increasing the proliferation and differentiation of B and/or T cells, infectious diseases may be treated, prevented, and/or diagnosed. The immune response may be increased by either enhancing an existing immune response, or by initiating a new immune response. Alternatively, polypeptide or polynucleotide and/or agonist or antagonist of the present invention may also directly inhibit the infectious agent, without necessarily eliciting an immune response.

Viruses are one example of an infectious agent that can cause disease or symptoms that can be treated, prevented, and/or diagnosed by a polynucleotide or polypeptide and/or agonist or antagonist of the present invention. Examples of viruses, include, but are not limited to Examples of viruses, include, but are not limited to the following DNA and RNA viruses and viral families: Arbovirus, Adenoviridae, Arenaviridae, Arterivirus, Birnaviridae, Bunyaviridae, Caliciviridae, Circoviridae, Coronaviridae, Dengue, EBV, HIV, Flaviviridae, Hepadnaviridae (Hepatitis), Herpesviridae (such as, Cytomegalovirus, Herpes Simplex, Herpes Zoster), Mononegavirus (e.g., Paramyxoviridae, Morbillivirus, Rhabdoviridae), Orthomyxoviridae (e.g., Influenza A, Influenza B, and parainfluenza), Papilloma virus, Papovaviridae, Parvoviridae, Picornaviridae, Poxviridae (such as Smallpox or Vaccinia), Reoviridae (e.g., Rotavirus), Retroviridae (HTLV-I, HTLV-II, Lentivirus), and Togaviridae (e.g., Rubivirus). Viruses falling within these families can cause a variety of diseases or symptoms, including, but not limited to: arthritis, bronchiolitis, respiratory syncytial virus, encephalitis, eye infections (e.g., conjunctivitis, keratitis), chronic fatigue syndrome, hepatitis (A, B, C, E, Chronic Active, Delta), Japanese B encephalitis, Junin, Chikungunya, Rift Valley fever, yellow fever, meningitis, opportunistic infections (e.g., AIDS), pneumonia, Burkitt's Lymphoma, chickenpox, hemorrhagic fever, Measles, Mumps, Parainfluenza, Rabies, the common cold, Polio, leukemia, Rubella, sexually transmitted diseases, skin diseases (e.g., Kaposi's, warts), and viremia. polynucleotides or polypeptides, or agonists or antagonists of the invention, can be used to treat, prevent, and/or diagnose any of these symptoms or

diseases. In specific embodiments, polynucleotides, polypeptides, or agonists or antagonists of the invention are used to treat, prevent, and/or diagnose: meningitis, Dengue, EBV, and/or hepatitis (e.g., hepatitis B). In an additional specific embodiment polynucleotides, polypeptides, or agonists or antagonists of the invention are used to treat patients nonresponsive to one or more other commercially available hepatitis vaccines. In a further specific embodiment polynucleotides, polypeptides, or agonists or antagonists of the invention are used to treat, prevent, and/or diagnose AIDS.

Similarly, bacterial or fungal agents that can cause disease or symptoms and that can be treated, prevented, and/or diagnosed by a polynucleotide or polypeptide and/or agonist or antagonist of the present invention include, but not limited to, include, but not limited to, the following Gram-Negative and Gram-positive bacteria and bacterial families and fungi: Actinomycetales (e.g., *Corynebacterium*, *Mycobacterium*, *Nocardia*), *Cryptococcus neoformans*, Aspergillosis, Bacillaceae (e.g., Anthrax, *Clostridium*), Bacteroidaceae, Blastomycosis, *Bordetella*, *Borrelia* (e.g., *Borrelia burgdorferi*), Brucellosis, Candidiasis, *Campylobacter*, Coccidioidomycosis, Cryptococcosis, Dermatocycoses, *E. coli* (e.g., Enterotoxigenic *E. coli* and Enterohemorrhagic *E. coli*), Enterobacteriaceae (*Klebsiella*, *Salmonella* (e.g., *Salmonella typhi*, and *Salmonella paratyphi*), *Serratia*, *Yersinia*), *Erysipelothrix*, *Helicobacter*, Legionellosis, Leptospirosis, *Listeria*, Mycoplasmatales, *Mycobacterium leprae*, *Vibrio cholerae*, Neisseriaceae (e.g., *Acinetobacter*, Gonorrhea, Meningococcal), *Meisseria meningitidis*, Pasteurellaceae Infections (e.g., *Actinobacillus*, *Haemophilus* (e.g., *Haemophilus influenza* type B), *Pasteurella*), *Pseudomonas*, Rickettsiaceae, Chlamydiaceae, Syphilis, *Shigella* spp., Staphylococcal, Meningiococcal, Pneumococcal and Streptococcal (e.g., *Streptococcus pneumoniae* and Group B Streptococcus). These bacterial or fungal families can cause the following diseases or symptoms, including, but not limited to: bacteremia, endocarditis, eye infections (conjunctivitis, tuberculosis, uveitis), gingivitis, opportunistic infections (e.g., AIDS related infections), paronychia, prosthesis-related infections, Reiter's Disease, respiratory tract infections, such as Whooping Cough or Empyema, sepsis, Lyme Disease, Cat-Scratch Disease, Dysentery, Paratyphoid Fever, food poisoning, Typhoid, pneumonia, Gonorrhea,

meningitis (e.g., meningitis types A and B), Chlamydia, Syphilis, Diphtheria, Leprosy, Paratuberculosis, Tuberculosis, Lupus, Botulism, gangrene, tetanus, impetigo, Rheumatic Fever, Scarlet Fever, sexually transmitted diseases, skin diseases (e.g., cellulitis, dermatocycoses), toxemia, urinary tract infections, wound infections.

- 5 Polynucleotides or polypeptides, agonists or antagonists of the invention, can be used to treat, prevent, and/or diagnose any of these symptoms or diseases. In specific embodiments, polynucleotides, polypeptides, agonists or antagonists of the invention are used to treat, prevent, and/or diagnose: tetanus, Diphtheria, botulism, and/or meningitis type B.

- 10 Moreover, parasitic agents causing disease or symptoms that can be treated, prevented, and/or diagnosed by a polynucleotide or polypeptide and/or agonist or antagonist of the present invention include, but not limited to, the following families or class: Amebiasis, Babesiosis, Coccidiosis, Cryptosporidiosis, Dientamoebiasis, Dourine, Ectoparasitic, Giardiasis, Helminthiasis, Leishmaniasis, Theileriasis,
- 15 Toxoplasmosis, Trypanosomiasis, and Trichomonas and Sporozoans (e.g., Plasmodium virax, Plasmodium falciparum, Plasmodium malariae and Plasmodium ovale). These parasites can cause a variety of diseases or symptoms, including, but not limited to: Scabies, Trombiculiasis, eye infections, intestinal disease (e.g., dysentery, giardiasis), liver disease, lung disease, opportunistic infections (e.g., AIDS
- 20 related), malaria, pregnancy complications, and toxoplasmosis. polynucleotides or polypeptides, or agonists or antagonists of the invention, can be used to treat, prevent, and/or diagnose any of these symptoms or diseases. In specific embodiments, polynucleotides, polypeptides, or agonists or antagonists of the invention are used to treat, prevent, and/or diagnose malaria.

- 25 Preferably, treatment or prevention using a polypeptide or polynucleotide and/or agonist or antagonist of the present invention could either be by administering an effective amount of a polypeptide to the patient, or by removing cells from the patient, supplying the cells with a polynucleotide of the present invention, and returning the engineered cells to the patient (ex vivo therapy). Moreover, the
- 30 polypeptide or polynucleotide of the present invention can be used as an antigen in a vaccine to raise an immune response against infectious disease.

Regeneration

A polynucleotide or polypeptide and/or agonist or antagonist of the present invention can be used to differentiate, proliferate, and attract cells, leading to the regeneration of tissues. (See, Science 276:59-87 (1997).) The regeneration of tissues
5 could be used to repair, replace, or protect tissue damaged by congenital defects, trauma (wounds, burns, incisions, or ulcers), age, disease (e.g. osteoporosis, osteoarthritis, periodontal disease, liver failure), surgery, including cosmetic plastic surgery, fibrosis, reperfusion injury, or systemic cytokine damage.

Tissues that could be regenerated using the present invention include organs
10 (e.g., pancreas, liver, intestine, kidney, skin, endothelium), muscle (smooth, skeletal or cardiac), vasculature (including vascular and lymphatics), nervous, hematopoietic, and skeletal (bone, cartilage, tendon, and ligament) tissue. Preferably, regeneration occurs without or decreased scarring. Regeneration also may include angiogenesis.

Moreover, a polynucleotide or polypeptide and/or agonist or antagonist of the
15 present invention may increase regeneration of tissues difficult to heal. For example, increased tendon/ligament regeneration would quicken recovery time after damage. A polynucleotide or polypeptide and/or agonist or antagonist of the present invention could also be used prophylactically in an effort to avoid damage. Specific diseases that could be treated, prevented, and/or diagnosed include of tendinitis, carpal tunnel
20 syndrome, and other tendon or ligament defects. A further example of tissue regeneration of non-healing wounds includes pressure ulcers, ulcers associated with vascular insufficiency, surgical, and traumatic wounds.

Similarly, nerve and brain tissue could also be regenerated by using a
polynucleotide or polypeptide and/or agonist or antagonist of the present invention to
25 proliferate and differentiate nerve cells. Diseases that could be treated, prevented, and/or diagnosed using this method include central and peripheral nervous system diseases, neuropathies, or mechanical and traumatic diseases, disorders, and/or conditions (e.g., spinal cord disorders, head trauma, cerebrovascular disease, and stroke). Specifically, diseases associated with peripheral nerve injuries, peripheral
30 neuropathy (e.g., resulting from chemotherapy or other medical therapies), localized neuropathies, and central nervous system diseases (e.g., Alzheimer's disease, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-

Drager syndrome), could all be treated, prevented, and/or diagnosed using the polynucleotide or polypeptide and/or agonist or antagonist of the present invention.

Chemotaxis

5 A polynucleotide or polypeptide and/or agonist or antagonist of the present invention may have chemotaxis activity. A chemotactic molecule attracts or mobilizes cells (e.g., monocytes, fibroblasts, neutrophils, T-cells, mast cells, eosinophils, epithelial and/or endothelial cells) to a particular site in the body, such as inflammation, infection, or site of hyperproliferation. The mobilized cells can then
10 fight off and/or heal the particular trauma or abnormality.

 A polynucleotide or polypeptide and/or agonist or antagonist of the present invention may increase chemotactic activity of particular cells. These chemotactic molecules can then be used to treat, prevent, and/or diagnose inflammation, infection, hyperproliferative diseases, disorders, and/or conditions, or any immune system
15 disorder by increasing the number of cells targeted to a particular location in the body. For example, chemotactic molecules can be used to treat, prevent, and/or diagnose wounds and other trauma to tissues by attracting immune cells to the injured location. Chemotactic molecules of the present invention can also attract fibroblasts, which can be used to treat, prevent, and/or diagnose wounds.

20 It is also contemplated that a polynucleotide or polypeptide and/or agonist or antagonist of the present invention may inhibit chemotactic activity. These molecules could also be used to treat, prevent, and/or diagnose diseases, disorders, and/or conditions. Thus, a polynucleotide or polypeptide and/or agonist or antagonist of the present invention could be used as an inhibitor of chemotaxis.

25

Binding Activity

 A polypeptide of the present invention may be used to screen for molecules that bind to the polypeptide or for molecules to which the polypeptide binds. The binding of the polypeptide and the molecule may activate (agonist), increase, inhibit
30 (antagonist), or decrease activity of the polypeptide or the molecule bound. Examples of such molecules include antibodies, oligonucleotides, proteins (e.g., receptors), or small molecules.

Preferably, the molecule is closely related to the natural ligand of the polypeptide, e.g., a fragment of the ligand, or a natural substrate, a ligand, a structural or functional mimetic. (See, Coligan et al., Current Protocols in Immunology 1(2):Chapter 5 (1991).) Similarly, the molecule can be closely related to the natural
5 receptor to which the polypeptide binds, or at least, a fragment of the receptor capable of being bound by the polypeptide (e.g., active site). In either case, the molecule can be rationally designed using known techniques.

Preferably, the screening for these molecules involves producing appropriate cells which express the polypeptide, either as a secreted protein or on the cell
10 membrane. Preferred cells include cells from mammals, yeast, *Drosophila*, or *E. coli*. Cells expressing the polypeptide (or cell membrane containing the expressed polypeptide) are then preferably contacted with a test compound potentially containing the molecule to observe binding, stimulation, or inhibition of activity of either the polypeptide or the molecule.

15 The assay may simply test binding of a candidate compound to the polypeptide, wherein binding is detected by a label, or in an assay involving competition with a labeled competitor. Further, the assay may test whether the candidate compound results in a signal generated by binding to the polypeptide.

Alternatively, the assay can be carried out using cell-free preparations,
20 polypeptide/molecule affixed to a solid support, chemical libraries, or natural product mixtures. The assay may also simply comprise the steps of mixing a candidate compound with a solution containing a polypeptide, measuring polypeptide/molecule activity or binding, and comparing the polypeptide/molecule activity or binding to a standard.

25 Preferably, an ELISA assay can measure polypeptide level or activity in a sample (e.g., biological sample) using a monoclonal or polyclonal antibody. The antibody can measure polypeptide level or activity by either binding, directly or indirectly, to the polypeptide or by competing with the polypeptide for a substrate.

Additionally, the receptor to which a polypeptide of the invention binds can be
30 identified by numerous methods known to those of skill in the art, for example, ligand panning and FACS sorting (Coligan, et al., Current Protocols in Immun., 1(2), Chapter 5, (1991)). For example, expression cloning is employed wherein

polyadenylated RNA is prepared from a cell responsive to the polypeptides, for example, NIH3T3 cells which are known to contain multiple receptors for the FGF family proteins, and SC-3 cells, and a cDNA library created from this RNA is divided into pools and used to transfect COS cells or other cells that are not responsive to the polypeptides. Transfected cells which are grown on glass slides are exposed to the polypeptide of the present invention, after they have been labelled. The polypeptides can be labeled by a variety of means including iodination or inclusion of a recognition site for a site-specific protein kinase.

Following fixation and incubation, the slides are subjected to autoradiographic analysis. Positive pools are identified and sub-pools are prepared and retransfected using an iterative sub-pooling and re-screening process, eventually yielding a single clones that encodes the putative receptor.

As an alternative approach for receptor identification, the labeled polypeptides can be photoaffinity linked with cell membrane or extract preparations that express the receptor molecule. Cross-linked material is resolved by PAGE analysis and exposed to X-ray film. The labeled complex containing the receptors of the polypeptides can be excised, resolved into peptide fragments, and subjected to protein microsequencing. The amino acid sequence obtained from microsequencing would be used to design a set of degenerate oligonucleotide probes to screen a cDNA library to identify the genes encoding the putative receptors.

Moreover, the techniques of gene-shuffling, motif-shuffling, exon-shuffling, and/or codon-shuffling (collectively referred to as "DNA shuffling") may be employed to modulate the activities of polypeptides of the invention thereby effectively generating agonists and antagonists of polypeptides of the invention. See generally, U.S. Patent Nos. 5,605,793, 5,811,238, 5,830,721, 5,834,252, and 5,837,458, and Patten, P. A., et al., Curr. Opinion Biotechnol. 8:724-33 (1997); Harayama, S. Trends Biotechnol. 16(2):76-82 (1998); Hansson, L. O., et al., J. Mol. Biol. 287:265-76 (1999); and Lorenzo, M. M. and Blasco, R. Biotechniques 24(2):308-13 (1998) (each of these patents and publications are hereby incorporated by reference). In one embodiment, alteration of polynucleotides and corresponding polypeptides of the invention may be achieved by DNA shuffling. DNA shuffling involves the assembly of two or more DNA segments into a desired polynucleotide

sequence of the invention molecule by homologous, or site-specific, recombination. In another embodiment, polynucleotides and corresponding polypeptides of the invention may be altered by being subjected to random mutagenesis by error-prone PCR, random nucleotide insertion or other methods prior to recombination. In
5 another embodiment, one or more components, motifs, sections, parts, domains, fragments, etc., of the polypeptides of the invention may be recombined with one or more components, motifs, sections, parts, domains, fragments, etc. of one or more heterologous molecules. In preferred embodiments, the heterologous molecules are family members. In further preferred embodiments, the heterologous molecule is a
10 growth factor such as, for example, platelet-derived growth factor (PDGF), insulin-like growth factor (IGF-I), transforming growth factor (TGF)-alpha, epidermal growth factor (EGF), fibroblast growth factor (FGF), TGF-beta, bone morphogenetic protein (BMP)-2, BMP-4, BMP-5, BMP-6, BMP-7, activins A and B, decapentaplegic(dpp), 60A, OP-2, dorsalin, growth differentiation factors (GDFs),
15 nodal, MIS, inhibin-alpha, TGF-beta1, TGF-beta2, TGF-beta3, TGF-beta5, and glial-derived neurotrophic factor (GDNF).

Other preferred fragments are biologically active fragments of the polypeptides of the invention. Biologically active fragments are those exhibiting activity similar, but not necessarily identical, to an activity of the polypeptide. The
20 biological activity of the fragments may include an improved desired activity, or a decreased undesirable activity.

Additionally, this invention provides a method of screening compounds to identify those which modulate the action of the polypeptide of the present invention. An example of such an assay comprises combining a mammalian fibroblast cell, a the
25 polypeptide of the present invention, the compound to be screened and 3[H] thymidine under cell culture conditions where the fibroblast cell would normally proliferate. A control assay may be performed in the absence of the compound to be screened and compared to the amount of fibroblast proliferation in the presence of the compound to determine if the compound stimulates proliferation by determining the
30 uptake of 3[H] thymidine in each case. The amount of fibroblast cell proliferation is measured by liquid scintillation chromatography which measures the incorporation of

3[H] thymidine. Both agonist and antagonist compounds may be identified by this procedure.

In another method, a mammalian cell or membrane preparation expressing a receptor for a polypeptide of the present invention is incubated with a labeled polypeptide of the present invention in the presence of the compound. The ability of the compound to enhance or block this interaction could then be measured. Alternatively, the response of a known second messenger system following interaction of a compound to be screened and the receptor is measured and the ability of the compound to bind to the receptor and elicit a second messenger response is measured to determine if the compound is a potential agonist or antagonist. Such second messenger systems include but are not limited to, cAMP guanylate cyclase, ion channels or phosphoinositide hydrolysis.

All of these above assays can be used as diagnostic or prognostic markers. The molecules discovered using these assays can be used to treat, prevent, and/or diagnose disease or to bring about a particular result in a patient (e.g., blood vessel growth) by activating or inhibiting the polypeptide/molecule. Moreover, the assays can discover agents which may inhibit or enhance the production of the polypeptides of the invention from suitably manipulated cells or tissues. Therefore, the invention includes a method of identifying compounds which bind to the polypeptides of the invention comprising the steps of: (a) incubating a candidate binding compound with the polypeptide; and (b) determining if binding has occurred. Moreover, the invention includes a method of identifying agonists/antagonists comprising the steps of: (a) incubating a candidate compound with the polypeptide, (b) assaying a biological activity, and (b) determining if a biological activity of the polypeptide has been altered.

Also, one could identify molecules bind a polypeptide of the invention experimentally by using the beta-pleated sheet regions contained in the polypeptide sequence of the protein. Accordingly, specific embodiments of the invention are directed to polynucleotides encoding polypeptides which comprise, or alternatively consist of, the amino acid sequence of each beta pleated sheet regions in a disclosed polypeptide sequence. Additional embodiments of the invention are directed to polynucleotides encoding polypeptides which comprise, or alternatively consist of,

any combination or all of contained in the polypeptide sequences of the invention. Additional preferred embodiments of the invention are directed to polypeptides which comprise, or alternatively consist of, the amino acid sequence of each of the beta pleated sheet regions in one of the polypeptide sequences of the invention. Additional
5 embodiments of the invention are directed to polypeptides which comprise, or alternatively consist of, any combination or all of the beta pleated sheet regions in one of the polypeptide sequences of the invention.

Targeted Delivery

10 In another embodiment, the invention provides a method of delivering compositions to targeted cells expressing a receptor for a polypeptide of the invention, or cells expressing a cell bound form of a polypeptide of the invention.

As discussed herein, polypeptides or antibodies of the invention may be associated with heterologous polypeptides, heterologous nucleic acids, toxins, or
15 prodrugs via hydrophobic, hydrophilic, ionic and/or covalent interactions. In one embodiment, the invention provides a method for the specific delivery of compositions of the invention to cells by administering polypeptides of the invention (including antibodies) that are associated with heterologous polypeptides or nucleic acids. In one example, the invention provides a method for delivering a therapeutic
20 protein into the targeted cell. In another example, the invention provides a method for delivering a single stranded nucleic acid (e.g., antisense or ribozymes) or double stranded nucleic acid (e.g., DNA that can integrate into the cell's genome or replicate episomally and that can be transcribed) into the targeted cell.

In another embodiment, the invention provides a method for the specific
25 destruction of cells (e.g., the destruction of tumor cells) by administering polypeptides of the invention (e.g., polypeptides of the invention or antibodies of the invention) in association with toxins or cytotoxic prodrugs.

By "toxin" is meant compounds that bind and activate endogenous cytotoxic effector systems, radioisotopes, holotoxins, modified toxins, catalytic subunits of
30 toxins, or any molecules or enzymes not normally present in or on the surface of a cell that under defined conditions cause the cell's death. Toxins that may be used according to the methods of the invention include, but are not limited to, radioisotopes

known in the art, compounds such as, for example, antibodies (or complement fixing containing portions thereof) that bind an inherent or induced endogenous cytotoxic effector system, thymidine kinase, endonuclease, RNase, alpha toxin, ricin, abrin, *Pseudomonas* exotoxin A, diphtheria toxin, saporin, momordin, gelonin, pokeweed
5 antiviral protein, alpha-sarcin and cholera toxin. By "cytotoxic prodrug" is meant a non-toxic compound that is converted by an enzyme, normally present in the cell, into a cytotoxic compound. Cytotoxic prodrugs that may be used according to the methods of the invention include, but are not limited to, glutamyl derivatives of benzoic acid mustard alkylating agent, phosphate derivatives of etoposide or
10 mitomycin C, cytosine arabinoside, daunorubisin, and phenoxyacetamide derivatives of doxorubicin.

Drug Screening

Further contemplated is the use of the polypeptides of the present invention, or
15 the polynucleotides encoding these polypeptides, to screen for molecules which modify the activities of the polypeptides of the present invention. Such a method would include contacting the polypeptide of the present invention with a selected compound(s) suspected of having antagonist or agonist activity, and assaying the activity of these polypeptides following binding.

20 This invention is particularly useful for screening therapeutic compounds by using the polypeptides of the present invention, or binding fragments thereof, in any of a variety of drug screening techniques. The polypeptide or fragment employed in such a test may be affixed to a solid support, expressed on a cell surface, free in solution, or located intracellularly. One method of drug screening utilizes eukaryotic
25 or prokaryotic host cells which are stably transformed with recombinant nucleic acids expressing the polypeptide or fragment. Drugs are screened against such transformed cells in competitive binding assays. One may measure, for example, the formulation of complexes between the agent being tested and a polypeptide of the present invention.

30 Thus, the present invention provides methods of screening for drugs or any other agents which affect activities mediated by the polypeptides of the present invention. These methods comprise contacting such an agent with a polypeptide of the

present invention or a fragment thereof and assaying for the presence of a complex between the agent and the polypeptide or a fragment thereof, by methods well known in the art. In such a competitive binding assay, the agents to screen are typically labeled. Following incubation, free agent is separated from that present in bound form, and the amount of free or uncomplexed label is a measure of the ability of a particular agent to bind to the polypeptides of the present invention.

Another technique for drug screening provides high throughput screening for compounds having suitable binding affinity to the polypeptides of the present invention, and is described in great detail in European Patent Application 84/03564, published on September 13, 1984, which is incorporated herein by reference herein. Briefly stated, large numbers of different small peptide test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The peptide test compounds are reacted with polypeptides of the present invention and washed. Bound polypeptides are then detected by methods well known in the art. Purified polypeptides are coated directly onto plates for use in the aforementioned drug screening techniques. In addition, non-neutralizing antibodies may be used to capture the peptide and immobilize it on the solid support.

This invention also contemplates the use of competitive drug screening assays in which neutralizing antibodies capable of binding polypeptides of the present invention specifically compete with a test compound for binding to the polypeptides or fragments thereof. In this manner, the antibodies are used to detect the presence of any peptide which shares one or more antigenic epitopes with a polypeptide of the invention.

Polypeptides of the Invention Binding Peptides and Other Molecules

The invention also encompasses screening methods for identifying polypeptides and nonpolypeptides that bind polypeptides of the invention, and the polypeptide of the invention binding molecules identified thereby. These binding molecules are useful, for example, as agonists and antagonists of the polypeptides of the invention. Such agonists and antagonists can be used, in accordance with the invention, in the therapeutic embodiments described in detail, below.

This method comprises the steps of:

- a. contacting a polypeptide of the invention with a plurality of molecules; and
- b. identifying a molecule that binds the polypeptide of the invention.

The step of contacting the polypeptide of the invention with the plurality of molecules may be effected in a number of ways. For example, one may contemplate
5 immobilizing the polypeptide of the invention on a solid support and bringing a solution of the plurality of molecules in contact with the immobilized polypeptide of the invention. Such a procedure would be akin to an affinity chromatographic process, with the affinity matrix being comprised of the immobilized polypeptide of the invention. The molecules having a selective affinity for the polypeptide of the
10 invention can then be purified by affinity selection. The nature of the solid support, process for attachment of the polypeptide of the invention to the solid support, solvent, and conditions of the affinity isolation or selection are largely conventional and well known to those of ordinary skill in the art.

Alternatively, one may also separate a plurality of polypeptides into
15 substantially separate fractions comprising a subset of or individual polypeptides. For instance, one can separate the plurality of polypeptides by gel electrophoresis, column chromatography, or like method known to those of ordinary skill for the separation of polypeptides. The individual polypeptides can also be produced by a transformed host cell in such a way as to be expressed on or about its outer surface (e.g., a recombinant
20 phage). Individual isolates can then be "probed" by the polypeptide of the invention, optionally in the presence of an inducer should one be required for expression, to determine if any selective affinity interaction takes place between the polypeptide of the invention and the individual clone. Prior to contacting the polypeptide of the invention with each fraction comprising individual polypeptides, the polypeptides
25 could first be transferred to a solid support for additional convenience. Such a solid support may simply be a piece of filter membrane, such as one made of nitrocellulose or nylon. In this manner, positive clones could be identified from a collection of transformed host cells of an expression library, which harbor a DNA construct encoding a polypeptide having a selective affinity for a polypeptide of the invention.
30 Furthermore, the amino acid sequence of the polypeptide having a selective affinity for the polypeptide of the invention can be determined directly by conventional means or the coding sequence of the DNA encoding the polypeptide can frequently be

determined more conveniently. The primary sequence can then be deduced from the corresponding DNA sequence. If the amino acid sequence is to be determined from the polypeptide itself, one may use microsequencing techniques. The sequencing technique may include mass spectroscopy.

5 In certain situations, it may be desirable to wash away any unbound polypeptide of the invention, or alternatively, unbound polypeptides, from a mixture of the polypeptide of the invention and the plurality of polypeptides prior to attempting to determine or to detect the presence of a selective affinity interaction. Such a wash step may be particularly desirable when the polypeptide of the invention
10 or the plurality of polypeptides is bound to a solid support.

 The plurality of molecules provided according to this method may be provided by way of diversity libraries, such as random or combinatorial peptide or nonpeptide libraries which can be screened for molecules that specifically bind to a polypeptide of the invention. Many libraries are known in the art that can be used, e.g., chemically
15 synthesized libraries, recombinant (e.g., phage display libraries), and in vitro translation-based libraries. Examples of chemically synthesized libraries are described in Fodor et al., 1991, *Science* 251:767-773; Houghten et al., 1991, *Nature* 354:84-86; Lam et al., 1991, *Nature* 354:82-84; Medynski, 1994, *Bio/Technology* 12:709-710; Gallop et al., 1994, *J. Medicinal Chemistry* 37(9):1233-1251; Ohlmeyer
20 et al., 1993, *Proc. Natl. Acad. Sci. USA* 90:10922-10926; Erb et al., 1994, *Proc. Natl. Acad. Sci. USA* 91:11422-11426; Houghten et al., 1992, *Biotechniques* 13:412; Jayawickreme et al., 1994, *Proc. Natl. Acad. Sci. USA* 91:1614-1618; Salmon et al., 1993, *Proc. Natl. Acad. Sci. USA* 90:11708-11712; PCT Publication No. WO 93/20242; and Brenner and Lerner, 1992, *Proc. Natl. Acad. Sci. USA* 89:5381-5383.

25 Examples of phage display libraries are described in Scott and Smith, 1990, *Science* 249:386-390; Devlin et al., 1990, *Science*, 249:404-406; Christian, R. B., et al., 1992, *J. Mol. Biol.* 227:711-718; Lenstra, 1992, *J. Immunol. Meth.* 152:149-157; Kay et al., 1993, *Gene* 128:59-65; and PCT Publication No. WO 94/18318 dated Aug. 18, 1994.

30 In vitro translation-based libraries include but are not limited to those described in PCT Publication No. WO 91/05058 dated Apr. 18, 1991; and Mattheakis et al., 1994, *Proc. Natl. Acad. Sci. USA* 91:9022-9026.

By way of examples of nonpeptide libraries, a benzodiazepine library (see e.g., Bunin et al., 1994, Proc. Natl. Acad. Sci. USA 91:4708-4712) can be adapted for use. Peptoid libraries (Simon et al., 1992, Proc. Natl. Acad. Sci. USA 89:9367-9371) can also be used. Another example of a library that can be used, in which the amide
5 functionalities in peptides have been permethylated to generate a chemically transformed combinatorial library, is described by Ostresh et al. (1994, Proc. Natl. Acad. Sci. USA 91:11138-11142).

The variety of non-peptide libraries that are useful in the present invention is great. For example, Ecker and Crooke, 1995, Bio/Technology 13:351-360 list
10 benzodiazepines, hydantoins, piperazinediones, biphenyls, sugar analogs, beta-mercaptoketones, arylacetic acids, acylpiperidines, benzopyrans, cubanes, xanthines, aminimides, and oxazolones as among the chemical species that form the basis of various libraries.

Non-peptide libraries can be classified broadly into two types: decorated
15 monomers and oligomers. Decorated monomer libraries employ a relatively simple scaffold structure upon which a variety functional groups is added. Often the scaffold will be a molecule with a known useful pharmacological activity. For example, the scaffold might be the benzodiazepine structure.

Non-peptide oligomer libraries utilize a large number of monomers that are
20 assembled together in ways that create new shapes that depend on the order of the monomers. Among the monomer units that have been used are carbamates, pyrrolinones, and morpholinos. Peptoids, peptide-like oligomers in which the side chain is attached to the alpha amino group rather than the alpha carbon, form the basis of another version of non-peptide oligomer libraries. The first non-peptide oligomer
25 libraries utilized a single type of monomer and thus contained a repeating backbone. Recent libraries have utilized more than one monomer, giving the libraries added flexibility.

Screening the libraries can be accomplished by any of a variety of commonly known methods. See, e.g., the following references, which disclose screening of
30 peptide libraries: Parmley and Smith, 1989, Adv. Exp. Med. Biol. 251:215-218; Scott and Smith, 1990, Science 249:386-390; Fowlkes et al., 1992; BioTechniques 13:422-427; Oldenburg et al., 1992, Proc. Natl. Acad. Sci. USA 89:5393-5397; Yu et al.,

1994, Cell 76:933-945; Staudt et al., 1988, Science 241:577-580; Bock et al., 1992, Nature 355:564-566; Tuerk et al., 1992, Proc. Natl. Acad. Sci. USA 89:6988-6992; Ellington et al., 1992, Nature 355:850-852; U.S. Pat. No. 5,096,815, U.S. Pat. No. 5,223,409, and U.S. Pat. No. 5,198,346, all to Ladner et al.; Rebar and Pabo, 1993, Science 263:671-673; and CT Publication No. WO 94/18318.

In a specific embodiment, screening to identify a molecule that binds a polypeptide of the invention can be carried out by contacting the library members with a polypeptide of the invention immobilized on a solid phase and harvesting those library members that bind to the polypeptide of the invention. Examples of such screening methods, termed "panning" techniques are described by way of example in Parmley and Smith, 1988, Gene 73:305-318; Fowlkes et al., 1992, BioTechniques 13:422-427; PCT Publication No. WO 94/18318; and in references cited herein.

In another embodiment, the two-hybrid system for selecting interacting proteins in yeast (Fields and Song, 1989, Nature 340:245-246; Chien et al., 1991, Proc. Natl. Acad. Sci. USA 88:9578-9582) can be used to identify molecules that specifically bind to a polypeptide of the invention.

Where the polypeptide of the invention binding molecule is a polypeptide, the polypeptide can be conveniently selected from any peptide library, including random peptide libraries, combinatorial peptide libraries, or biased peptide libraries. The term "biased" is used herein to mean that the method of generating the library is manipulated so as to restrict one or more parameters that govern the diversity of the resulting collection of molecules, in this case peptides.

Thus, a truly random peptide library would generate a collection of peptides in which the probability of finding a particular amino acid at a given position of the peptide is the same for all 20 amino acids. A bias can be introduced into the library, however, by specifying, for example, that a lysine occur every fifth amino acid or that positions 4, 8, and 9 of a decapeptide library be fixed to include only arginine. Clearly, many types of biases can be contemplated, and the present invention is not restricted to any particular bias. Furthermore, the present invention contemplates specific types of peptide libraries, such as phage displayed peptide libraries and those that utilize a DNA construct comprising a lambda phage vector with a DNA insert.

As mentioned above, in the case of a polypeptide of the invention binding

molecule that is a polypeptide, the polypeptide may have about 6 to less than about 60 amino acid residues, preferably about 6 to about 10 amino acid residues, and most preferably, about 6 to about 22 amino acids. In another embodiment, a polypeptide of the invention binding polypeptide has in the range of 15-100 amino acids, or 20-50 amino acids.

The selected polypeptide of the invention binding polypeptide can be obtained by chemical synthesis or recombinant expression.

Antisense And Ribozyme (Antagonists)

In specific embodiments, antagonists according to the present invention are nucleic acids corresponding to the sequences contained in SEQ ID NO:X, or the complementary strand thereof, and/or to nucleotide sequences contained a deposited clone. In one embodiment, antisense sequence is generated internally by the organism, in another embodiment, the antisense sequence is separately administered (see, for example, O'Connor, Neurochem., 56:560 (1991). Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988). Antisense technology can be used to control gene expression through antisense DNA or RNA, or through triple-helix formation. Antisense techniques are discussed for example, in Okano, Neurochem., 56:560 (1991); Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988). Triple helix formation is discussed in, for instance, Lee et al., Nucleic Acids Research, 6:3073 (1979); Cooney et al., Science, 241:456 (1988); and Dervan et al., Science, 251:1300 (1991). The methods are based on binding of a polynucleotide to a complementary DNA or RNA.

For example, the use of c-myc and c-myb antisense RNA constructs to inhibit the growth of the non-lymphocytic leukemia cell line HL-60 and other cell lines was previously described. (Wickstrom et al. (1988); Anfossi et al. (1989)). These experiments were performed in vitro by incubating cells with the oligoribonucleotide. A similar procedure for in vivo use is described in WO 91/15580. Briefly, a pair of oligonucleotides for a given antisense RNA is produced as follows: A sequence complimentary to the first 15 bases of the open reading frame is flanked by an EcoRI site on the 5' end and a HindIII site on the 3' end. Next, the pair of oligonucleotides is

heated at 90°C for one minute and then annealed in 2X ligation buffer (20mM TRIS HCl pH 7.5, 10mM MgCl₂, 10mM dithiothreitol (DTT) and 0.2 mM ATP) and then ligated to the EcoR1/Hind III site of the retroviral vector PMV7 (WO 91/15580).

For example, the 5' coding portion of a polynucleotide that encodes the mature polypeptide of the present invention may be used to design an antisense RNA oligonucleotide of from about 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription thereby preventing transcription and the production of the receptor. The antisense RNA oligonucleotide hybridizes to the mRNA in vivo and blocks translation of the mRNA molecule into receptor polypeptide.

In one embodiment, the antisense nucleic acid of the invention is produced intracellularly by transcription from an exogenous sequence. For example, a vector or a portion thereof, is transcribed, producing an antisense nucleic acid (RNA) of the invention. Such a vector would contain a sequence encoding the antisense nucleic acid of the invention. Such a vector can remain episomal or become chromosomally integrated, as long as it can be transcribed to produce the desired antisense RNA. Such vectors can be constructed by recombinant DNA technology methods standard in the art. Vectors can be plasmid, viral, or others known in the art, used for replication and expression in vertebrate cells. Expression of the sequence encoding a polypeptide of the invention, or fragments thereof, can be by any promoter known in the art to act in vertebrate, preferably human cells. Such promoters can be inducible or constitutive. Such promoters include, but are not limited to, the SV40 early promoter region (Bernoist and Chambon, *Nature*, 29:304-310 (1981)), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto et al., *Cell*, 22:787-797 (1980)), the herpes thymidine promoter (Wagner et al., *Proc. Natl. Acad. Sci. U.S.A.*, 78:1441-1445 (1981)), the regulatory sequences of the metallothionein gene (Brinster et al., *Nature*, 296:39-42 (1982)), etc.

The antisense nucleic acids of the invention comprise a sequence complementary to at least a portion of an RNA transcript of a gene of interest. However, absolute complementarity, although preferred, is not required. A sequence "complementary to at least a portion of an RNA," referred to herein, means a sequence having sufficient complementarity to be able to hybridize with the RNA,

forming a stable duplex; in the case of double stranded antisense nucleic acids of the invention, a single strand of the duplex DNA may thus be tested, or triplex formation may be assayed. The ability to hybridize will depend on both the degree of complementarity and the length of the antisense nucleic acid. Generally, the larger the hybridizing nucleic acid, the more base mismatches with a RNA sequence of the invention it may contain and still form a stable duplex (or triplex as the case may be). One skilled in the art can ascertain a tolerable degree of mismatch by use of standard procedures to determine the melting point of the hybridized complex.

Oligonucleotides that are complementary to the 5' end of the message, *e.g.*, the 5' untranslated sequence up to and including the AUG initiation codon, should work most efficiently at inhibiting translation. However, sequences complementary to the 3' untranslated sequences of mRNAs have been shown to be effective at inhibiting translation of mRNAs as well. See generally, Wagner, R., *Nature*, 372:333-335 (1994). Thus, oligonucleotides complementary to either the 5' - or 3' - non-translated, non-coding regions of a polynucleotide sequence of the invention could be used in an antisense approach to inhibit translation of endogenous mRNA. Oligonucleotides complementary to the 5' untranslated region of the mRNA should include the complement of the AUG start codon. Antisense oligonucleotides complementary to mRNA coding regions are less efficient inhibitors of translation but could be used in accordance with the invention. Whether designed to hybridize to the 5' -, 3' - or coding region of mRNA, antisense nucleic acids should be at least six nucleotides in length, and are preferably oligonucleotides ranging from 6 to about 50 nucleotides in length. In specific aspects the oligonucleotide is at least 10 nucleotides, at least 17 nucleotides, at least 25 nucleotides or at least 50 nucleotides.

The polynucleotides of the invention can be DNA or RNA or chimeric mixtures or derivatives or modified versions thereof, single-stranded or double-stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone, for example, to improve stability of the molecule, hybridization, etc. The oligonucleotide may include other appended groups such as peptides (*e.g.*, for targeting host cell receptors *in vivo*), or agents facilitating transport across the cell membrane (see, *e.g.*, Letsinger et al., *Proc. Natl. Acad. Sci. U.S.A.* 86:6553-6556 (1989); Lemaitre et al., *Proc. Natl. Acad. Sci.*, 84:648-652 (1987); PCT Publication

NO: WO88/09810, published December 15, 1988) or the blood-brain barrier (see, e.g., PCT Publication NO: WO89/10134, published April 25, 1988), hybridization-triggered cleavage agents. (See, e.g., Krol et al., BioTechniques, 6:958-976 (1988)) or intercalating agents. (See, e.g., Zon, Pharm. Res., 5:539-549 (1988)). To this end,
5 the oligonucleotide may be conjugated to another molecule, e.g., a peptide, hybridization triggered cross-linking agent, transport agent, hybridization-triggered cleavage agent, etc.

The antisense oligonucleotide may comprise at least one modified base moiety which is selected from the group including, but not limited to, 5-fluorouracil,
10 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xantine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine,
15 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil,
20 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine.

The antisense oligonucleotide may also comprise at least one modified sugar moiety selected from the group including, but not limited to, arabinose,
25 2-fluoroarabinose, xylulose, and hexose.

In yet another embodiment, the antisense oligonucleotide comprises at least one modified phosphate backbone selected from the group including, but not limited to, a phosphorothioate, a phosphorodithioate, a phosphoramidothioate, a phosphoramidate, a phosphordiamidate, a methylphosphonate, an alkyl
30 phosphotriester, and a formacetal or analog thereof.

In yet another embodiment, the antisense oligonucleotide is an a-anomeric oligonucleotide. An a-anomeric oligonucleotide forms specific double-stranded

hybrids with complementary RNA in which, contrary to the usual b-units, the strands run parallel to each other (Gautier et al., Nucl. Acids Res., 15:6625-6641 (1987)).

The oligonucleotide is a 2'-O-methylribonucleotide (Inoue et al., Nucl. Acids Res., 15:6131-6148 (1987)), or a chimeric RNA-DNA analogue (Inoue et al., FEBS Lett. 5 215:327-330 (1987)).

Polynucleotides of the invention may be synthesized by standard methods known in the art, e.g. by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, etc.). As examples, phosphorothioate oligonucleotides may be synthesized by the method of Stein et al. 10 (Nucl. Acids Res., 16:3209 (1988)), methylphosphonate oligonucleotides can be prepared by use of controlled pore glass polymer supports (Sarin et al., Proc. Natl. Acad. Sci. U.S.A., 85:7448-7451 (1988)), etc.

While antisense nucleotides complementary to the coding region sequence of the invention could be used, those complementary to the transcribed untranslated 15 region are most preferred.

Potential antagonists according to the invention also include catalytic RNA, or a ribozyme (See, e.g., PCT International Publication WO 90/11364, published October 4, 1990; Sarver et al, Science, 247:1222-1225 (1990). While ribozymes that cleave mRNA at site specific recognition sequences can be used to destroy mRNAs 20 corresponding to the polynucleotides of the invention, the use of hammerhead ribozymes is preferred. Hammerhead ribozymes cleave mRNAs at locations dictated by flanking regions that form complementary base pairs with the target mRNA. The sole requirement is that the target mRNA have the following sequence of two bases: 5' -UG-3' . The construction and production of hammerhead ribozymes is well 25 known in the art and is described more fully in Haseloff and Gerlach, Nature, 334:585-591 (1988). There are numerous potential hammerhead ribozyme cleavage sites within each nucleotide sequence disclosed in the sequence listing. Preferably, the ribozyme is engineered so that the cleavage recognition site is located near the 5' end of the mRNA corresponding to the polynucleotides of the invention; i.e., to 30 increase efficiency and minimize the intracellular accumulation of non-functional mRNA transcripts.

As in the antisense approach, the ribozymes of the invention can be composed of modified oligonucleotides (e.g. for improved stability, targeting, etc.) and should be delivered to cells which express the polynucleotides of the invention in vivo.

DNA constructs encoding the ribozyme may be introduced into the cell in the same

5 manner as described above for the introduction of antisense encoding DNA. A preferred method of delivery involves using a DNA construct "encoding" the ribozyme under the control of a strong constitutive promoter, such as, for example, pol III or pol II promoter, so that transfected cells will produce sufficient quantities of the ribozyme to destroy endogenous messages and inhibit translation. Since
10 ribozymes unlike antisense molecules, are catalytic, a lower intracellular concentration is required for efficiency.

Antagonist/agonist compounds may be employed to inhibit the cell growth and proliferation effects of the polypeptides of the present invention on neoplastic cells and tissues, i.e. stimulation of angiogenesis of tumors, and, therefore, retard or
15 prevent abnormal cellular growth and proliferation, for example, in tumor formation or growth.

The antagonist/agonist may also be employed to prevent hyper-vascular diseases, and prevent the proliferation of epithelial lens cells after extracapsular cataract surgery. Prevention of the mitogenic activity of the polypeptides of the
20 present invention may also be desirable in cases such as restenosis after balloon angioplasty.

The antagonist/agonist may also be employed to prevent the growth of scar tissue during wound healing.

The antagonist/agonist may also be employed to treat, prevent, and/or
25 diagnose the diseases described herein.

Thus, the invention provides a method of treating or preventing diseases, disorders, and/or conditions, including but not limited to the diseases, disorders, and/or conditions listed throughout this application, associated with overexpression of a polynucleotide of the present invention by administering to a patient (a) an antisense
30 molecule directed to the polynucleotide of the present invention, and/or (b) a ribozyme directed to the polynucleotide of the present invention.

invention, and/or (b) a ribozyme directed to the polynucleotide of the present invention

Other Activities

5 The polypeptide of the present invention, as a result of the ability to stimulate vascular endothelial cell growth, may be employed in treatment for stimulating re-vascularization of ischemic tissues due to various disease conditions such as thrombosis, arteriosclerosis, and other cardiovascular conditions. These polypeptide may also be employed to stimulate angiogenesis and limb regeneration, as discussed
10 above.

 The polypeptide may also be employed for treating wounds due to injuries, burns, post-operative tissue repair, and ulcers since they are mitogenic to various cells of different origins, such as fibroblast cells and skeletal muscle cells, and therefore, facilitate the repair or replacement of damaged or diseased tissue.

15 The polypeptide of the present invention may also be employed stimulate neuronal growth and to treat, prevent, and/or diagnose neuronal damage which occurs in certain neuronal disorders or neuro-degenerative conditions such as Alzheimer's disease, Parkinson's disease, and AIDS-related complex. The polypeptide of the invention may have the ability to stimulate chondrocyte growth, therefore, they may
20 be employed to enhance bone and periodontal regeneration and aid in tissue transplants or bone grafts.

 The polypeptide of the present invention may be also be employed to prevent skin aging due to sunburn by stimulating keratinocyte growth.

 The polypeptide of the invention may also be employed for preventing hair
25 loss, since FGF family members activate hair-forming cells and promotes melanocyte growth. Along the same lines, the polypeptides of the present invention may be employed to stimulate growth and differentiation of hematopoietic cells and bone marrow cells when used in combination with other cytokines.

 The polypeptide of the invention may also be employed to maintain organs
30 before transplantation or for supporting cell culture of primary tissues.

 The polypeptide of the present invention may also be employed for inducing tissue of mesodermal origin to differentiate in early embryos.

The polypeptide or polynucleotides and/or agonist or antagonists of the present invention may also increase or decrease the differentiation or proliferation of embryonic stem cells, besides, as discussed above, hematopoietic lineage.

5 The polypeptide or polynucleotides and/or agonist or antagonists of the present invention may also be used to modulate mammalian characteristics, such as body height, weight, hair color, eye color, skin, percentage of adipose tissue, pigmentation, size, and shape (e.g., cosmetic surgery). Similarly, polypeptides or polynucleotides and/or agonist or antagonists of the present invention may be used to
10 modulate mammalian metabolism affecting catabolism, anabolism, processing, utilization, and storage of energy.

Polypeptide or polynucleotides and/or agonist or antagonists of the present invention may be used to change a mammal's mental state or physical state by influencing biorhythms, cardiac rhythms, depression (including depressive diseases,
15 disorders, and/or conditions), tendency for violence, tolerance for pain, reproductive capabilities (preferably by Activin or Inhibin-like activity), hormonal or endocrine levels, appetite, libido, memory, stress, or other cognitive qualities.

Polypeptide or polynucleotides and/or agonist or antagonists of the present invention may also be used as a food additive or preservative, such as to increase or
20 decrease storage capabilities, fat content, lipid, protein, carbohydrate, vitamins, minerals, cofactors or other nutritional components.

Other Preferred Embodiments

Other preferred embodiments of the claimed invention include an isolated
25 nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least about 50 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1.

Also preferred is a nucleic acid molecule wherein said sequence of contiguous nucleotides is included in the nucleotide sequence of SEQ ID NO:X in the range of
30 positions beginning with the nucleotide at about the position of the 5' Nucleotide of

the Clone Sequence and ending with the nucleotide at about the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X in Table 1.

Also preferred is a nucleic acid molecule wherein said sequence of contiguous nucleotides is included in the nucleotide sequence of SEQ ID NO:X in the range of
5 positions beginning with the nucleotide at about the position of the 5' Nucleotide of the Start Codon and ending with the nucleotide at about the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X in Table 1.

Similarly preferred is a nucleic acid molecule wherein said sequence of contiguous nucleotides is included in the nucleotide sequence of SEQ ID NO:X in the
10 range of positions beginning with the nucleotide at about the position of the 5' Nucleotide of the First Amino Acid of the Signal Peptide and ending with the nucleotide at about the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X in Table 1.

Also preferred is an isolated nucleic acid molecule comprising a nucleotide
15 sequence which is at least 95% identical to a sequence of at least about 150 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X.

Further preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least about 500 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X.

A further preferred embodiment is a nucleic acid molecule comprising a
20 nucleotide sequence which is at least 95% identical to the nucleotide sequence of SEQ ID NO:X beginning with the nucleotide at about the position of the 5' Nucleotide of the First Amino Acid of the Signal Peptide and ending with the nucleotide at about the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X
25 in Table 1.

A further preferred embodiment is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to the complete nucleotide sequence of SEQ ID NO:X.

Also preferred is an isolated nucleic acid molecule which hybridizes under
30 stringent hybridization conditions to a nucleic acid molecule, wherein said nucleic acid molecule which hybridizes does not hybridize under stringent hybridization

conditions to a nucleic acid molecule having a nucleotide sequence consisting of only A residues or of only T residues.

Also preferred is a composition of matter comprising a DNA molecule which comprises a human cDNA clone identified by a cDNA Clone Identifier in Table 1,
5 which DNA molecule is contained in the material deposited with the American Type Culture Collection and given the ATCC Deposit Number shown in Table 1 for said cDNA Clone Identifier.

Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least 50 contiguous
10 nucleotides in the nucleotide sequence of a human cDNA clone identified by a cDNA Clone Identifier in Table 1, which DNA molecule is contained in the deposit given the ATCC Deposit Number shown in Table 1.

Also preferred is an isolated nucleic acid molecule, wherein said sequence of at least 50 contiguous nucleotides is included in the nucleotide sequence of the
15 complete open reading frame sequence encoded by said human cDNA clone.

Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to sequence of at least 150 contiguous nucleotides in the nucleotide sequence encoded by said human cDNA clone.

A further preferred embodiment is an isolated nucleic acid molecule
20 comprising a nucleotide sequence which is at least 95% identical to sequence of at least 500 contiguous nucleotides in the nucleotide sequence encoded by said human cDNA clone.

A further preferred embodiment is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to the complete
25 nucleotide sequence encoded by said human cDNA clone.

A further preferred embodiment is a method for detecting in a biological sample a nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X
30 wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table

1; which method comprises a step of comparing a nucleotide sequence of at least one nucleic acid molecule in said sample with a sequence selected from said group and determining whether the sequence of said nucleic acid molecule in said sample is at least 95% identical to said selected sequence.

5 Also preferred is the above method wherein said step of comparing sequences comprises determining the extent of nucleic acid hybridization between nucleic acid molecules in said sample and a nucleic acid molecule comprising said sequence selected from said group. Similarly, also preferred is the above method wherein said step of comparing sequences is performed by comparing the nucleotide sequence
10 determined from a nucleic acid molecule in said sample with said sequence selected from said group. The nucleic acid molecules can comprise DNA molecules or RNA molecules.

A further preferred embodiment is a method for identifying the species, tissue or cell type of a biological sample which method comprises a step of detecting nucleic
15 acid molecules in said sample, if any, comprising a nucleotide sequence that is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained
20 in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

The method for identifying the species, tissue or cell type of a biological sample can comprise a step of detecting nucleic acid molecules comprising a nucleotide sequence in a panel of at least two nucleotide sequences, wherein at least
25 one sequence in said panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from said group.

Also preferred is a method for diagnosing in a subject a pathological condition associated with abnormal structure or expression of a gene encoding a secreted protein identified in Table 1, which method comprises a step of detecting in a
30 biological sample obtained from said subject nucleic acid molecules, if any, comprising a nucleotide sequence that is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a

nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

5 The method for diagnosing a pathological condition can comprise a step of detecting nucleic acid molecules comprising a nucleotide sequence in a panel of at least two nucleotide sequences, wherein at least one sequence in said panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from said group.

10 Also preferred is a composition of matter comprising isolated nucleic acid molecules wherein the nucleotide sequences of said nucleic acid molecules comprise a panel of at least two nucleotide sequences, wherein at least one sequence in said panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID
15 NO:X wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1. The nucleic acid molecules can comprise DNA molecules or RNA molecules.

20 Also preferred is an isolated polypeptide comprising an amino acid sequence at least 90% identical to a sequence of at least about 10 contiguous amino acids in the amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1.

 Also preferred is a polypeptide, wherein said sequence of contiguous amino acids is included in the amino acid sequence of SEQ ID NO:Y in the range of
25 positions beginning with the residue at about the position of the First Amino Acid of the Secreted Portion and ending with the residue at about the Last Amino Acid of the Open Reading Frame as set forth for SEQ ID NO:Y in Table 1.

 Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 30 contiguous amino acids in the
30 amino acid sequence of SEQ ID NO:Y.

Further preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 100 contiguous amino acids in the amino acid sequence of SEQ ID NO:Y.

Further preferred is an isolated polypeptide comprising an amino acid
5 sequence at least 95% identical to the complete amino acid sequence of SEQ ID NO:Y.

Further preferred is an isolated polypeptide comprising an amino acid sequence at least 90% identical to a sequence of at least about 10 contiguous amino acids in the complete amino acid sequence of a secreted protein encoded by a human
10 cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is a polypeptide wherein said sequence of contiguous amino acids is included in the amino acid sequence of a secreted portion of the secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in
15 Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 30 contiguous amino acids in the amino acid sequence of the secreted portion of the protein encoded by a human cDNA
20 clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 100 contiguous amino acids in the amino acid sequence of the secreted portion of the protein encoded by a human
25 cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to the amino acid sequence of the secreted portion of the protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1
30 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Further preferred is an isolated antibody which binds specifically to a polypeptide comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Further preferred is a method for detecting in a biological sample a polypeptide comprising an amino acid sequence which is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1; which method comprises a step of comparing an amino acid sequence of at least one polypeptide molecule in said sample with a sequence selected from said group and determining whether the sequence of said polypeptide molecule in said sample is at least 90% identical to said sequence of at least 10 contiguous amino acids.

Also preferred is the above method wherein said step of comparing an amino acid sequence of at least one polypeptide molecule in said sample with a sequence selected from said group comprises determining the extent of specific binding of polypeptides in said sample to an antibody which binds specifically to a polypeptide comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is the above method wherein said step of comparing sequences is performed by comparing the amino acid sequence determined from a polypeptide molecule in said sample with said sequence selected from said group.

Also preferred is a method for identifying the species, tissue or cell type of a biological sample which method comprises a step of detecting polypeptide molecules in said sample, if any, comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is the above method for identifying the species, tissue or cell type of a biological sample, which method comprises a step of detecting polypeptide molecules comprising an amino acid sequence in a panel of at least two amino acid sequences, wherein at least one sequence in said panel is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the above group.

Also preferred is a method for diagnosing in a subject a pathological condition associated with abnormal structure or expression of a gene encoding a secreted protein identified in Table 1, which method comprises a step of detecting in a biological sample obtained from said subject polypeptide molecules comprising an amino acid sequence in a panel of at least two amino acid sequences, wherein at least one sequence in said panel is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

In any of these methods, the step of detecting said polypeptide molecules includes using an antibody.

Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a nucleotide sequence encoding a polypeptide wherein said polypeptide comprises an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence

selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number
5 shown for said cDNA clone in Table 1.

Also preferred is an isolated nucleic acid molecule, wherein said nucleotide sequence encoding a polypeptide has been optimized for expression of said polypeptide in a prokaryotic host.

Also preferred is an isolated nucleic acid molecule, wherein said polypeptide
10 comprises an amino acid sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Further preferred is a method of making a recombinant vector comprising
15 inserting any of the above isolated nucleic acid molecule into a vector. Also preferred is the recombinant vector produced by this method. Also preferred is a method of making a recombinant host cell comprising introducing the vector into a host cell, as well as the recombinant host cell produced by this method.

Also preferred is a method of making an isolated polypeptide comprising
20 culturing this recombinant host cell under conditions such that said polypeptide is expressed and recovering said polypeptide. Also preferred is this method of making an isolated polypeptide, wherein said recombinant host cell is a eukaryotic cell and said polypeptide is a secreted portion of a human secreted protein comprising an
25 amino acid sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y beginning with the residue at the position of the First Amino Acid of the Secreted Portion of SEQ ID NO:Y wherein Y is an integer set forth in Table 1 and said position of the First Amino Acid of the Secreted Portion of SEQ ID NO:Y is defined in Table 1; and an amino acid sequence of a secreted portion of a protein
30 encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1. The isolated polypeptide produced by this method is also preferred.

Also preferred is a method of treatment of an individual in need of an increased level of a secreted protein activity, which method comprises administering to such an individual a pharmaceutical composition comprising an amount of an isolated polypeptide, polynucleotide, or antibody of the claimed invention effective to
 5 increase the level of said protein activity in said individual.

The above-recited applications have uses in a wide variety of hosts. Such hosts include, but are not limited to, human, murine, rabbit, goat, guinea pig, camel, horse, mouse, rat, hamster, pig, micro-pig, chicken, goat, cow, sheep, dog, cat, non-human primate, and human. In specific embodiments, the host is a mouse, rabbit,
 10 goat, guinea pig, chicken, rat, hamster, pig, sheep, dog or cat. In preferred embodiments, the host is a mammal. In most preferred embodiments, the host is a human.

In specific embodiments of the invention, for each "Contig ID" listed in the fourth column of Table 7, preferably excluded are one or more polynucleotides
 15 comprising, or alternatively consisting of, a nucleotide sequence referenced in the fifth column of Table 7 and described by the general formula of a-b, whereas a and b are uniquely determined for the corresponding SEQ ID NO:X referred to in column 3 of Table 7. Further specific embodiments are directed to polynucleotide sequences excluding one, two, three, four, or more of the specific polynucleotide sequences
 20 referred to in the fifth column of Table 7. In no way is this listing meant to encompass all of the sequences which may be excluded by the general formula, it is just a representative example. All references available through these accessions are hereby incorporated by reference in their entirety.

TABLE 7

| Gene No. | cDNA Clone ID | NT SEQ ID NO: X | Contig ID | Public Accession Numbers |
|----------|---------------|-----------------|-----------|---|
| 1 | HWEAC77 | 11 | 984653 | AI085609 and AA429294. |
| 2 | HRAEH37 | 40 | 904323 | H17862, AI222550, AI207344, AW068428, AA397646, AA581204, R88190, AA864871, T60301, R88480, T74353, AA206937, AW295179, AA058486, AW379072, R87289, F12597, AI862896, AA340088, AA333098, AW337695, AI636228, AI199183, T29710, |

| | | | | |
|---|---------|----|--------|---|
| | | | | AI245728, AA527474, AI658678, and M95549. |
| 2 | HRAEH37 | 41 | 904324 | AI860759, AI884486, AW193525, AA846599, AI554646, AI799706, AA523949, AW001228, H17863, AI003564, AI524226, AA448141, AA767653, AI567123, AA206837, AA773851, AA993496, AW168928, H29750, AI679819, AI889279, AA421148, AI679243, AI923054, AA447998, AI290066, AA058367, AI358091, W89146, AA576924, AW067953, AA599007, AA700339, AA705404, AW238218, AA564967, H61752, AA969775, AA652622, H61751, AA593588, W89145, AW169777, AI940516, AI696808, AI277898, T53109, AA402926, AI805502, AI249113, W33071, T49465, AA887759, AA581976, F10215, AA350860, AI866503, AI804832, T49466, T89112, AA887749, AW351458, C14005, AA299510, AI860478, AI658678, AA527474, AW295179, AI199183, AI619916, T61678, R23078, AA364590, AI214409, AA581255, AI701266, AA715116, AW188721, W35420, AW087438, AI207344, AW375741, AA570117, AA471194, T73430, AA548627, T53110, AI884326, AI222550, AA581204, T29710, AI245728, AI800473, AA827697, AW169784, AA767924, AI539260, AI565147, AI559752, AA503384, AL047100, M95549, E01349, I28326, AF177401, I08319, A12558, AL117416, AF090903, AL023657, D83032, AL136884, X95310, and AL137533. |
| 4 | HDQHY04 | 14 | 903650 | AI871408, AA278445, AA479276, AI038602, N52018, AA706346, R80332, AI559411, AA305170, AI915377, AW440856, AI671553, AI809685, AA744961, AA309735, AA479371, AI940805, AI928450, N66171, AW103125, AA333461, AA857635, T80638, T80747, AA370795, AW088932, AA344676, AA279061, AA331630, R80226, AL119457, AL119399, AI270183, AI491775, AL042544, AL042382, AI863321, AI619502, AL079794, AI536685, AI538716, AI421903, AI862139, AW131954, AI869367, AI857296, AI863014, AI799199, AI499463, AI920968, AI500553, AI696398, AI564719, AW002342, AI282903, AI628316, AI572787, AI873644, AI567351, AI811344, AI637584, AI476046, AW104724, AW148320, AI889376, AW050522, AI654750, AI678989, AI590227, AL041772, AW131294, AI702406, AW075381, AI269205, AI434468, AI619716, AI569328, AW169527, AI280747, AI432969, AI702433, AI824557, AI828731, AI433157, AW075413, AI923357, AI702068, AI636719, AI539153, AI280637, AI287326, AI866002, AW169653, AI699857, AI590021, AI824648, AI499131, |

| | | | | |
|--|--|--|--|---|
| | | | | AI572676, AI569583, AI888501, AI950892, AI366900, AW102785, AI540832, AI537303, AI453322, AI801608, AI560012, AL036187, AI475451, AI804983, AI921753, AI640379, AI445025, AI439087, AW149869, AI568855, AI271786, AI873731, AI634737, AI251434, AI282281, AI802542, AI269696, AI439762, AI597918, AI812080, AI677796, AA449768, AI635942, AI573032, AI862144, AI289937, AI583065, AI687127, AI800185, AL037081, AI492528, AI873704, AW195957, AI630928, AI281773, AW087207, AI439089, AW029611, AI249257, AW274192, AI470648, AI436644, AI446606, AI872711, AI922365, AI687728, AL040243, AI924971, AI554821, AI800411, AI580984, AI480118, AI273142, AI610307, AI673256, AA470491, AW075667, AI283941, AI499285, AW170635, AI874261, AI569616, AI670002, AI926790, AI097248, AI446003, AI679724, AI609592, AI702073, AI932638, AI591420, AI680498, AI868831, AI282355, AI493567, AI281762, AI274508, AI801793, AI570781, AI538085, AI536638, AI274769, AW243820, AI446248, AW166970, AI648684, AW189268, AI624548, AI612913, AI866608, AW071417, AI539808, AI471712, AW167448, AW301409, AI812107, AI687465, AI826225, AW090071, AI634345, AI633125, AI681985, AI811353, AI811785, AW168802, AI859733, AI633419, AI475430, AI475371, AI918435, AW192652, AI445992, AI680138, AI636372, AI475134, AI433976, AI689248, AI357316, AI539771, AL036396, AI475817, AI174394, AI274745, AL045500, AW085799, AI247193, AI500077, AI801213, AI925156, AI697137, AI784252, AA835801, AI537244, AI499512, AI682971, AI520931, AI312542, AI340603, AI922665, AI818683, AI566670, AA833760, AI284020, Y11587, AF090900, I89947, AL117460, I48979, AL080060, AL049314, Y16645, AF113694, AF113690, AL133016, E07108, AF090934, AF104032, AL110221, AL117457, AL050108, S68736, AL133557, AF097996, AF090901, AL137527, U42766, AL110196, AL050146, AF090943, I48978, AL122050, AF125949, AF078844, AR011880, AF113013, AL133640, AF118070, A08916, AL050393, S78214, AB019565, AL050116, I89931, AL035458, AL122093, AL137550, AF090896, X84990, AL133075, L31396, AL050149, L31397, AF113689, AR059958, AF177401, AL133606, AF113676, AF118064, I33392, AF113691, A65341, AJ000937, AF113019, AF090903, AL096744, A08913, AL049938, U49908, AL117435, AL049452, X72889, AJ242859, |
|--|--|--|--|---|

| | | | | |
|---|---------|----|--------|---|
| | | | | AF017152, AL133565, AF106862, AF087943, AF113699, AF113677, X70685, U35846, Y11254, AL050277, AF146568, AL133113, A08910, AF158248, AL137557, E02349, E03348, AL122123, AL133560, AL080137, A08909, AL080124, AL049466, AL133080, AL137459, AL049382, AF026124, A93016, A77033, A77035, AF111851, AL049283, AL117394, A12297, A03736, AL122121, AF017437, AL133093, I49625, AL137271, AL122110, AF091084, AL137523, X63574, AL050024, AL137283, AL117583, E07361, A58524, A58523, AF125948, AF057300, AF057299, Z82022, AJ238278, X82434, U00763, AF079765, I00734, AL049430, AL110225, AL110280, AL049464, AL049300, AL133072, AL137521, S61953, AF067728, AL050138, I03321, AL122098, AF118094, U67958, AF183393, E00617, E00717, E00778, AL137648, AL137538, U80742, U78525, AL137463, U91329, AL117585, X96540, AF111849, U77594, AL050172, X65873, U58996, A08912, X93495, U72620, AL080159, AL110197, A93350, X98834, I42402, AL133014, E05822, AF111112, AL122049, I66342, AF026816, AJ012755, AF061943, AR013797, AL080127, AC004883, AL050155, Z37987, AF119337, AL080074, AL137560, AF106657, AF081197, E15569, I26207, AL133067, I09360, AF106827, AL133568, AF185576, E08263, E08264, X79812, AF008439, AR038969, AL133104, M30514, AF079763, AC006840, Y14314, AF095901, U96683, A07647, AF182215, Y09972, AR000496, U39656, AL133077, AF081195, AL035587, AL137533, AF153205, AL137480, AL137526, AL137479, AL117440, AL137556, AF100931, AL022147, E02221, AL137476, AR038854, AF003737, Z72491, AF134726, AL137294, AR029490, AF061573, E12747, AP000130, AP000208, L19437, AL133098, A90832, AF132676, AF061836, AL137488, AJ006417, AL137292, AL022723, AC007172, AP000247, AF162270, and N55769. |
| 4 | HDQHY04 | 45 | 904769 | AI871408, AA278445, AA479276, AI038602, N52018, AA706346, R80332, AI559411, AA305170, AI915377, AW440856, AI671553, AI809685, AA744961, AA309735, AA479371, AI940805, AI928450, N66171, AW103125, AA333461, AA857635, T80638, T80747, AA370795, AW088932, AA344676, AA279061, AA331630, R80226, AI421903, AI912434, AW149925, AI950892, AI887163, AI524179, AW089009, AI686576, AW073996, AI680221, AI620056, AI673278, AA731711, AL035458, AL050155, |

| | | | | |
|---|---------|----|---------|---|
| | | | | AL078630, AL117457, A12522, AF185614, AL122050, Z82206, X59813, A12297, A12558, AL049283, AR020905, AL049423, AL117394, AL050116, AF113677, X79812, I33392, Y13350, AR022283, AL117635, AF177401, AL049430, AC006112, AL049557, AF106657, AR034821, AF003737, A59344, AR029490, AL035587, Z13966, AF172400, and AF111849. |
| 4 | HDQHY04 | 46 | 852335 | AA333461, T80638, T80747, and N55769. |
| 5 | HWDH38 | 15 | 1028519 | AA410788, AW069227, AI278089, AW054995, AI733856, AI890324, AI923052, AA547979, AI634187, AI056177, AA847499, AA916430, AI457313, AA228778, AI732389, AI754653, AI537020, AA158190, AI679002, AW168433, AI891080, AL079734, AA515728, AA491423, AA832145, AI049630, AW068596, AA579179, AW004884, AW084445, AI923050, AI282479, AL120282, AW338035, AW338021, AL038842, AA719073, T09219, AW338179, AA669155, AA502991, AI279417, AA491955, AA236867, AA640710, AI254770, AA223174, AI821714, AI791913, AI284126, AI792133, AI821785, AI251034, AA176604, AW302315, AA569648, AA714110, AI133164, AI284543, AA513851, AW088846, AA557911, AA856866, AI921765, AI636627, AA290878, AI669589, AI360558, AW026305, AI752365, AI821768, AW265688, AI791185, AI421950, AI419337, AA181917, AW275432, AA668455, AI245693, R98218, AI499941, AI251284, AI251203, AA666332, AA568754, AI114557, AI361090, AW303098, AC006057, AL049776, AC002544, AL031311, AC002045, AP001051, AC004882, AP000046, AF129756, AC007021, AC007216, Z83844, AF038458, Z83845, AC007686, AL050318, AC004024, AL022323, AC002073, Z95115, AC003667, AC004531, AC005037, AL031666, AC002553, AF047825, AC005015, U95740, AL133163, AC004228, AC005049, AC002365, AP000356, AC005280, AL033518, AC005488, AC005972, AC006441, AL024498, AJ251973, U95742, AC006487, AC005746, AP000113, AP000045, AC003101, AC006511, AC005182, AC006064, AC004865, Z95114, AL022476, AC005668, AL031602, AC005695, AC008018, AC004098, AC004491, AC004921, AC005291, AC006530, AF001549, AC007263, AL049760, AC016830, AB023049, AF053356, AF024533, AP000557, AL080243, AC011718, AC007842, AC004821, AC005696, AC007450, AC003664, AC003098, AC004812, AC004020, AL133500, AC007376, AF111169, |

| | | | | |
|---|---------|----|---------|---|
| | | | | AF001548, AL009181, AC004590, AL035072, AC005519, AF139813, AP000116, AL049869, AL080242, Z83826, AC007371, AP000043, AP000111, AC007151, AL136295, AC016025, AC002549, AC007390, AL024474, AC007541, AL022238, AC006356, AL109798, Z98200, AC005295, AC008055, AL049636, AD000092, AC006101, AF031078, AL034549, AL021937, Z85996, AF111168, AC007790, AC002565, AP000692, AC005071, AC006079, AL035410, AF196972, AL035415, AC002395, AC004883, AC002400, AC006262, AC006236, U85195, Z95116, AC005874, AF134471, AC007066, AC004682, AC005514, AP000501, AL024509, AC002316, AC003665, Z81364, AL096791, AL034429, AL022334, AC003012, Z84469, U47924, AC005255, AF030876, AC007308, AL022322, AC006952, AL033521, AC004526, Z82206, AC005004, AC005747, AC004084, AC004149, AL049874, AL020993, AE000658, AC007312, AC007057, AL022237, AC005520, AC005823, AC005846, AC009509, AC005899, AC005921, AJ003147, AF207550, AC008012, AP000689, L47234, AP000010, AC003029, AP000036, AC006480, AL031985, AC004891, Z82217, AC005279, AL022329, AC004099, AC004671, AC002527, AC004983, AC003037, AL049569, AC005940, AC006948, AC005412, AC005011, AC004765, AC007227, AC005358, AC007917, AJ011930, Z99128, AC006211, AL022153, AC003110, AJ229041, U91326, AC003104, AB020868, AC008101, AP000359, AC007193, AL021918, AC006581, AC002039, AL049712, AC004967, AC006023, AC000003, AC004841, AC005786, AL008730, AP000193, AC006468, AC006449, AL133246, AC005243, AC005251, AC005288, AL096677, AC007191, AL121825, AC005102, AL109628, AF196970, Z93244, AF109907, AC003109, AL078634, AC004984, AL035443, AB001523, U82828, AC002044, AL034420, U63721, AC005722, AL117337, U95743, AC005225, and AC007314. |
| 5 | HWD4H38 | 47 | 889281 | AW392670, AL119443, AW363220, AW384394, U46341, AL119497, AW372827, AL119319, AL119335, AL119457, AL119324, AL119341, AL119483, Z99396, U46347, AL119484, AL119363, AL119391, AL119355, U46350, U46351, U46349, AL119444, AL119396, U46346, AL119522, AL134518, AL119439, and AL119418. |
| 6 | HHESQ62 | 16 | 1032630 | AA584489, AI133164, AI445582, AA838140, AW021583, AL046409, AI753365, AL120343, |

| | | | | |
|--|--|--|--|--|
| | | | | <p> AW021917, AW271904, AL039187, AI284640, AI733856, AW157005, AW341903, AA527961, F36273, AW247819, AA657835, F24728, AI281881, AI431303, AW276827, AI040051, AL038705, AI754037, AI028510, AI434695, AI017251, AW088846, AA992126, AI345654, AW270619, AW193265, AW275719, AA469451, AA582554, AI821044, AI805363, AW438643, H88666, AL038842, AA572685, AA516047, AA579566, AI963720, AI350211, AI923052, AI564185, AA634837, AI817230, AA955031, AI613280, AI370878, AI755214, AI754567, AW276935, AA613627, AI064864, AW270270, AI754653, AL040921, AA579179, R92640, AA587604, AI581486, AW276435, AW103509, AI754105, AA488746, AI623423, AA805841, AA503298, AI081147, AI888518, AA340747, AW265009, AL040054, AW338086, AA865262, AI446464, AW103383, AA720732, AI343143, AI375710, AA486559, AI537955, AI192631, H79308, AI085719, AA702605, AA664604, AI285696, AI696962, AA878140, AI360514, AA720702, AA502155, AL079734, AW167154, AI619997, AI075876, AI276298, AA085587, AI469599, AI537506, AL046205, AW277171, AI536625, AI270117, AI598003, AA470969, AW328331, AI753488, AI754955, AI471691, AI367523, AI927094, AI051037, AI821881, AA121777, AI821918, AI859280, AA569190, AI338426, AI471481, AA527209, AW081194, AI801591, AA236867, AW383952, AI110770, AA728937, AL037683, AA586458, AC004921, Z97054, AF001549, Z97989, AL049647, AL096791, D83253, AC007685, AC006080, AC008064, AC006367, AL049837, AL033525, AC012085, AC003036, AC002492, AL022163, AC003684, AL008720, AL121580, AL035588, AC005375, AL035443, AC006060, AC005244, AL008715, AF093117, AC005799, AC007384, AL096703, AL121578, AP000035, AL135783, Z95114, AC003048, AC002430, AC006343, AC004552, AL049610, AC006427, AC005081, AC006454, AL031291, AP000120, AP000320, AC007360, AC004695, AC009731, AL080274, AC006582, AP000167, AP000052, AC004865, AP000009, AC004453, AF124731, AF019664, AC004615, AC006213, AL031311, AC005539, AC002462, AC007372, AC005378, AC012384, AL121657, AC003035, AP000086, AP000517, AP000014, AC002365, AC005399, AC007955, AC006063, AC004659, AC005296, AL133289, AC007221, </p> |
|--|--|--|--|--|

| | | | | |
|---|---------|----|--------|--|
| | | | | AL035698, AC007388, AC004901, AC003685, AP000097, AC007344, AL132987, AL078638, AL133238, AC007298, AP000353, AC005325, AL049832, AC007228, AC005589, AC003046, AL033397, AB023054, Z82194, AC004067, AL031983, AL049761, AL020993, AL049633, AL022328, AC003954, AC000353, AL009031, AP000280, AC005200, AC007114, AL049829, AL109628, AP000134, AP000212, AC006012, AC008125, AC009263, AC007731, AL023775, AC005216, AC005728, AL137191, AL031010, AP000107, AC006211, AL022396, AC004486, AL049839, AL079342, AC006222, AL035608, AC008372, AL030996, AL033376, AC006305, AF200465, AC005500, AL117356, AP000245, AC006137, AL023574, D86992, AC004616, AL021391, AP000207, Z94801, AC005187, AC005233, AL121653, AF205588, AL008718, AC007091, AF165142, U62293, AC005751, AL031664, AL034424, AC005154, AL031848, L78810, AC004976, AC008273, AC004554, AL031904, AL035684, AC008055, AP000473, AC005695, AC004895, AC004805, AC004832, AC002287, AJ229043, AC006115, AC007058, AP000129, AF172277, AC002549, AC006026, AC006538, AJ229042, AC005552, AC005155, AP000967, D88270, AC005226, AC002525, AC004804, AF051976, AC009516, AC003013, AL050302, AL049643, AC007784, AC005291, AC004656, AL049833, AL022322, AC022517, AP000704, Y10196, AP000553, AC005962, AC006050, X87344, AC001228, U95740, AC003664, AC005192, AC005284, AC005288, AF196969, AL031905, AC004887, AC004914, AC007363, AC008033, AL049911, AC004837, AC005034, AC003682, AC005899, AC002123, AL049646, AP000696, AJ011930, AC005874, AF134471, AC007043, AC004946, AC005158, AL031774, AC004967, AL121998, AL022323, AL117337, AP000151, AC010200, AL034384, Z49258, AJ006345, AL035400, AC007707, AC005755, AC004913, Z92542, AC002080, AC006390, AC005736, and AC006195. |
| 6 | HHESQ62 | 51 | 839399 | AI445582, AL038705, AA527961, AA584489, AA838140, AW247819, AI733856, AW021917, AI754037, AW271904, AW341903, AW021583, AW157005, AL039187, AW276827, AA657835, AI133164, AA483223, AI434695, AI017251, AA598545, AI753365, AA582554, AA493708, AI040051, AW270619, AL040921, AI923052, AA634837, AI028510, |

| | | | | |
|--|--|--|--|---|
| | | | | AI276298, AI754653, AL120343, AA623002, AW088846, AI284640, AA854632, AA955031, AW275719, AA503298, AA470969, F24728, AA992126, AA121777, AI755214, AI754567, AW193265, AA133928, AA665021, AA507282, AL038842, AA469451, H79308, AA347114, AI079154, AL046409, AI431303, AL079734, F36273, AA490183, AI754105, AA577906, AW276935, AA584145, AA643770, AA586458, AA515905, AI581486, AA503019, AI805363, AA527209, AW438643, AW103383, AA720732, AW103509, AA516047, AI598003, AA565426, AI224619, AI343143, AA640685, AA846876, AA410788, AW270270, AA579566, R92640, AW276435, AW117740, AA878140, AA613627, AI564185, AW167154, AI350211, AI821881, AA488746, AI821918, AA865262, AA579179, AA502155, AA486559, AW162442, AA632648, AI821044, AI963720, AI623423, AI613280, AI281881, AA828042, AI469599, AW238278, AA587604, AI081147, AI753488, AI270117, AI859438, AA857486, AA805841, AL040054, AA503475, AA669155, AI537506, AA806796, AA664604, AI754955, AA533036, AA599920, AI338426, AI471691, AA721645, AA452887, AI696962, AW270771, AW265009, AW081194, AI051037, AA862243, AA569190, AA714282, AW328331, AI345654, AC004921, Z97989, AF001549, Z97054, AC012085, AC007685, AL096791, AL049837, AC008064, AL049647, AC003036, D83253, AC006367, AL033525, AL008720, AC002492, AL121580, AL008715, AL035588, AC005375, AL022163, AC009731, AL135783, AC006343, AL031291, AC006060, AF093117, AP000035, AL035443, AF124731, AC006427, AC004615, AC005244, AC002430, AC005799, AC007360, Z95114, AC007384, AL121578, AC006080, AC004865, AL096703, AC006454, AC003048, AL049761, AL121657, AL033397, AP000009, AL049610, AC004901, AC005325, AC004695, AL023775, AC003035, AP000120, AC004453, AL080274, AC007372, AC007221, AC006213, AC007344, AC002462, AC003685, AP000167, AP000052, AP000320, AP000097, AC005539, AC012384, AL133289, AC004067, AC002365, AC003046, AC004659, AC009516, AP000517, AC007298, AC004552, AL078638, AL031311, AC005399, AC008125, AL031010, Z82194, AL035698, AP000353, AL008718, AB023054, AL132987, AC003684, AC004486, AL049829, AF019664, |
|--|--|--|--|---|

| | | | | |
|---|---------|----|--------|--|
| | | | | AC005233, AC022517, AL023574, AL133238, AP000553, AC002287, AP000280, AC007955, AP000245, AC005552, AC005378, AL020993, AC006026, AC006137, AC005200, AC000353, AL049839, AC009263, AC004805, AL137191, AL049633, AL117356, AL031664, AL049832, AP000212, AP000134, AC005751, L78810, AL034424, AC007091, AC007114, AL009031, AC002115, AC006012, AL030996, AC004914, AF172277, AC007731, AC005500, AC006305, AC007228, AL079342, X87344, AC006115, AC004895, AC006211, AC005589, AL035608, AP000207, AC005296, AC004837, AL031904, AC007388, AC005284, AC005216, AL031848, AP000473, AC007058, AC006390, AC005192, AC002123, AC004967, AC005158, AL049833, AC005154, AC007784, AJ229042, Z94801, AC002523, AL121653, AC006088, AC002549, AC007707, AC005736, AP000014, AC006222, AP000107, AC005728, AC003013, AC008033, AF205588, AL031905, AL035684, AC004976, AC003093, AL096701, AC006568, AL049643, AL034371, AC005668, AC006538, AP000129, AC005187, AL049867, Y10196, AL049911, AC004554, AC008273, AF196969, AP000173, AC005082, AC006112, AF002223, AC008372, AL050318, AC004000, AC005553, AC008055, AL049874, AC004616, AC005226, AL121998, AL109628, AJ011930, AL035403, AC005089, AL031983, AF165142, AP000289, AP000086, AC002525, AC004832, AC005899, AC005695, AC005588, AL022323, AC004887, AC005800, AC003983, AC004508, AP000110, AP000042, AC007043, AC003682, AL035420, AC005962, AC007363, AL050302, AL049569, AC007630, AC006958, AC005057, AC003969, AC006288, AC005697, AC007637, AC005808, AL050317, AP000555, AC004033, AL049759, AF200465, AC001228, AP000403, and AC007327. |
| 7 | HDPSH53 | 52 | 882768 | AW392670, U46346, AL119457, AL119497, U46350, AL134528, AW372827, AL119391, Z99396, AL119335, AL119496, AW363220, AW384394, AL119363, AL119319, AL042965, AL119443, U46347, AL119439, U46349, AL119484, AL119444, U46351, AL134533, AL119324, AL134518, AL134524, AL119341, AL119355, AL119483, U46341, AL119399, AL119401, U46345, AL042433, AL119522, AL119396, AL134531, AL042614, AL042975, AL142132, AL134538, AL037205, |

| | | | | |
|---|---------|----|--------|---|
| | | | | AL134542, AL039857, AL042450, AL042984, AL042970, AL043029, AL042544, AL042542, AL043003, AL043019, AL042551, AL119464, AB026436, A81671, AR054110, AR066494, AR060234, and AR069079. |
| 9 | HLWCP78 | 19 | 941963 | AI018217, AI379675, AA131537, AA626816, AA954979, AI042383, AW016404, AA314725, AI539461, AI927251, AW131258, AW089331, AA339721, AI804427, AI783704, AA331134, AA335129, AI933420, N44789, AA316112, AA588075, AA131692, R58887, AA324444, AA777086, AA588662, R54764, R54952, AI383301, AW380879, AA335130, AA335009, AA335010, C02235, AI858465, AI632365, AI359586, AI961589, AI811845, AI565128, AI277008, AI472536, AI690813, AI963846, AI913312, AI554821, AI609069, AI500061, AA908294, AA848053, AW083750, AI114703, AW129106, AI633061, AI564426, AI537837, AI583578, AI571439, AI274745, AI633125, AI473528, AI568138, AI587156, AI955906, AI571699, AW082532, AI524179, AI538716, AI561356, AL041772, AI498067, AI886206, AW007555, AI288285, AI640370, AW167222, AI365256, AI553645, AI631216, AW081231, AW167146, AI524654, AI950729, AI249946, AW151785, AI476620, AI687362, AI689463, AW084425, AI886415, AI539690, AI699020, AI434693, AW169039, AI273856, AW081866, AI678446, AI625464, AI687568, AI476021, AI698403, AL134830, AI683940, AW082033, AI609409, AW161735, AI446785, AW243741, AI521799, AI680221, AI597731, AI624950, AI686808, AA761557, AW263569, AI635492, AW088560, AI273085, AI648454, AW151451, AW168452, AI619691, AL043345, N21402, AI439452, AW084117, AI655841, AI284060, AI917252, AW262042, AI583065, AA975588, AI569637, AI620003, AI702073, AI817552, AI623662, AW169671, AI814218, AI538850, AI669612, AI367364, AI433157, AI802542, AI613038, AI758437, AW118382, AI636507, AW169234, AI453229, AW198090, AI564719, AI499986, AA479803, AI570909, AI969641, AI245008, AI763414, AW172878, AW105330, AW075667, AA911767, AW166612, AI521005, AI690411, AW131288, AI919534, AI677796, AI521128, AL120706, AI951076, AW026905, AI815232, AI474146, AI284131, AW188382, AI582871, AI866901, AL138457, AI889189, AW083175, AW087199, AA835966, W48671, AI932794, AI479292, AI971587, AI927755, AI612913, AI624693, AI270183, AI590423, AW021717, AI873644, AI954945, AI538342, AI280637, |

| | | | | |
|--|--|--|--|---|
| | | | | AI625079, R10067, AW104724, AI624304, AI690946, AI679916, AI474107, AW104827, AW073865, AI497733, AI955943, AW075608, AI335208, AW084447, AI619502, AI261589, AI918677, AW263796, AW148408, AI862142, AI567243, AI637584, AI469112, AL120831, AW193530, AW073270, AI300294, AI610690, AI587114, AI469532, AI874243, AI865906, AI698391, AW157096, AW129597, AW089471, AI538564, AW087207, AW152182, AI247293, AW149925, AI591075, AI670009, AI361586, AI819976, AI301710, AI890907, AI874109, AI699865, AI648684, AW129722, AW169291, AL080140, AL117435, I33391, AR068466, Z13966, AL035407, AL137459, AF159615, AL133619, AF090934, AL122118, AL137548, AF113676, AF019767, AF118094, AF107847, AF115392, I89947, AF078844, E02152, AF126247, X76228, AL133640, A93350, AL080163, S83456, AF079765, AR013797, I26207, AL023657, AF090901, A08910, Z35309, A08909, AF132676, AF061836, AF074604, L13297, A08908, AR005011, AL137256, S61953, S63521, AL133070, L40363, AL133645, E12580, AF176651, AL117416, AR029490, AB019565, E02349, A08907, E12579, S77771, E00984, I04527, AL080126, AL110222, AL133558, E05822, E03671, AL137640, A08913, AF118090, I42402, AB007812, AL122045, U62807, AL110218, AR038854, U62317, AL133080, A77033, A77035, A08912, AJ006039, A08911, A18777, I89931, AL133104, A83556, AF205861, X06146, AL050092, I49625, AL049283, AL049382, AL137538, AF090886, AL137495, S76508, D16301, X98066, AL122049, AF113677, AF200416, U54559, AL122121, AL133624, S69510, I89934, AR029580, AL137530, Y11254, U67958, AL050277, AF102578, AF109155, I30339, I30334, E06788, E06790, E06789, AF002672, M79462, M19658, A12297, U49434, Y18680, I36502, AL137488, AR034821, AL137292, AL137558, AF008439, X99226, U88966, AF113691, AP000020, AL079293, I48978, AF089818, AF141289, A27171, AL133075, AF139986, A58524, A58523, AL137539, A21103, AF067790, Y10655, AL137556, A08456, A31057, AL049339, S65585, AL122104, AL137271, AL117460, I17544, I09499, I25048, I25049, AL049466, AJ004832, X83508, U75604, AR050959, I80064, X63410, AL137480, AL133067, AL137526, Y16645, AR000496, AR009628, AF067420, U39656, AL137281, AL117578, AF177401, AL137527, A03736, |
|--|--|--|--|---|

| | | | | |
|----|---------|----|--------|---|
| | | | | AF032666, AF000145, X57961, S78214, AF199027, AF030513, A30330, A30331, AR020905, AF113694, AL137429, AF059612, Y10823, A65340, E02253, AF155119, AF068615, X93495, A93914, AF146568, AL050116, AF058921, AL117394, AL034400, AL050393, I03321, A18788, A08916, I22272, AR016469, I41145, Y11435, AF100931, AR068751, AL096751, AF097996, X79812, AR053103, AF124435, AR068753, M85165, AB028451, U55017, X67688, AL117585, AC006501, X92070, AL137574, AF061795, AF151685, U49908, U37312, U83172, AL137275, U80742, AL137705, AL049426, U42766, U53505, AL080074, I89944, E12747, U87620, E03348, AF114170, AF113689, A86558, S36676, E03349, U83980, AL050172, Z82022, AF079763, AF017152, and J05032. |
| 9 | HLWCP78 | 55 | 901953 | AI018217, AA626816, AA314725, AA335129, AA331134, AA324444, AI383301, AW380879, AA335009, and R58887. |
| 10 | HTJML75 | 57 | 873355 | AW024974, AW303401, AI274799, AW074233, AI679074, AI499422, AI827575, AI928361, AA160606, AA315776, AI016059, AW249651, AW450819, AW079518, AI365272, AI338083, AA729126, AA831884, AI440443, AI283488, AI925468, AI274761, AW300875, AI983318, R42025, AA313959, AI240111, R20845, R37892, AI187826, R33244, R64261, R33245, AA862592, AA984343, AW087285, AI984649, AA580394, AI795901, H29720, R25583, AA283458, AA928568, AI796242, and AF191337. |
| 11 | HWBFES7 | 21 | 907063 | AF140342. |
| 11 | HWBFES7 | 58 | 907067 | AI419066, AI733552, AI554506, AI076219, AI400695, AI831940, AI248699, AI374872, AI968601, AI632124, AA761451, AW168136, AA811169, AI668630, AI637619, AI668632, H45142, H21585, AW058169, AA181557, H45443, AA182451, T88968, H45265, AA703073, T87799, T87800, AI686511, AW083175, AW090726, AI565128, AI745656, AI815018, AW129171, AW169604, AI274541, AI251906, AI819970, AI445131, AI669609, AW075413, AW118414, AW026087, AW090700, AW029081, AW167776, AW161735, AI818977, AI345261, AI345005, AI860370, AW188382, AI468959, AI345014, AI308790, AI609592, AW168718, AA903067, AI685798, AI859429, AW194441, AI349957, AI564515, AW081606, AW169149, AW170741, AW129918, AI653541, AI580927, AI554186, AW002838, AI924911, AI863133, AW130785, AI871923, AW151892, |

| | | | | |
|----|---------|----|---------|--|
| | | | | AI690490, AI702301, AI829327, AI635038, AI887247, AI608932, AI290154, AI744330, AL137481, AF140342, A70386, AR050959, AB026995, L12407, AF115410, AF148129, X60769, AF199509, U57352, Y14634, AL133054, E15324, AL079293, X73361, AL096709, AL133644, AL137558, AF091084, AF141315, D44497, D89079, AL137534, AL080140, U80919, AL133010, AL049452, E02914, AF161699, AF117959, AJ238278, AF137367, AR066486, A15345, X63410, M69222, A07588, S36676, Y11254, Y17607, AF082324, L31396, U68387, AF146568, A21625, AF124396, L31397, AL117693, AL080126, X79812, A23327, I08608, AF118558, D38178, U75932, AL079309, Y10655, S73498, AF113694, AF113677, X54971, U49908, X74593, AF058921, AL110159, S61953, AF188712, X53587, S70057, E16086, and AF102166. |
| 11 | HWBFE57 | 59 | 876136 | AC004947, AL078604, AL009028, AP000948, AE000659, AC007156, and AL022399. |
| 12 | HDPLO25 | 22 | 1032410 | AI096624, AI742426, AI343524, AW206464, AI743004, AW207506, AI971875, AI378547, AI763134, AA452699, AI306659, AI741119, AI808184, AW071446, AI681435, AI800948, AI923411, R55837, AA258683, AI912546, N41370, N24991, AW188250, AI686596, AI979273, AA450060, AI421707, AW365059, AW014260, AW136651, AA700541, AA977327, AA721031, AA485062, AI561015, AW139628, AW405442, AI863644, AI077467, AA001396, AI638834, AA361681, AW300126, AA485157, AW300113, R35711, R67953, AA974156, AI151115, R66863, AA976977, F32847, AI640210, AA017578, AA280510, F29704, Z44280, AW189156, AA569526, AI201767, R17530, AA329679, AI929037, AA888581, Z19983, C01202, AW080076, F08051, AI251434, AW167228, AL039086, AI932794, AI537261, AI802542, AI869367, AW022699, AI537677, AI269862, AW129106, AI499963, AW148356, AW071417, AL042628, AA833760, AW151714, AA225339, AI886181, AI624548, AI281772, AW198090, AI623682, AL110306, AL079963, AI929108, AI916419, AW087938, AI863321, AI270706, AW054931, AI571909, AL134712, AW079572, AI470651, AI310155, AW022682, AI932915, AW167918, AA427700, AI352497, AI620089, AI927755, AA641818, AI612759, AI886192, AW118518, AI932638, AW149994, AI783861, AL119863, AW051258, AI637748, AW117743, AW189270, AI288285, AI554821, AI886415, AW169604, AI862144, AI610362, AI343059, |

| | | | | |
|--|--|--|--|---|
| | | | | AI284131, AI889133, AI824576, AL038445, AI567582, AI538829, AI349933, AW105601, AL036638, AI280732, AI921281, AW008353, AI890214, AI633125, AI671663, AI570807, AI174394, AI343091, AL036396, AI923989, AI345416, AI345612, AL119791, AI800411, AI345253, AI366992, AW151136, AI698391, AW083573, AL037582, AL037602, AI637584, AI921379, AL043293, AW163834, AW020693, AI744185, AI345415, AI934035, AI569309, AI440239, AI500706, AI619502, AI871923, AI889189, AI500662, AI934011, AI473536, AI207510, AI073952, Z99428, AI673363, H89138, AI862135, AW167410, AI859464, AL043981, AW268302, AI677796, AI680162, AW072719, AI499285, AI306705, AW169653, AI866082, AI538716, AI923370, AI473799, AW073697, AI567351, AI587606, AI280661, AI758437, AI537617, AI494201, AI670009, AI537515, AI569583, AI433157, AL135025, AI702073, AL040241, AW026882, AW189415, AW090393, AI608936, AW020561, AI590686, AI468872, AL038605, AL036980, AI805688, AW129230, AI826225, AI811785, AI590120, AW079409, AI493576, AI249877, AI308032, AI685106, AL036631, AI886123, AW059713, AA806720, AI344785, AL041772, AI863240, AL036214, AL037454, AW160916, AI915291, AW152182, AI683492, AA908294, AI812015, AW117746, AL048644, AI921248, AL021578, AL137476, AL122098, AF008439, AF017437, AJ000937, AL133067, AL133075, AF125948, I89947, A08913, AL137292, AR038969, AF118094, Z82022, A77033, A77035, X93495, A08916, AJ012755, I48978, Y11254, AL110197, AL050116, A08910, AL137550, I89931, AL110196, A08909, I49625, L31397, I03321, I33392, A08912, AR038854, E02349, AF183393, AL133016, AL122093, AL122049, AF090934, AF100931, AL050024, AF113699, AF162270, AL137459, AL137533, AF087943, AL133557, Y14314, AF146568, AL122121, X72889, L19437, AL080159, AF111851, U58996, L31396, X82434, AL133014, AF090896, U35846, AL137560, A58524, A58523, AL049466, AF113019, AL133080, AL080060, AL122110, AL137556, I42402, AF026124, AF090900, AL050146, AL117440, AF067790, AF067728, L30117, U96683, AL050149, AL133093, I00734, AL050277, X65873, I48979, A65341, AL049283, AL117460, AL049464, Y09972, E00617, E00717, E00778, AF158248, AL050138, AL137521, AF061573, AF113694, E03348, AF113689, Y11587, AR000496, AL133640, U39656, E15569, AR059958, AF090903, |
|--|--|--|--|---|

| | | | | |
|----|---------|----|--------|--|
| | | | | <p>S68736, U78525, AL117432, AL133565, E02221, AL137557, AL122050, AF132676, AF061836, AL117585, AF185576, U80742, AL133113, AL137527, AF104032, A93016, AL049300, AF090943, Z37987, AF113013, AL117457, U68387, AL080137, AJ242859, AF111849, AJ238278, X84990, AL133077, AF090901, AL050393, AL133560, AF026816, A03736, AL137463, AF111112, I26207, AF091084, AL049452, X87582, E05822, AF118070, E04233, U67958, AL049430, AL137271, AF153205, AL137538, U49908, A93350, AF061795, AF151685, AF113676, A45787, AF177401, AL080124, AF057300, AF057299, AL110280, S61953, U00763, Y07905, AF119337, AL137526, AF118064, I09360, AL137478, AL080074, AL133098, AF017152, AL080127, AL096744, AL133568, AF106862, U72620, AL049382, E07108, AF113691, A07647, AL117394, X63574, AJ006417, AF126247, AF003737, A90832, AL110221, I09499, AF030513, AB019565, AF113677, AL117583, S78214, A12297, U91329, X96540, AF061943, AL050366, AF078844, AF113690, Y16645, AF097996, Z72491, AL050108, AF061981, AL133072, U42766, AF079765, AR011880, AL133104, AL117435, X70685, AL049314, M30514, AL137648, AF079763, E08631, AL023657, AF125949, AL110225, AL122118, AL133606, X98834, AL122123, E08263, E08264, E07361, AC002467, AL050172, AL049938, AF139986, AL137294, AL080086, AF210052, AL137273, AF081197, X53587, AR020905, X92070, AL080148, E12747, AL050092, AL137480, and X81464.</p> |
| 12 | HDPLO25 | 60 | 874064 | <p>AI096624, AI742426, AW206464, AI743004, AI378547, AW207506, AI343524, AI763134, AW071446, AI923411, AI912546, AI971875, AW188250, AA452699, AA258683, AI979273, AI421707, AI306659, AI681435, AI741119, AA977327, AA700541, AW136651, AI808184, AA721031, AI561015, AI800948, N24991, AA485062, R55837, AA450060, AW139628, AI077467, AI863644, AI686596, AW365059, AW014260, N41370, AW300126, AI638834, AW300113, AI151115, AA974156, AA976977, F32847, AA001396, AW405442, AA361681, AA485157, R67953, R35711, F29704, AI640210, AW189156, R66863, AA017578, AI913608, AA280510, AA569526, AI201767, Z44280, AA329679, R17530, AI929037, AA888581, Z19983, F08051, C01202, N77044, AL021578, and AF131789.</p> |
| 12 | HDPLO25 | 61 | 902188 | <p>T29199, AA362689, AA362688, AA362690, AL044652, AA725783, AW303008,</p> |

| | | | | |
|----|---------|----|--------|--|
| | | | | AA826358, AI049508, AI886654, AI055937, AI635819, AI088768, AA653154, AI630176, AA774949, AA349927, AA569471, AA099939, AI174766, AI793028, AL135262, W55874, AA662225, AW265244, AW338201, AI086530, AA484145, AA527602, T60166, AW265186, AI918465, AA601356, AL038705, AA702729, AA719407, AL037383, H96467, H96480, AW243793, AL042753, AI217937, AA292178, T06754, AL038799, AA309844, R17793, T94249, A49045, I66487, A94048, A94061, AR035224, I09252, I09251, I07209, I07249, I09270, I09268, I09269, AR068508, AR068510, AR067734, AR068509, A63954, A49701, AR067731, A29109, A32111, I66495, AR068550, A23373, AR068551, I66494, AR067732, E01324, I08638, A94046, A94054, I58322, I58323, I66497, I66496, I66486, A27169, A27170, A39929, A83151, I09267, I66498, AR038321, AR038307, I91969, AR051537, AR051524, AR051528, AR051547, Z95152, AC005920, AC005694, AL049712, AC005529, Y18000, AC002470, AC000052, AB001523, AC005913, AC002563, Z94044, AL035562, AC005527, AC004087, Z83845, AC002365, AC004019, AF053356, AC006961, U95743, L10641, AP001054, AL078583, AF217403, AC007191, AC005695, AL008637, AC006035, AL022313, A42964, AC004525, Y10196, AL049874, U91318, AL049569, AL049690, AL031768, AC006530, AC004812, AC007308, U66059, AC009516, AC002112, Z86090, Z94277, AC005004, AL109758, Z97054, AC004149, AC004879, AC002990, U95740, AL023574, AF001549, and AL035469. |
| 13 | HDPWN93 | 23 | 992925 | AA432202, AI890824, AI753494, AI651671, AA993211, AW001898, AL039524, AI800905, AI246773, AI682295, AI658613, AI631136, AW189302, AI372827, AI050708, AA531521, AI346388, AI683842, AW296359, AW372955, AI685246, AI589722, AW271749, AA622365, AI352313, AI088281, AI372826, AI015389, AI539826, AA349564, AA448189, AI760986, H15544, H15545, AA349563, H09846, AI991731, AA320029, AA383782, H15604, W67789, H15603, AI015277, R33930, AA543091, AI469944, R48594, H09761, AA830547, AA505499, AA320560, AA326027, R79459, AA322654, L32015, C20992, AI828309, AW117647, AW190887, AW262975, AA317254, AA143736, AW103622, AW050451, AI609346, N21451, AI905534, AA282625, AW007401, AA284991, AA621245, |

| | | | | |
|----|---------|----|--------|---|
| | | | | AA143707, AI811818, AA393671, AI678229, AA429592, AW407359, AC004590, AF086245, AP000161, AP000020, and AL137367. |
| 13 | HDPWN93 | 62 | 887914 | AA282625, AW407359, and AC004590. |
| 13 | HDPWN93 | 63 | 905983 | AA432202, AI890824, AI753494, AI651671, AA993211, AW001898, AL039524, AI800905, AI246773, AI682295, AI658613, AI631136, AW189302, AI372827, AI050708, AA531521, AI346388, AI683842, AW296359, AW372955, AI685246, AI589722, AW271749, AA622365, AI352313, AI088281, AI372826, AI015389, AI539826, AA349564, AA448189, AI760986, H15544, H15545, AA349563, H09846, AI991731, AA320029, AA383782, H15604, W67789, H15603, AI015277, R33930, AA543091, AI469944, R48594, H09761, AA830547, AA505499, AA320560, AA326027, R79459, AA322654, L32015, C20992, AI828309, AW117647, AW190887, AW262975, AA317254, AA143736, AW103622, AW050451, AI609346, N21451, AI905534, AA282625, AW007401, AA284991, AA621245, AA143707, AI811818, AA393671, AI678229, AA429592, AW407359, AC004590, AF086245, AP000161, AP000020, and AL137367. |
| 14 | HWHPM16 | 24 | 903814 | AI961281, W25575, W73855, AA025948, W69100, W95776, W92535, W69380, W68286, AA359882, AA022503, AI131566, AA846828, AA706316, AA777022, AA480817, W95987, W73821, W32894, AI141167, C00662, W95733, AI161236, AW001367, W58747, Z99396, AI148339, AA854719, AW009909, W94659, AI092860, AW392670, AL134902, AI142132, AL134525, AW372827, AL119355, AL119484, AL119457, AI150077, AL119319, U46349, AL119324, AL119483, AW363220, AW384394, AL119391, AL119439, AL119497, U46351, AL042978, AL119363, U46350, U46347, AL042614, AL119443, U46341, AL119444, AL119522, AL119341, AL037205, AL119401, AL119396, AL043003, U46346, AL134528, AL119335, AI144221, AW009219, AL043033, AL043011, AL134538, AL042544, AI142137, AL119418, AL119399, AL042984, AL119496, AL134533, AL134536, U46345, AL043019, AL043029, AL043039, AL119464, AL042450, AL042542, AL042965, AL042975, AL042551, AD001502, AF086315, I25027, AR054109, I44515, I26928, I26930, I26927, I25041, I44516, AR009152, AR019098, AR035224, I85513, AR009151, AR027099, A94046, A94054, I05393, A10617, AR028792, A01324, |

| | | | | |
|----|---------|----|--------|---|
| | | | | A01323, AR019094, AR034783, I63120, A92666, A92668, I49890, A92667, AR064322, AR064323, AR064321, A94048, A94061, A92665, AR038321, AR038307, AR020199, AR020200, AR001287, A49045, AR020198, AR020197, AR067734, AR067731, A32110, AR067732, AR067733, AR064320, A83642, A83643, AR029418, I89986, AR029417, A70359, I05430, AR028791, AR028793, I09121, AB026436, AR055065, A30600, AR055066, AR066494, AR060234, AR054110, A81671, and AR069079. |
| 14 | HWHPM16 | 64 | 762226 | AI554127, AL049113, AI623461, N27660, AA714202, AW027142, AI146746, AA234520, N62741, AW149632, AA024451, AA971630, R49567, AI278865, AI559459, AA526772, AA333352, AA663249, AI133119, AA547956, AA363341, R42517, AA548156, AA548037, AA347611, T55030, AA372715, AA804413, AI610110, AA206665, AA642366, AA359112, AL035410, AL030998, AC003093, AC007319, AC004707, AL022147, Z83820, AC003983, AF196971, AC004691, AC005406, AC010209, AC003678, AC004584, AC003689, U77841, AC011604, AC005908, Z97054, AP000277, AP000037, AP000105, AR036572, U91328, AL031073, M10976, AL118518, AP000473, AL023280, AP000083, AC007206, AC007073, AC005304, AL049713, AL009174, AF064859, AL034343, AC006145, AL049543, AC007204, Z84718, AP000351, AC005871, Z97180, AP000305, AC006205, AP000047, AC005599, M74509, AP000115, AC004706, Z99572, AC002310, AF196779, AP000128, AP000206, AC002094, AC004985, AC005181, AP000245, AL034376, AC007324, AC008101, AC006484, AC008079, Z75896, AL109756, AC000357, AJ249921, AL096677, AC004455, AC005775, AL021395, AL121578, Z97195, AC004813, AF050154, AF187320, D00591, AC005538, AC004998, AF002993, AC003080, Z69722, AC004386, AC005023, AC002418, L47334, AL035555, AF026257, AC004953, AP000705, AL021391, D90086, AL021940, AC004131, S74906, AF118808, AL078624, AL136501, AC006481, AC012039, AC006198, AL137711, D10450, AC005326, and Z56880. |
| 16 | HDPXY01 | 26 | 879048 | AW176604, AJ230819, AL133649, and D29801. |
| 16 | HDPXY01 | 67 | 904768 | AW131590, AI926671, AI624771, AI978673, AI681035, AI570135, AW027636, AA883228, AW297661, AI434191, AW270077, AI745206, AA471014, AI810158, AI377790, AI983421, AW167619, AI870703, AA703646, AA972392, AA147535, AW117883, |

| | | | | |
|----|---------|----|--------|---|
| | | | | AI432478, AA703546, T49102, AI355795, AI623870, AA233934, AA234204, AA309040, AJ230819, AI243001, AA293599, AW176604, AA147657, AW193857, AA857348, AI480215, AW205948, AL133649, D29801, and I22312. |
| 16 | HDPXY01 | 68 | 895716 | AI681035, AI983421, AI377790, AW270077, AW027636, AW297661, AI480215, AW117883, AI434191, AI432478, AI978673, AW131590, AI624771, AA972392, AI926671, AA471014, AI355795, AI745206, AW167619, AI570135, AA703646, AA147535, AI623870, AA883228, AA233934, AI810158, AW205948, AW188390, AA703546, AI243001, AA234204, AW193857, AI870703, AA857348, AA678514, I22312, and AL133649. |
| 16 | HDPXY01 | 69 | 895715 | AW131590, AI926671, AI624771, AI978673, AI570135, AA883228, AI434191, AI745206, AW297661, AI681035, AA471014, AW027636, AI810158, AW270077, AI983421, AW167619, AA703646, AI870703, AA972392, AI377790, AA147535, AW117883, AI432478, AA703546, T49102, AI355795, AI623870, AA233934, AA234204, AA309040, AI243001, AA293599, AW193857, AA857348, AW205948, AI480215, AL133649, I22312, and AA147657. |
| 17 | HWBEM18 | 27 | 949402 | AA502912, AI050870, AW006390, AL119604, AW401673, AI440347, AA310795, AA524413, AA524099, AI688921, AW167025, AI970597, AI028774, AW188383, AW150222, AW083366, AA394112, AI802599, AA515127, AA420978, AA632926, AW081261, R82177, N55034, AI814038, AW407844, AA504752, AA446015, AA652891, AI000287, R82219, F06324, AA478999, AI191730, AA456551, AW406901, AA577322, R11808, AA504654, AA661852, AA995856, AA321769, AI672210, AW270191, AA357086, AA477935, AA292315, AL039259, AW403756, AA470467, H62806, W76432, AA278264, H91643, T99718, AA347047, AA902263, AA278730, W04875, AI953093, AI200728, AA687290, AI018458, AA486391, W72039, AI675899, AI870318, AI923553, AW027693, AF113751, Y00826, AB020713, AL117527, and X87546. |
| 17 | HWBEM18 | 70 | 906580 | AA995856, AI672210, AW270191, AL119604, T99718, Y00826, AF113751, AB020713, AA255512, AA282876, AA283032, AA420593, AA483157, AA483242, AA491774, AA491825, AA491980, AA493272, AA493718, AA502863, AA505839, AA507521, |

| | | | | |
|--|--|--|--|---|
| | | | | AA515147, AA515158, AA547997, AA548059, AA548259, AA548549, AA552753, AA552844, AA554985, AA557741, AA563828, AA564736, AA564831, AA565136, AA565516, H56134, H78512, H92599, H95100, H96481, N22409, AA581666, AA581901, AA581910, AA582052, AA584907, AA583392, AA583603, AA584459, AA584534, AA584627, AA584743, AA584747, AA584834, AA586654, AA586665, AA587520, AA593542, AA594081, AA594147, AA594706, AA595827, AA601406, AA601412, AA601418, AA601495, AA601625, AA601628, AA603182, AA604932, AA613153, AA613193, AA618000, AA618459, AA631445, AA572730, AA577765, AA577773, AA577803, AA577825, AA577875, AA577884, AA577921, AA577931, AA578885, AA661560, AA664727, AA665322, AA713821, AA713895, AA714107, AA714486, AA715716, AA715807, AA720777, AA720805, AA720943, AA721030, AA721049, AA730122, AA736954, AA737245, AA736489, AA742398, AA745295, AA749350, AA761416, AA767964, AA769182, AA804967, AA804973, AA808402, AA824585, AA829583, AA834071, AA834194, AA836427, AA836838, AA856868, AA872859, AA873328, AA911390, AA935394, AA937357, AA971198, AA976274, AA983272, AA983440, AA989133, N29555, N75439, N84325, N85750, W37681, W90097, W93753, N86042, N89024, N89311, AA017128, AA018677, AA082639, AA084139, AA088273, AA091426, AA093759, AA094022, AA095145, AA096335, AA121766, AA121767, AA121839, AA121840, AA121876, AA132944, AA136395, AA136629, AA136943, AA136977, AA157151, AA159480, AA160931, AA165176, AA164621, AA167809, AA167240, AA167241, AA167453, AA167491, AA167492, AA167468, AA167543, AA167544, AA167579, AA169847, AA169141, AA169228, AA169233, AA169557, AA169344, AA169346, AA169351, AA176467, AA176906, AA179044, AA179174, AA179365, AA179963, AA179985, AA180281, AA180454, AA181443, AA188613, AA188692, AA188716, AA188953, AA188838, AA191062, |
|--|--|--|--|---|

| | | | | |
|--|--|--|--|---|
| | | | | AA191204, AA194957, AA195197, AA206780, AA206908, AA643815, AA649905, AA651698, AA211180, AA211914, AA214036, AA218670, AA218754, AA218852, AA218913, AA218880, AA219034, AA219167, AA219168, AA219209, AA219222, AA219479, AA077547, AA227287, AA227893, AA249258, AA252123, AA257010, AA399244, AA421028, AA437400, AA448567, AA451922, AA458671, AA458672, AA458716, AA458783, AA464904, AA464929, AA464930, AA485341, AA485874, AA487777, AA487851, AA487953, AA487954, AA486097, AA486160, AA486768, AA486808, AA487304, AA487519, AA491346, AA489850, AA489899, AA489968, AA489989, AA490015, AA490032, AA490150, AA490183, AA490246, AA496279, AA504660, AA504676, AA504875, AA504927, AA505040, C75155, AA598981, AA634823, AA662979, AA662985, AA663003, AA663004, AA663029, AA663091, AA663223, AA663367, AA663500, AA668151, AA668186, AA668235, AA668252, AA668281, AA668292, AA668327, AA668372, AA668412, AA668496, AA668699, AA680277, AA456204, AA456920, AA457089, AA457176, AA457219, AA457283, AA457322, AA457507, AA457548, AA457599, AA457601, AA434080, AA434354, AA679143, AA679325, AA708186, AA716403, AA719211, AA722310, AA683065, AA683238, AA758129, AA774908, AA775210, AA775329, AA776006, AA778726, AA778953, AA778965, AA780571, AA780730, AA780751, AA782269, AA782279, AA782321, AA833652, AA773127, T03057, T03259, AA984451, AA984452, AI026095, T39228, T40391, T40631, T41239, T48646, T48647, T49954, T49955, T55332, T56669, T57324, T59577, T59685, T59821, T59968, T60702, T63021, T63167, T63720, T63875, F06978, T65750, T65825, F01046, T89256, T92131, T93559, T94346, T94701, T94872, T94919, T96260, T99749, R05946, R06051, R14500, T68944, AA772877, AA773954, AA774057, AA774071, AA694367, AI246137, AI270057, AI298238, AI356307, AI370186, AI453652, AI475824, AI468889, AI565003, AI623435, AI627881, AI174659, AI185790, AI198511, AI201329, AI204938, AI654729, AI219575, |
|--|--|--|--|---|

| | | | | |
|----|---------|----|---------|---|
| | | | | AI242491, AI267152, AI267267, AI267283, AI267387, AI267399, AI267434, and AI267690. |
| 17 | HWBEM18 | 71 | 877573 | AA502912, AI050870, AW006390, AI953093, AI675899, AI870318, AI923553, AW401673, AA310795, AI440347, AW027693, AA524413, AI075145, AI359954, AA524099, AI597640, AI083744, W72039, AA456551, AI688921, AW167025, AI970597, AI028774, AW083366, AA702590, AW008239, AW188383, AW150222, W04875, AA133158, AI075281, AI380071, AI696180, AI919128, AI458590, AI690287, AW341136, AI290772, AA394112, AI564066, AI802599, N50705, AI912716, AI290867, AA515127, AI697475, AI459065, AI218744, AI214605, AI697999, AA632926, AI630774, AI268530, D52783, AI692351, AI373390, AI304923, AA420978, AI342694, AI348602, AI760118, AW081261, N70505, AI342675, AI348362, AI302027, AI218972, AI340052, AI702729, AI342752, AA459322, AA923137, AI300921, AI818912, AW166144, AA971517, AI458565, AW341160, AI214348, AI348572, AI867102, AI032703, AI761002, AA436286, AI369001, R82177, N55034, AI807719, AI814038, AW407844, AA504752, H62722, AA446015, AA909765, R82219, AI000287, AA652891, AA829731, F06324, AA770635, T33986, Z39227, AA478999, AI702141, W76432, AI268398, H91642, AA987310, AI191730, AI268346, AA459103, AA808738, AA995397, H91643, R11808, AA931687, AA996234, D81323, AA988933, AI215821, AI216187, AA977023, AA577322, AA976720, AA916695, AA436160, AA504654, AA661852, AI197990, AA938371, R43824, AA973530, D53592, AI002429, AA321769, AA279841, AA524872, AA885803, D80894, AA512952, AA789129, AA971520, AI860335, AA357086, AA477935, H62806, AA923086, AA292315, AA974038, AA618436, AA931944, AA844341, AI300623, AA989091, AL039259, AA939247, AA470467, AA250870, R37724, T35082, AW059597, AI023554, AA347047, AA916526, AA902263, R28861, AA278730, AA687290, AI826758, AA486391, AA937729, AW406901, AI002329, AB020713, AL117527, Y00826, AA968738, AI291215, AI299477, AI350166, and AI299545. |
| 18 | HDTFE17 | 28 | 1043391 | W35383, AA015734, AA015831, AA018179, AA020824, AA020847, AA021564, AA021617, AA088190, AA088363, AA261917, AA262421, AA418866, AA426526, AA430530, AA613360, AA573487, AA577277, AA761384, |

| | | | | |
|----|---------|----|--------|---|
| | | | | AA831386, AA864866, AA902924, AA922543, AA934011, AI052766, AA481843, AA481990, AA433981, AA844171, AI004425, AI033046, AI073848, AI087207, AI088106, AI093860, Z38440, Z42185, AA683532, AI272861, AI285507, AI301768, AI343925, AI351185, AI356409, AI359625, AI479005, AI569081, AI582503, AI147150, AI623660, AI184904, AI193793, AI264308, AI273503, AI587448, AI351022, AI632302, AI621103, AI670930, AI698414, AI701275, AI801704, AI868864, AI888089, AI916106, AI949047, AI963403, AI970843, AI989788, AW073000, AW132028, AW196624, AW207687, AW193329, AW511737, AW590114, and AW590183. |
| 18 | HDTFE17 | 72 | 874477 | AW374334, T08905, AI652677, and AF196972. |
| 18 | HDTFE17 | 73 | 892317 | AA621888, AA621871, T56553, AW087213, AI673648, AW246689, AI655004, AW250829, AW170222, AI140471, R94667, AA149673, AW183216, AW189831, AA612723, AA149754, AW374334, AI815239, AI440263, AI872423, AI859991, AI371265, AI539771, AI537677, AI494201, AI500659, AI815232, AI801325, AI500523, AI582932, AI923989, AI284517, AI500706, AI445237, AI491776, AW151138, AI521560, AI889189, AI500662, AI866786, AW172723, AI284509, AI889168, AI866573, AI633493, AI434256, AI805769, AI888661, AI284513, AI888118, AI436429, AI889147, AI371228, AI440252, AL047422, AI860003, AI610557, AI887499, AI963849, AW409775, AI798359, AI866465, AW151136, AI538850, AI890907, AI539800, AW050725, AI866469, AI434242, AI355779, AI581033, AI242736, AI539781, AI371251, AI866510, AI440260, AI559957, AI521571, AI923046, AI554821, AI433157, AI345649, AI758988, AI583578, F26535, AW151132, AI538885, AI804505, AI356470, AI866691, AW191003, AI539707, AI887775, AI590043, AI926593, AI289791, AI285417, AW151974, AI623302, AI582912, AI927233, AW058275, AL042750, AW131994, AI500714, AI285439, AI358271, AI623736, AW194509, AW404239, AI805688, AI491710, AI431307, AI801286, AI440238, AI567971, AI431316, AI539260, AI828574, AI273179, AW151979, AI499104, AI431238, AL047152, AI702065, AL048375, F27428, AI885949, AW089557, AI285419, AI432666, AI469775, AI866581, AI340603, AL047398, AW074057, AI567953, AI815150, AI371442, AI446495, AI952433, AI867068, AI889191, AI309443, AI225248, AL038778, AI433976, AI559976, AI698352, AI282249, AL042377, |

| | | | | |
|----|---------|----|---------|--|
| | | | | AI371229, AI287476, AI954721, AL079910, AI366900, AW089732, AW083804, AI049851, AA593940, AL041918, AI343030, AA715307, AI275175, AL047763, AA954134, AI860027, AI499463, AI783569, F36033, AW024921, AL038575, AI432644, AI610362, AI648567, AI620284, AA760727, AI924051, AI440239, AI560514, AI349246, AI469516, AA642381, AI537273, AW265004, AI436456, AI963846, AI567940, AI817244, AI612913, AI567993, AI285826, AI863014, AI521594, AI499512, AI815233, AW090387, AI889133, AA641818, AI249946, AL048323, AL042787, AL042572, AI798571, AI610402, AI276331, AI434223, AI500458, AI610429, AL045500, AI628850, AI433968, AI312432, AI872315, AI539632, AI683821, AI889148, AF196972, M81784, AL009029, AF098162, AL022315, AF126372, AC005743, AC004093, AC004617, AL022723, X72889, AL132985, AC005773, I66342, U96074, AL133015, AF109906, AL133608, AL122101, AL080146, L40363, AF132676, AF061836, I48978, AC007458, AL110196, AL137459, Y13653, I22020, E12580, AF199027, E02221, AF177401, I80062, AF192557, I42402, and AR008521. |
| 19 | HBXAB02 | 29 | 1018820 | AW293312, AI937384, AI871895, AW084788, AI804641, AI357795, AA984106, AI184076, AI083793, AL118920, AI635836, AI808745, AA576156, N95023, AA242997, AA741070, AI174555, W04870, AA536061, AI702093, AI025407, AA412707, AA405096, T03377, AI806457, H23645, AA730162, H07961, AW386950, AI682515, AI242242, T33589, AA598585, AA291077, AA781341, AI991764, AA042883, AA029249, AA677827, AI991911, W51892, Z46068, H46700, T66367, AW005882, T15492, AW005939, AA350854, F12308, AA598784, Z40425, AW373072, F09934, AI866250, AI801018, F37442, AA378758, AA043009, AA861126, AW176443, N70500, AI380101, AI682087, F27243, AW337643, AI277155, AW386952, H07870, F33259, H23604, AA770644, AI587570, AA243080, AA676757, AW393033, AA326720, AW104785, W23868, AA284131, AA627281, AI583578, AA551491, AI890574, AW176422, AI439978, AI682971, AI587156, AI950729, AI570807, AW169234, AW104641, AI498067, AI932794, AW007555, AI702073, AI700159, AW082532, AI961589, AW198090, AL079963, AI637584, AW104724, AI433157, AI254727, AA514684, AW075381, AI270183, AI611738, AI633125, AI923370, AI434468, AI924701, AI286256, AW192652, AI584140, |

| | | | | |
|--|--|--|--|--|
| | | | | <p> AW080327, AI274655, AI591420, AI698391, AI889189, AI768496, AI282903, AI801793, AL119863, AI927233, AI056328, AI539800, AI570909, AI922689, AI926367, AI624693, AW074374, AI670009, AW081866, AI538259, AW103878, AI468872, AW026882, AW129230, AI818353, AW051059, AI590227, AI284084, AI581362, AI815232, AI445432, AI915291, AW089275, AI288285, AW080090, AI832245, AL045500, AW166583, AW148363, AI619748, AW169132, AI571867, AA908294, AI619691, AL121041, AW051258, AI281687, AI590686, AI439452, AI648454, AW163834, AL040390, AI500061, AI684234, AI524677, AI678688, AI864836, AI677796, AI934259, AW075667, AI267162, AI522052, AI933992, AW118518, AI859915, AI678446, AI567128, AI697137, AI609375, AW087199, AI582932, AI291973, AW148294, AI355827, AI684244, AI497733, AI613038, AI659334, AI963191, AI955943, AI493567, AI571439, AI862139, AI474146, AI802542, AI951076, AI471712, AL110306, AI609409, AI619502, AI889376, AW193530, AW132056, AI799470, AI866465, AI648509, AI572021, AI564719, AI287326, AI929108, AI440448, AI702068, AI478723, AI696398, AW080746, AI633062, AI554821, AA848053, AI423326, AI445611, AI469470, AW268964, AI288305, AW168452, AI933783, AI952145, AW020693, AL047100, AW081653, AW078895, AW152182, AI539780, AI612913, AI573060, AW088560, AI679550, AI270099, AI863241, AI866751, AW168677, AW131165, AW083573, AI627893, AI571909, AI640704, AB020649, AL137297, AL137459, A65340, AL080159, AI8788, I89947, AL050366, AF069506, AF159615, AR034821, AF215669, AL133619, I48978, I66342, AL080139, A77033, A77035, AF177401, AF017437, A08916, A08913, AL049283, AL133016, S36676, X82434, AF100931, Z37987, AL050149, AL133113, A08910, M92439, A08909, AL137526, AL117460, AF102578, AI8777, AL050138, X79812, I17544, I09499, U35846, AL049339, Y16645, I33392, AL137271, AB007812, AL133075, AF104032, AL110296, AF183393, AL122100, AF026816, E15582, AL117435, AF087943, U37359, I00734, AF091084, I89931, X70685, Z82022, Y14314, E00617, E00717, E00778, E02221, AL137539, I49625, AF044323, AL137548, A15345, AL122110, AL137550, AL133568, AL050092, AF032666, X72889, AF113013, AL136884, AF113677, AL050024, AR038854, AL137463, AF067728, Y11587, AL117626, AF115392, AF031147, </p> |
|--|--|--|--|--|

| | | | | |
|----|---------|----|--------|---|
| | | | | AL110225, AL110280, AF113019, AF028823, AL133558, I48979, AF118090, X63162, AF125949, AL137558, U42031, AF090901, Z97214, AL133560, E05822, AL122050, AF090900, A65341, AL122106, I03321, AF111849, S69510, AF139986, AL133010, X06146, AF008439, AF090903, AL133093, U72621, AF182215, E01573, E02319, AF118094, AF065135, AL133645, AL117583, AF176651, AJ238278, AL133557, AL110221, AL117457, A08912, S68736, AL080126, A03736, S61953, AL117648, AL050277, AF126247, AL137530, A76335, X80340, AF111851, U53505, AL137538, X84990, AL050116, AL137488, AF030513, AL137294, AF113691, AJ005690, AL137479, AF067790, AL133104, AF113690, AF090934, AL080074, S63521, A08908, AF141289, AL133565, AF079763, AL049452, Y09972, AL122045, A52563, L04504, S76508, AF061943, AR011880, A58524, I89934, A58523, AF180525, I26207, AL133640, E02349, AL117416, AL080140, AL133665, AL080148, AL110224, AF106862, AF036941, AB016226, AF106697, A57389, AF090943, AL049938, Y11254, AL049382, Z13966, AF192557, X52128, E01314, AL137529, M27260, AL137256, S77771, AL137658, A27171, S78214, X60786, U75932, Y10655, AF113694, S78453, AL137557, AL133080, L04849, AL049430, L40363, AF113699, AF132676, AF061836, U49434, AF004162, U80742, AL050155, AL050393, D83032, E02152, AF114170, AL137283, L31396, I17767, AL050146, AF057300, AF057299, AL133112, U91329, X53587, U88966, A86558, U87620, X57961, AL080137, AL137555, AL133606, L19437, AF097996, AL049347, AL049314, E06743, AR059958, AL122098, L31397, AF017152, AF113676, and A45787. |
| 19 | HBXAB02 | 74 | 890192 | AA984106, AL118920, H07961, AW386950, AA242997, Z46068, T66367, AA350854, F12308, AW373072, AA378758, W04870, AW176443, AW386952, AI380101, H23645, AA291077, AA326720, W23868, W51892, AW176422, T73545, AB020649, and AL137297. |
| 19 | HBXAB02 | 75 | 901261 | AW293312, AA707908, AA412707, AI937384, AI871895, AW084788, AA741070, AI804641, AI357795, AA536061, AI025407, AI808745, AI184076, AL118920, AI083793, AI635836, N95023, AA576156, AA242997, AI174555, W04870, AA405096, AA781341, AI702093, AA042883, T03377, H23645, AA730162, AI806457, T33589, AI242242, AA598585, AI380101, AI682515, AA291077, |

| | | | | |
|--|--|--|--|---|
| | | | | AA677827, AA029249, AI991764, AI991911, W51892, Z46068, H46700, AA984106, T66367, AW005882, AA861126, AA350854, AW005939, T15492, N70500, F12308, Z40425, F09934, AA598784, F37442, AI866250, H07870, AI801018, F27243, H23604, AW176443, AW337643, AI682087, AA043009, H07961, F33259, AI277155, AW386950, AI587570, AA243080, AA676757, AA627281, AW386952, AA326720, AW393033, AA770644, AW104785, W23868, AA284131, AI583578, AA551491, AW373072, AI890574, AW176422, AI439978, AI682971, AI587156, AI950729, AI570807, AW169234, AW104641, AI498067, AI932794, AW007555, AI702073, AI700159, AW082532, AI961589, AW198090, AL079963, AI637584, AW104724, AI433157, AI254727, AA514684, AW075381, AI270183, AI611738, AI633125, AI923370, AI434468, AI924701, AI286256, AW192652, AI584140, AW080327, AI274655, AI591420, AI698391, AI889189, AI768496, AI282903, AI801793, AL119863, AI927233, AI056328, AI539800, AI570909, AI922689, AI926367, AI624693, AW074374, AI670009, AW081866, AI538259, AW103878, AW026882, AI468872, AW129230, AI818353, AW051059, AI590227, AI284084, AI815232, AI445432, AI915291, AW089275, AI288285, AW080090, AI832245, AL045500, AW166583, AW148363, AI619748, AW169132, AI571867, AA908294, AI619691, AL121041, AW051258, AI281687, AI439452, AI590686, AI648454, AW163834, AL040390, AI500061, AI684234, AI524677, AI678688, AI677796, AI864836, AI934259, AW075667, AI267162, AI522052, AW118518, AI933992, AI859915, AI678446, AI567128, AI697137, AI609375, AW087199, AI582932, AI291973, AW148294, AI355827, AI684244, AI497733, AI613038, AI659334, AI963191, AI955943, AI493567, AI571439, AI862139, AI474146, AI802542, AI951076, AI471712, AL110306, AI609409, AI619502, AI889376, AW193530, AW132056, AI799470, AI866465, AI648509, AI572021, AI564719, AI287326, AI929108, AI440448, AI478723, AI702068, AI696398, AW080746, AI554821, AA848053, AI423326, AI445611, AW268964, AI288305, AW168452, AI933783, AI952145, AW020693, AL047100, AW081653, AW078895, AI633062, AW152182, AI539780, AI612913, AI573060, AW088560, AI270099, AI679550, AI863241, AI866751, AW168677, AW083573, AI627893, |
|--|--|--|--|---|

| | | | | |
|--|--|--|--|---|
| | | | | AI571909, AI280661, AI640704, AI432736, AI537617, AB020649, AL137297, AL137459, A65340, AL080159, A18788, I89947, AL050366, AF069506, AF159615, AR034821, AF215669, AL133619, I48978, I66342, AL133093, AL080139, A77033, A77035, AF017437, A08916, A08913, AL049283, S36676, AL133016, AF100931, Z37987, AL050149, AL133113, M92439, A08910, U00763, X82434, AL137526, A08909, AL117460, AF102578, A18777, AL050138, AF177401, X79812, I17544, I09499, U35846, AL049339, Y16645, I33392, AL137271, AB007812, AL133075, AL110296, AF183393, AL122100, AF026816, E15582, AF087943, AL117435, U37359, I00734, AF091084, I89931, X70685, Z82022, Y14314, E00617, E00717, E00778, E02221, AL137539, I49625, U72620, AF044323, AL137548, A15345, AL122110, AL137550, AL133568, AL050092, AF032666, AF008439, X72889, AL136884, AF113677, AL050024, AR038854, AL137463, AF113013, AL137480, AF067728, Y11587, AL117626, AF115392, AF031147, AL110225, AL110280, AF113019, AF028823, AL133558, I48979, AF118090, X63162, AF125949, AL137558, AF090901, Z97214, AL133560, E05822, AL122050, AF090900, U42031, A65341, AL122106, I03321, AF111849, S69510, AF139986, AL133010, AF104032, X06146, AF090903, U72621, AF182215, E01573, E02319, AL137665, AF118094, AF065135, AL133645, AL117583, AF176651, AJ238278, AL133557, AL110221, AL117457, A08912, S68736, AL080126, A03736, S61953, AL117648, AL050277, AF126247, AL137530, A76335, X80340, AF111851, U53505, AL137538, X84990, AL050116, AL137488, AF030513, AL137294, AJ005690, AL137479, AF067790, AL133104, AF113690, AF090934, AL080074, S63521, A08908, AF141289, AF079763, AL049452, Y09972, AL122045, A52563, L04504, S76508, AF061943, AR011880, A58524, A58523, I89934, AF113691, I26207, AL133640, E02349, AL117416, AL133565, AL080140, AL133665, AL080148, AL110224, AF106862, AF036941, X60786, AB016226, A57389, AF090943, AL049938, Y11254, AF180525, AL049382, Z13966, AF192557, X52128, E01314, AL137529, M27260, AL137256, S77771, AL137658, A27171, S78214, U75932, Y10655, AF113694, S78453, AL137557, AF106697, AL133080, AL049430, L40363, AF113699, AF132676, AF061836, U49434, U80742, AL050155, AL050393, D83032, U88966, E02152, AF114170, AL137283, |
|--|--|--|--|---|

| | | | | |
|----|---------|----|--------|---|
| | | | | I17767, L04849, AL050146, AF057300, AF057299, AL133112, X53587, A86558, L31396, U87620, X57961, AL080137, AL137555, AL133606, U91329, L19437, AF097996, AL049347, AL049314, E06743, AR059958, AL122098, and AF017152. |
| 20 | HDPSB18 | 78 | 732097 | AA640305, AA180056, AA595661, AA199578, AI523205, AA631915, AA568303, AI270280, AI086603, AI754064, AI348780, AA489390, AI753131, AA493546, AW239465, AA177130, AI570067, AI690379, AI567676, F34376, AA157876, N27737, AA730530, AI174703, AW272606, AI345256, AI610814, C15966, AA599712, AA608729, AW238242, AL039436, AL121652, AL009181, AC006449, AL133448, AP000048, AP000116, AL022316, Z99716, AL034549, AC005519, AC004522, AL035086, AC003689, AF207550, AC005962, AC005015, AF196779, AJ003147, AC007225, AL035458, AC007546, AC002470, AP000065, AC008072, AC005081, AC005067, AC007934, AF053356, AL031681, AC003101, Z84466, AL133371, AL008718, AC004150, AF038458, AP000505, AB023048, AC003029, AP000692, AC002504, AP000691, AP000503, AC005332, AC016025, AC003665, AC006115, AL049569, AC005228, AL139054, AC006597, AC007227, AF134726, AC004125, AC006013, AL050332, AL137100, AL117258, AL020997, U80017, Z93023, AC009516, AC010205, AC005225, AC004230, AC006014, AL008582, AC005520, AC006285, AC005041, AC002984, AD001527, AC004967, Z93244, AC005031, AL049636, AC002310, AC007055, AC002350, AL034420, AC007298, AC007021, AC005037, AP000694, U85195, AC005184, AC004477, AC005377, AC005412, AL132712, AC005488, AC005921, AL022721, AP000512, AC003982, AC005695, AC005722, AP000555, AP000556, AC004812, AC005004, AC005785, AL031733, AC006948, AL136504, AC005730, AC006538, AE000658, AL031680, AC005971, AC012627, AC006042, AC007151, AL049694, AF001548, AC005913, AC004019, AL022315, AL050318, AC005180, AL121603, AC004771, AF111169, AC008055, Z95116, AC007676, AL136295, AF196971, AC004878, AC005071, AC004854, AC005531, AB017602, AC004000, AF111168, AC009247, Z83838, AC003043, Z97632, AC002314, AL022326, AC003962, AP000140, AC006468, AC004491, AL035659, AC006441, AL031311, Y14768, AC007308, |

| | | | | |
|----|---------|----|---------|---|
| | | | | AL035460, AC007030, AC006509, AC004099, AC004821, AP000030, AC005821, AC006211, AL031284, Z93928, AC006312, AC005300, AC005778, AC006236, AC004815, AP000696, AC005049, AF111167, AC000353, AL079342, U52112, U91325, AF129756, AC005529, AC007066, AC003108, AL109758, AC007201, AC018633, AC005399, AC002316, AL022163, AC004876, AC007686, AC005808, AC000052, AL022323, AP000088, AC005527, AC002551, Z97056, AL078581, AC004913, AC006160, AC005736, AL109798, AC005089, AC007216, AC005940, Z97054, AF205588, AP000355, AF200465, AC005086, AC004531, AC007536, AP000501, AC004638, AP000553, AC005011, U07000, Z98941, AP000040, and AP000108. |
| 21 | HDPTW65 | 31 | 912262 | AA312170, AA019750, AI681557, AL136168, AL096705, AC007786, and AC002357. |
| 21 | HDPTW65 | 79 | 906717 | AA312181, AA019750, AI681557, AL096705, Z98046, AC007786, and AC002357. |
| 21 | HDPTW65 | 80 | 762318 | AA312170, AI432644, AI623302, AW081103, AI432653, AI431357, AI431337, AI432665, AW128900, AI431241, AI432654, AI431312, AI431345, AI431353, AI431307, AI431316, AI492519, AI431354, AI431310, AI431255, AI432676, AI431243, AI432661, AI431351, AI432649, AI431347, AI431230, AI431248, AI431328, AI432655, AI431330, AI432672, AI432674, AI431346, AI432651, AI432647, AI432677, AI432675, AI431254, AI432662, AI432673, AI432658, AI431340, AI432650, AW128846, AI432664, AI791349, AW128897, AI492509, AI492520, AI431751, AI431236, AI431308, AW128884, AI492510, AI432643, AI432657, AI431335, AI431247, AI432645, AI431342, AW129223, Y17793, and AF064854. |
| 22 | HOHDF66 | 32 | 1020499 | AA742544, AA009717, R81588, AA009827, AI381577, AI677718, and AI685149. |
| 22 | HOHDF66 | 81 | 906092 | AA742544, AA009717, R81588, AA009827, AI381577, and AI677718. |
| 23 | HDPTW24 | 33 | 843591 | AL046746, AW302903, AI270019, AW189113, AI819140, AI744348, AA683238, AA493695, AA809546, AA526630, AI433008, AI345157, AA279421, AA634889, AW002350, AA782526, AW408047, R93627, AA181478, AI110595, AW328331, AL041619, AW407578, AI687343, AA757775, AI270117, AW069769, AI890324, AI611506, AI271028, AW022897, AI268334, AI268336, AI305766, AI963720, AI953275, AI932599, AW021735, AA490183, AI056913, |

| | | | | |
|----|---------|----|---------|--|
| | | | | AA661921, AI309375, AA486131, AI933579, AW275510, AL119915, AA503600, AA747070, AA236903, AW168420, AW068596, W45625, T70715, AL119691, AA092571, AA847181, AI754653, AI254913, AA468022, AA663512, AW028392, AA631251, R24887, AA318397, AW271069, AI821877, AW360896, AC016831, Z98043, AC008249, AC005082, Z94056, AC000355, AC004595, AL121825, Z76735, AC004134, AL035696, AC006511, AC003665, AL031848, AC007285, AC004821, AC005193, AC006373, AC002067, AL049745, Z84814, AC004525, AC002377, AL031846, AC011311, AL118497, AL121576, AC005233, Z93023, AL022328, AC002470, AL049647, AC004773, AC003681, AC000120, AC005209, AL096791, AC004750, AC005826, AC004985, AC006379, Z84469, AC007115, AC004895, AL049831, AL133500, AL109847, AC005664, AL033521, AL033397, AL109653, AL022336, AC004612, AC002127, AC000097, Z95118, AC005325, AC006547, AC004806, Z99716, AL035458, AC005503, AC002369, AC005862, AL008629, AL031428, AL034384, AF049895, AL133249, AC005666, AC006059, AC006120, AC002565, AC006449, AC002536, AL049825, AC000372, AC004104, AC009330, AL031670, AL022165, AC002451, AC007848, AC006479, AC005971, AC005261, AC005901, AC005730, AC003695, AC002115, AL109865, AC005342, AC005237, AC006430, AC002541, AC000073, AC004552, AC007055, AC004816, AC005084, Z82198, AC003664, AC006071, AC004647, Z98046, AL023578, AC004470, AC005019, AC003109, AC008372, AC008012, AC006374, Z85986, AL049543, AC004881, AC006017, AL122126, AC004448, AC010197, AL049844, AC005699, Z98742, AL031733, AC002036, AC000083, U82828, Z98052, AC009248, AL049546, Z78022, AL031277, AC004409, AF001550, AC005518, AC007237, AC004875, AL035423, AC005324, AL109753, AC005993, AC007253, AC004603, AC006539, AL049874, AC000085, AC004986, AL023807, Z98304, AL121652, AC010200, AC006070, AP000011, AP000152, AL117355, AC009223, AC002487, AC003071, AC006207, AC004933, AC005280, AC002538, AC009542, AC002300, AB015752, and U57005. |
| 24 | HRADO01 | 34 | 1018804 | H68792, AL037632, AI110760, AA577824, |

| | | | |
|--|--|--|---|
| | | | AI054414, AA223932, AL047602, AW102811, AA493641, AI016704, AW406162, AA601355, AI457389, AW079809, AI216054, AA010672, AI567674, AL135377, AA487475, AL046746, AI734144, AA650623, AA643454, AA643457, AA634991, AA157876, AA046906, AA665449, AA569182, AW243793, AA808796, AW055226, AA601278, AA643452, AA508036, N66012, AA323343, AA569284, AA857296, AL135698, AL045053, AW265073, AA504951, AI433104, AA812139, AA126635, AA613164, AA019793, AW085751, AI446474, AI791227, AI445815, AL041013, AW089861, AI580906, AI734070, AI144101, AI859834, AI907046, AW268973, H45320, AI125167, AI865213, AI653542, AW402458, AC002070, AL009181, AF064857, Z99570, AC005229, AL022323, AF017104, AC004382, AC002538, AC004929, Z84469, AL050318, AC018633, AF003626, AP000365, AC007649, AC002378, AL132994, Z97181, U91321, AF111167, AC005844, AP000688, AL033525, AF141325, AC005224, AC002350, AL024508, AC007899, AC005971, AC002287, Z85996, AC004531, AC005531, AC005822, AC005553, AC003695, AC004448, AC005291, Z98751, AC007240, AC008009, AL031775, AC004967, AL031664, AC005046, AC006948, AC005324, AP001037, AC007225, AC008101, AL035420, AC005863, AL034384, AC005274, AC007065, AL022165, Z95152, AC008042, AC006120, Z98946, AL050321, AF111168, AC004600, AC002316, AC005271, AL078593, Z98949, U07000, AC004042, Z93017, AC005181, AL023807, AC004408, AC007363, AP000030, AL034399, AL031597, AC002077, AC005516, AC002377, AL031655, AC008033, AL096712, AC005666, AC005703, Z95113, AC007314, AC000094, AC004966, AF196970, AL024507, AL049709, AC005914, AC005284, AP000347, AC006211, AC003046, AC005736, AC006501, AC005018, AP000346, AL050305, AC002091, AL031120, Z98052, AC006445, AC007406, AC004259, AF067844, AL034553, AC002302, AL133321, AC005094, AP000033, AC006026, AC005826, AC006121, AL009179, AL035088, AC006450, AL034379, Z98304, AC000085, AC007878, AC005728, AF099810, AL049759, AC006379, Z95889, AC004112, AC004905, AL023876, AL008723, AC006441, AP000350, AC004491, AC009069, AC012384, AC006057, AC003684, AC005193, AL021391, AF207550, |
|--|--|--|---|

| | | | | |
|----|---------|----|--------|--|
| | | | | AL049869, AC005879, AL049873, AC002563, AL049872, AP000031, AL049779, AC009464, AL021453, AL049836, Z98036, AC007285, AC002300, AL034555, AL078621, AC005017, AL035587, Z99916, AC007386, AL049569, AP000115, AL021368, Z98752, AC004087, AC009181, AL031665, AC004817, AC006059, AL031257, AC005539, AC007277, AC003003, AC007666, AC006226, AJ006345, AC005747, AL049646, AC006530, AP000557, AC006080, AL031311, AL031123, AC004020, AL033381, AC007358, AC003025, AF134726, AC006115, U91327, Z93016, AC005155, AC006481, AP000354, AL022336, AC005920, Z83845, AL049832, AC002565, AF139813, AP000500, AC003982, Z93930, AL035653, AP000326, AL031575, AC006387, AP000356, AC004941, AC004816, AC005036, AC010202, AC005152, AC004662, AC004228, AF003529, AC004675, AC005740, AL035409, AC007463, AC008498, AP000169, AP000054, AL035089, AC002369, AC006360, AC004066, AC006254, AC005323, AF107885, AC005969, AC006203, AL022322, Z97055, AC005900, AC004168, AC005358, AC004983, AF109907, Z85987, AL121748, AC007546, Z86090, Z95116, AF088219, and AL109654. |
| 24 | HRADO01 | 84 | 800740 | H68792, AW102811, AA601355, AA493641, AL135377, AA577824, AA634991, AA569284, AA569182, AA010672, AA508036, AW055226, AI433104, AA643454, AA643457, AW265073, AI580906, AI457389, AA613954, AA643452, AA808796, AA223932, AW089861, AA577906, AL041013, AI446474, AI653542, AI859834, AI284640, AI865213, AW402458, AL046409, AA601680, AI431303, AI340641, AW021917, AA501614, AA857296, AW075979, AI754291, AA984263, AI890348, AA847333, AW193265, AI952900, AI623899, AI251111, AI345157, AW088846, AI859946, AW276827, AA188940, AA349366, AW062724, AI350211, AI053445, AI753951, AI754013, AW192330, AI049996, AI679713, AI358513, AI613280, AI885572, AW327624, AI493264, AI628224, AI755236, AA525293, AI754923, AA682912, AI016704, AI862802, AL046746, AI343143, AI298246, AA812058, W23546, AA628627, AC004106, AL121653, AC006011, Z86090, AC008372, AC005839, AC004381, AC002470, AC005527, AC005081, AL031311, AC004596, AL022320, AP000555, L47234, AC006449, AL034420, AC006480, AC000025, |

| | | | | |
|--|--|--|--|---|
| | | | | AC005288, AC005387, AC005971, AC002544, AC004887, Z83840, AC004386, AC004263, U91326, AL035249, AC009464, Z95331, AC006254, AC003029, AC016025, AC004854, AL031281, AC008249, AC005632, AL121655, AL049776, AF134726, AC006285, AC003051, AC006312, AL020993, Z84469, AF015149, AC005037, AC006257, AC005057, AP000365, AF064863, AC005921, AC003101, AC005375, AC005488, AC005899, AL021397, AL035681, AC004814, AC006468, AF111168, AC004927, AC005412, AC002425, AC005215, Z99716, AC005225, AC002350, AL132777, AL031283, AP000356, AL034429, AC004967, AL109952, AL135744, AC007207, AP000556, AC006511, AP000557, AP000248, U80017, AL022323, AC004819, AC007055, AC002996, AL096701, AC007536, AC005291, AL024507, AC005280, AC005911, AC007690, AC004257, AL049779, AC005520, AC005519, AL121694, AL133245, AC006077, AC005531, AL008582, AC000353, Z98051, AC008101, AC005324, AC005529, AJ003147, Z98304, AL008635, AC006064, AC002430, AC005193, AB023050, AL049745, AF109907, AC005678, AC007324, AC005261, AC005972, AC002543, AC006441, Z83844, AL022328, AC004019, U95742, AL049760, AC004590, AC009516, AC004821, AC005354, AC005406, AL031659, AC005277, AC004024, U91325, AC005837, AC007384, AC007686, AC002418, AL080243, AF196969, AC004999, AC009247, AL133249, AC007066, AC008079, AC002369, AC005512, AC004056, AC006120, AC007227, AC004638, Z84480, Z99755, AC003098, AF091512, AC004905, AC004859, AP000207, AP000129, AP000503, Z98752, AC005102, AC007999, U47924, AL121748, AC002365, AC005768, U95740, AC002312, AP000502, AC007528, AC006487, AL049780, AL035659, AC004983, AC009181, AC002301, AC005581, AP000961, AF038458, Z98742, AP000030, AL121657, AC007161, AP000347, AL034423, AL035419, Z95114, AL031666, AC005409, AC007919, AL049761, AC000003, AC005393, AL078581, AC002563, AL109798, AC004816, AC007051, AF165926, AC004491, Y10196, AL109627, AC007216, AC005740, AC007312, AL031282, AC020663, AC005562, AP000553, AC005399, AL078475, AC003007, Z97054, AF196971, AC006132, |
|--|--|--|--|---|

| | | | | |
|----|---------|----|--------|---|
| | | | | AC005701, AC005919, AC005231, AB023049, AC004796, U85195, AC004858, AC005589, AC004895, AL031848, AC004797, AL096791, AL035415, AL023553, AL023577, AL049576, AC004953, AL031650, Z84814, AC006539, AC006211, U35114, L44140, U91323, AP000692, AC004227, AC004815, AL022318, AF001549, AL022165, AL023882, and AD000092. |
| 25 | HTJNX29 | 35 | 904831 | AA699986, AI831572, AA707224, AA316671, AI826553, AW062507, AA494546, AW362519, W23003, AW118600, AI276610, T98080, T98002, AI761193, W88598, W89099, R08175, R11209, R11267, R08229, T85456, AA694114, R29134, AA377260, AI365256, AI224373, AI886055, AW089275, AI348777, AI335426, AI677797, AW054939, AI400898, AI656270, AW089006, AW151034, AI567302, AI784028, AI345415, AI689614, AI829495, AW149069, AA765656, AI680498, AA911767, AI633419, AW087863, AL040011, N80094, AI582954, AW301865, AW029457, AI911620, AI932877, AA190341, AI287476, AI915443, AI360195, AA514684, AI434242, AI921155, N89459, AA641644, AI872148, AW008166, AW003866, AI824360, AI523771, AI432532, AI682891, AW019985, AW089932, AI919345, AI084188, AW080379, AI371984, AI660432, AI860830, AA627701, AI364135, AW265004, AI283996, AI472536, AW025279, AL043288, AI687465, AW080076, AW083750, AW327693, AI932739, AA693347, AI628325, AI702540, AW089293, H43387, AW085786, AI250627, AI074150, AW163823, AI560023, AW162189, AI471548, AI421662, AI419440, AI345612, AI499483, AI339115, AI918554, AI873704, AI633061, AW089179, AI345416, AI433611, AI861982, AI934147, AW085373, AI635528, AW087837, AI872722, AI250646, AI784214, AW263569, AA575874, AL042729, AI879377, AI921386, AW194014, AI702091, AI285417, AW151451, AW074702, AA665669, AA743385, AW008434, AI887268, AI697324, AI571699, AI452993, AW132001, AI636187, AI678446, AI648502, AI344911, AI571868, AI446698, AI866465, AI570966, AI559863, AW080374, AI886355, AA999906, F26535, AW022955, AI611215, AW162071, AI453328, AI306706, AW074605, AW149236, AI241792, AW149221, AI627714, AA743531, AW029579, AI961631, AI537617, AI953803, AI690410, AA587590, AI799244, H42948, AW079768, AW078680, AI688894, AW102900, AW020419, AW087928, |

| | | | | |
|--|--|--|--|---|
| | | | | AI699154, AI628875, AI539111, AI818213, AI335338, AA830821, AW302924, AI888621, AA808175, AI887785, AI446775, AI247298, AI569294, AI520809, AI251221, AI539153, AI339932, AI287862, AI589391, AW150511, AI379711, AI814896, AI933992, AI597805, AW149876, AW190297, AI469505, AI689096, AW080766, AI873638, AI586984, AW167422, AW132104, AA130341, AA916133, AL048499, AI370812, AA693331, AW148303, AI307505, AL079794, AA928539, F37364, AI345745, AI921167, D12620, AB002454, D26480, AF054821, AF133298, U02388, M94548, U39206, D12621, U39208, AC004523, U39207, AD000685, AD000091, AC005336, AC004597, AB002461, AL137712, AB002456, AB002460, AB015302, AB015298, AB015299, AB002457, AB015305, AF094480, I17544, E03348, E03349, AL117460, Y13350, Y11587, U77594, AF067728, U42031, AL137658, AL122098, AF054988, AJ003118, AF158248, AF036941, AF082526, E01812, S77771, AL137429, AL050155, AL137476, AF077051, AF054289, E04233, AL050172, AF043642, AL137557, AF090901, X96540, AR029580, AL137271, AF065135, A90844, U90884, E01963, U75370, X79812, AF030165, AL050278, I66342, S69407, AL133088, M86826, AL137574, AL137256, AC004227, Y08769, A18777, AF067790, E12580, AF090886, I30339, I30334, AL137300, AF038847, A65336, A08913, U68233, I92592, AR068466, E02221, AR038854, U95739, I22272, AF081366, U83980, E12579, S69385, AF155119, A08912, AL137711, AF199027, AL137254, X59414, A08910, AF032666, A08911, A08909, AL137627, AC006039, AF118094, AF054986, AL133099, U96683, AL137641, E06743, AL137548, AR038904, A08907, A08908, L10353, AL080154, AL137536, AF114784, AL117585, AF109155, S76508, AL137480, AF016628, I48978, AL137523, AF111851, AL117416, AF026008, M64936, AF035161, U92992, L19437, Y18680, AL133070, AF089818, AL133645, AF126488, S83440, U73682, AL080150, AF185576, AL137521, AF100931, AF017437, A65341, AL050024, AL080146, U02475, AR000496, X62580, U39656, Z82022, AL117649, AF047716, J05032, AL080162, M27260, AF061795, AF151685, AL049426, X83508, X72889, U49908, AP000697, Z49258, AL050170, X66871, X70514, AB007812, AF002672, U88966, X99226, AF126372, AL133084, AL117587, Z72491, |
|--|--|--|--|---|

| | | | | |
|----|---------|----|--------|---|
| | | | | AL110221, AF162270, AC004686, E01614, E13364, AL137268, E02152, X53587, AC002467, AL117648, and AR013797. |
| 25 | HTJNX29 | 85 | 906748 | AI826553, AW362519, AW062507, AI761193, T85456, AA316671, R08229, AW118600, AI400898, AC004523, D26480, AF054821, AF133298, AD000685, AC005336, U02388, D12621, AB002454, D12620, AD000091, AB002459, AB015302, AB015299, and AB002457. |
| 25 | HTJNX29 | 86 | 906746 | AA699986, AI831572, AA707224, AA316671, AI826553, AW062507, AA494546, AW362519, W23003, AW118600, AI276610, T98080, T98002, AI761193, W89099, R11209, R08175, W88598, R11267, R08229, T85456, AA694114, AA377260, R29134, AI400898, D12620, AB002454, D26480, AF054821, AF133298, U02388, M94548, U39206, U39208, AC004523, AD000685, AC005336, AD000091, U39207, D12621, AC004597, AB002461, AB002456, AB002460, AB015302, AB015298, AB015299, AB002457, and AB015305. |
| 27 | HLDQA07 | 37 | 992926 | AW235556, AW207411, W72972, AI580378, AA563952, AI655152, AI301835, AW129556, AI804914, AI279596, AI620860, AI680592, AA932836, AW024966, AA578896, AI912213, AI671481, AI308062, AW339355, W37838, AW189426, AA886938, W76209, AI061211, AI739650, AI537273, AI799408, AI802542, AI927755, AI921248, AI500061, AW170635, AW151786, AW198090, AI306613, AI620089, AI627988, AI670009, AI632408, AI678357, AI269862, AI933589, AI281772, AI886181, AI445992, AI613038, AI890507, AI635067, AI611738, AI636588, AI335426, AI521560, AI348777, AW238688, AI619502, AW118518, AI633125, AI559586, AW054964, AI932794, AI366900, AI524671, AW051258, AW117746, AI677796, AI352497, AA449768, AI288305, AI570807, AW026882, AI923370, AW050522, AI433157, AI288285, AI702073, AI627893, N33175, AW268220, AI334450, AI874166, AW268302, R32821, AW051088, AW072719, AI698391, AW264727, AI915291, AW152182, AI768496, AI582932, AI889376, AI889189, AI554821, AI261344, AI445829, AW149925, AL121286, W74529, AI886123, AW148536, AI275640, AI570861, AI591420, AI681985, AI886753, AI537261, AI620284, AI281782, AW169604, AI280732, AI680388, AW081036, AI866770, AI690748, AI568374, AA806720, AI890214, AI934011, AI568138, AI569309, AI799273, AI869377, AL046944, AI564719, AI859464, AI358701, AL036923, AL045500, AI868740, AI274508, AI702019, |

| | | | | |
|--|--|--|--|--|
| | | | | <p> AW161156, AW083573, AI963193, AI624293, AI439745, AI620075, AI571909, AI619607, AI680162, AI564749, AI569583, AW084425, AI635016, AL039132, AI174394, AW090393, AL036634, AI868931, AI565062, AI917252, AI688858, AI873604, AI648509, AI572021, AI537677, AI871977, AI432030, AW104108, AI868204, AI799183, AI499285, AW168485, AI824576, AW131294, AI783504, AL134259, AL119863, AI608936, AI269205, AI798456, AW268122, AI445990, AW080357, AI335209, AI609127, AI345148, AI308032, AI926790, AI863382, AI491897, AW301513, AI932638, AI475408, AW088134, AI783997, AI873644, AI306705, AW151136, AI431975, AI344785, AI815232, AW029197, AI620003, AW163834, AI686817, AI270183, AI611354, AW150453, AI432218, AI473536, AW148363, AI287489, AI890833, AI580984, AI537837, AI922365, AW079075, AI452707, AI590415, AI637748, AL036673, AI281762, AI497733, AL040241, AW059713, AW151893, AW264895, AI310575, AL037582, F27788, AI925196, AL037602, AA579618, AW080327, AI609409, AI241923, AL039086, AW076093, AI801460, AI637584, AA470491, AI340533, AI521103, AI804983, AI824746, AI538829, AA279293, AW166970, AW118496, AI801152, AI874261, AI432040, AI800155, AW083778, AI863321, AI963846, AI955917, AI696398, AA225339, AL137448, AL080159, I89947, AL110280, Z82022, I48978, Y14314, AR038854, AF026816, U67958, A77033, A77035, A08913, AF090903, AL023657, A08916, AL080074, A08910, I48979, A08909, AL133640, AF183393, AL080124, U80742, AL137271, I89931, AF106862, AL049452, I49625, U35846, A58524, A58523, AF185576, AL137480, AF119337, AF087943, AL050149, AF118094, A08912, AL049382, AL117460, I33392, AF111849, AL117435, AL137463, X82434, AL050138, AF091084, AL133067, AF113677, E15569, AF079763, L19437, AL137550, A93350, AF125948, AJ012755, AF113019, A65341, AJ238278, AF026124, E07108, AF177401, AL050277, AL050393, AF153205, AL117440, X72889, AL049938, AF104032, AL133075, A45787, AL133560, AF061573, AF113690, AF090934, AF017437, E04233, AL122050, AF113013, AL137533, U00763, AF061943, AR011880, AF078844, AL110196, AF097996, Y11254, Z72491, AF111851, AL137459, AL122093, U42766, AL049283, Y09972, AF162270, A03736, Y11587, X96540, AF113691, AF090900, AL050116, L31396, L31397, X63574, </p> |
|--|--|--|--|--|

| | | | | |
|----|---------|----|--------|--|
| | | | | AL137476, A90832, E02349, AJ242859, AL049464, AL133016, I00734, AL137488, S61953, AF113694, AL137556, E03348, AF090943, AF067728, AL050024, AJ000937, AL049430, AF113699, AL137560, A07647, E00617, E00717, E00778, AL050146, AF090901, AL122121, AL122123, AB019565, AL133104, AF003737, AL137557, AL133080, X70685, AL049314, M30514, AL137648, AL137538, AL117585, AF017152, AF125949, AF113676, U68387, AF158248, AL110225, AL117394, AL133113, A12297, X65873, AL133606, U91329, AF057300, AF057299, X98834, I03321, L30117, S78214, AL117457, AF061981, AF146568, AF111112, AL122110, AL049466, AF113689, X87582, E05822, AR059958, X84990, AL110221, AL133565, I26207, AR038969, Y16645, AR000496, U39656, I42402, AL117583, AL133557, AL050108, AF090896, Y07905, X93495, AL133072, AF079765, AL137521, AL080060, AL122049, AL137526, AL049300, AF118064, I09360, AL133093, AF118070, U58996, AL133098, AL122098, AL080127, U96683, AL096744, AL133077, S68736, AL133568, AL133014, AL080137, AL137527, AF008439, U72620, E08263, E08264, E07361, A93016, AR020905, AL137273, AL137478, AL050172, U78525, X81464, AL110197, AL117416, Z37987, AL050092, AL137292, AL133665, AF061795, E08631, AF151685, AF139986, AJ006417, AF100931, AL133081, AF032666, AL137294, AL080148, AL080086, AL117432, E02221, AF051325, AL137529, AF210052, AL122118, E12747, AF067790, U49908, AF106827, X53587, and AL137283. |
| 27 | HLDQA07 | 89 | 882769 | AW207411, AW129556, AI279596, AI620860, AI671481, AW024966, AA563952, AI804914, AI912213, W76209, W72972, AI680592, AI301835, AI308062, AW189426, AA578896, AI580378, AW235556, AW339355, AI739650, W37838, AI655152, AA932836, AA886938, AI061211, AI537273, AI627988, AI802542, AW151786, AI921248, AI500061, AW170635, AW198090, AI799408, AI306613, AW149925, AI670009, AI632408, AW050522, AI678357, AI269862, AI933589, AI281772, AI445992, AI635067, AI611738, AI521560, AI636588, AI619502, AW118518, AI633125, AI564719, AI591420, AI889376, AI932794, AI524671, AW051258, AW117746, AI677796, AI352497, AA449768, AI288305, AI570807, AW026882, AI923370, AI627893, AI620089, AI433157, AI288285, AI702073, AI681985, AW268220, N33175, AI334450, AI874166, AW268302, AI698391, AI927755, AI915291, AI582932, AI768496, |

| | | | | |
|--|--|--|--|---|
| | | | | AI889189, AI261344, AI554821, AL121286, AW081036, AI570861, AW148536, AI917252, AI886753, AI620284, AI281782, AW169604, AI886181, AI866770, AA806720, AI274508, AI890214, AI926790, AI934011, AI569309, AI799273, AI868740, AI702019, AI613038, AI335426, AI348777, AL045500, AI890507, AI680388, AI619607, AI470648, AI270183, AI963193, AI439745, AI571909, AI680162, AI569583, AL039132, W74529, AW084425, AI174394, AI890833, AI648509, AI635016, AI688858, AI873604, AI565062, AI572021, AW238688, AI537677, AL134259, AI269205, AI499285, AI799183, AI824576, AI620003, AI609127, AI783504, AI559586, AI798456, AW054964, AI366900, AI804983, AI608936, AL119863, AW268122, AI491897, AI335209, AI280732, AI431975, AW080357, AI308032, AI932638, AW088134, AI873644, AI306705, AW151136, AI432218, AI344785, AI815232, AW152182, AW163834, AI686817, AW150453, AI609594, AI287489, AI611354, AW148363, AW051088, AW083573, R32821, AI580984, AW072719, AW079075, AI452707, AI590415, AI281762, AI637584, AI497733, AL040241, AW059713, AI925196, AW080327, AI310575, F27788, AL037582, AL037602, AI521103, AA470491, AW076093, AW161156, AI609409, AL039086, AI696612, AW264727, AI801460; AI955917, AI538829, AI340533, AI824746, AI537244, AI801152, AL036923, AW118496, AA279293, AI874261, AW166970, AL046944, AL036634, AI445829, AI452560, AI800155, AI963846, AW132056, AA225339, AI862139, AI689420, AI610307, AI696398, AW118557, AI284131, AI886123, AI619737, AI868204, AI680463, AI932966, AI559296, AI285735, AI475806, AI286256, AL079963, AI860609, AI275640, AI340582, AI537303, AI445975, AL134999, AI280637, AI445990, AI521012, AI471909, AW151714, AW192652, AW071417, AI978703, AI922365, AI687375, AI569637, AI668893, AL137448, AL080159, I89947, AL110280, Z82022, I48978, A08913, AF026816, U67958, A77033, A77035, A08916, I48979, AF090903, Y14314, A08910, A08909, AL133640, AF183393, AL080124, AR038854, AL137271, I89931, AF106862, AL049452, I49625, U35846, A58524, A58523, U80742, AF119337, AF087943, AL050149, A08912, AF118094, AL049382, AL117460, I33392, AL137463, AL117435, X82434, AL023657, AL050138, AF091084, AF113677, AL137550, A93350, AF125948, X72889, AF113019, A65341, AJ238278, AF026124, AL050277, |
|--|--|--|--|---|

| | | | | |
|----|---------|----|---------|--|
| | | | | <p>E07108, AF177401, AL050393, AF153205, AL049938, AL133075, AL133560, AF113690, AF090934, AF017437, AL122050, U00763, AF079763, AF113013, AF185576, AL137480, AJ012755, AF061943, AF078844, AR011880, AF097996, Y11254, AF111851, AL137459, AL122093, U42766, X96540, AL049283, A03736, Y11587, AF111849, AF113691, AF090900, AL050116, L31396, L31397, X63574, E02349, AJ242859, AL049464, AL133016, AF113694, E03348, AF090943, AF067728, AL050024, AJ000937, AL049430, AF113699, AL137560, E15569, AL050146, AF090901, AL122121, AB019565, AL122123, AL133067, AL137557, AL133080, X70685, AL049314, AL137648, AL137538, AL117585, AF017152, AF125949, AF113676, AF158248, AL110225, AL117394, AL133113, A12297, X65873, AL133606, L19437, I03321, S78214, AL117457, AF146568, AF111112, AL122110, AL049466, AF113689, AR059958, X84990, AL110221, AL117440, AL133565, I26207, Y16645, AL110196, AL080074, AR000496, U39656, I42402, AL117583, AL133557, A07647, A45787, AL050108, AF090896, X93495, AL133072, AF079765, AL137521, AF061573, AF104032, AL080060, AL122049, AL049300, AF118064, I09360, AL133093, AF118070, E04233, AL122098, AL080127, AL096744, AL137533, AL133077, S68736, AL133014, AL080137, AL137527, U91329, U72620, E08263, E08264, E07361, A93016, Z72491, AF057300, AF057299, AF162270, Y09972, AL137273, S61953, L30117, U78525, AL137476, AL137556, A90832, AL110197, I00734, AL137488, X98834, AL133104, AF003737, E00617, E00717, E00778, M30514, U68387, AL050172, AF061981, X87582, E05822, U58996, AR038969, Y07905, AL137526, AL133098, U96683, AL133568, AF008439, AR020905, AL137283, AL137478, X81464, AL117416, Z37987, AL050092, E02221, AL137292, AL133665, AF061795, E08631, AF151685, AF139986, AJ006417, AF100931, AL133081, AF032666, AR013797, AF106827, AL137294, AL080148, AL080086, AL117432, AF051325, X92070, AL137529, AF210052, AL122118, E12747, and AF067790.</p> |
| 28 | HWDAD17 | 38 | 1001601 | <p>AW206008, AA421139, AI016583, AW129761, AA861899, AI161232, AA662496, AI394438, AW006729, AI633261, AI761456, AW193773, AI468509, AA632307, R82202, R63479, AA397617, AA400157, AW242664, AI422618, AI433392, AW003755, AA502416, AA400332, R36142, AA026216, AL121365, AW268072,</p> |

| | | | | |
|--|--|--|--|---|
| | | | | AI312542, AW020693, AI889376, AI677796, AL036396, AI569945, AI251221, AW152186, AI783504, AI343112, AW301300, AI349598, AA572758, AI348854, AW075207, AI564719, AI345735, AI340519, AI340603, AA579232, AW075413, AI313320, AI679764, AA420722, AL048656, AI923989, AI312146, AI312339, AI345258, AI950664, AI241763, AI345347, AI918449, AI923357, AI611743, AI334884, AI433157, AI702073, AI480118, AL039086, AI500706, AW075351, AI345745, AW022682, AI610488, AL036403, AL120853, AI439087, AW074172, AI815855, AW081255, AL045500, AI590530, AI251830, AL036631, AL079963, AW074993, AI349614, AI581033, AI873731, AW268253, AW083804, AL041150, AI349256, AI312152, AI824648, AI310575, AI923370, AI473554, AA427700, AW075084, AI620284, AI254727, AL119863, AI445992, AI358701, AI349937, AA977767, AI340533, AI538342, AI307708, AI866770, AI312325, AL036664, AI680435, AW152469, AI309401, AW403717, AW302965, AW410430, AL038605, AI587121, AA640779, AI307520, AI284517, AI334893, AA613907, AI499986, AI537677, AI500061, AI689420, AW302988, AI890806, AI889168, AI633125, AL119836, AI955906, AI698391, AI815232, AI801325, AI500523, AI915291, AI582932, AW088903, AW089572, AI345737, AI348897, AI521560, AI889189, AI500662, AA814407, AL036802, AI344935, AI866573, AI345736, AI635067, AW026882, AI582558, AI345397, AI888661, AA494167, AI446373, AI468872, AA287231, AI537515, AI573032, AI608936, AI364788, AI336585, AL036980, AI349787, AI345370, AI538885, AL041772, AI312428, AI310925, AI863321, AI866608, AL038565, AL036638, AI654750, AI619502, AI288285, AI745713, AW161579, AI539153, AI349226, AW078839, AI884469, AI432040, AI933589, AI445990, AW238730, AI307543, N71180, AI654389, AI439920, AA833760, AI866798, AL043975, AI671642, AI475451, AI431424, AL047344, AI859464, AW193134, AW105601, AW302992, AL036904, AI567612, AL036274, AI570909, AI570807, AI637584, AI554821, AI267502, AI589267, AW151136, AI867042, H89138, AI818225, AI926800, AI567351, AA012905, AA835801, AI349645, AW020095, AI680498, AW167918, AI862144, AI590423, AI500659, AI570989, AI805638, AW269097, AI690312, AI886753, AI873644, AL042382, AI612885, AI539771, AI540759, AL040277, AW023859, AW071380, AI345608, AI866510, |
|--|--|--|--|---|

| | | | | |
|--|--|--|--|--|
| | | | | <p> AW074869, AI274541, AI445237, AI611728, I48978, A08916, I89947, A08913, AL137523, A08910, A08909, I89931, AL049452, I49625, I48979, AL050024, AL049283, AL050149, AF177401, AL122050, AF090900, I00734, E00617, E00717, E00778, I09360, AL137550, X96540, AL133560, X82434, AF113699, AL049382, AL133568, AF090896, AF104032, Y09972, Y07905, A93016, AF113694, I03321, X52128, AF118064, AL133080, AL096751, Y16645, AF097996, Y11254, AL049430, X70685, AJ238278, AF090903, AL122093, X65873, AF026816, AF113677, AF067728, A65341, AF113690, A77033, A77035, AF087943, AL137271, E02349, AF106827, Z82022, AL137459, A93350, AF090901, AL050393, AL117435, U42766, AJ012755, Y11587, AL080124, AL137283, AL133558, AL110296, I17767, AL117460, AL050146, AR011880, AF113019, X84990, AL133075, U35846, AL137521, U91329, U88966, U00763, AL050277, AF078844, AL137526, AL049466, AF031147, AF017152, AL110221, AL096744, E15569, AL050108, AL117394, A08912, AF017437, AL133093, AL137560, AL133640, AL117583, AL122098, AL133557, AL050116, AL133016, AL137658, AF146568, S78214, AL133565, AL133031, AL049464, AB019565, AL122110, AF119337, AL133104, AF090934, AL110196, AJ000937, I33392, I42402, AF183393, AL137529, AL117457, L31396, AF113013, AL133072, AF079765, A03736, L31397, S61953, AF069506, AL049314, AF111851, AL110225, S68736, AB016226, AF090943, X80340, AJ242859, AL137527, AF061943, AF091084, AL049300, AF118094, AL137557, AF118070, AL137478, E06743, AL137648, AL117585, E07108, A12297, AF113691, AL133606, X63574, AF106862, X98834, U72620, AL122123, AF111112, E03348, I26207, AF113689, AL049938, U67958, AL080159, AF153205, AR059958, AF125948, AF125949, AL137705, AL133113, X93495, AL122121, X72889, A58524, AL137463, A58523, AL080060, AL137538, AF026124, Z37987, AL080127, AF113676, I66342, AL133077, AF158248, U80742, AL080137, AL050138, E07361, AF079763, AL110197, AF065135, AR000496, U39656, AL050172, U87620, A08908, AL137665, M92439, AR038854, AL133081, Y14314, AL110222, AL137479, AL110280, E01573, E02319, AF028823, AF100931, AL122049, AL133014, AF185576, AL080126, AF057300, AF057299, E08263, E08264, Y10655, A18777, AF132676, AF061836, AF192557, AL133098, AF017790, AF090886, </p> |
|--|--|--|--|--|

| | | | | |
|----|---------|----|--------|---|
| | | | | I89934, AF003737, AL137556, AF126247, E05822, AL080234, A07647, AL137488, Z97214, A90832, I09499, AL137539, AF008439, U75932, AL035458, AL133637, I80064, AL122111, X92070, E01614, E13364, and AR013797. |
| 28 | HWDAD17 | 90 | 883548 | AI394438, AW129761, AI761456, AI016583, AA421139, AA400332, AW206008, N73907, H93265, AA662496, AI161232, AA397617, AA861899, AI041233, AI468509, AW006729, AW193773, H93264, AA632307, AI633261, R82202, R63478, R82256, R63479, AA384241, D61543, AA400157, AA298161, AW242664, AI433392, AW003755, AA502416, AI422618, R36142, R36420, AA026216, H79569, AA977767, AI610488, AI619502, AI564719, AW075413, AI624548, AI636719, AI312542, AA833760, AL036187, AI826225, AI680498, AI828731, AA449768, AI569583, AW104724, AI561356, AI554821, AI802542, AI619716, AI633125, AI873644, AI174394, AW026882, AI863321, AI863382, AI433157, AI590021, AI445025, AI537307, AI269580, AI567351, AI569328, AL045500, AW149925, AI554218, AI702073, AI524671, AW081528, AW148408, AW075667, AA835801, AI476046, AI249962, AI583065, AI439920, AI634737, AI868204, AI625079, AI648663, AI569945, AI536638, AI824648, AI628205, AI537244, AI654750, AI498067, AW104827, AW262983, AI815237, AW152182, AW149311, AI538716, AI469532, AI635464, AI677796, AI873731, AI689248, AI434468, AI348854, AI367680, AI539808, AI889376, AI912288, AI274013, AI536685, AI610690, AI570989, AW129722, AI812080, AI612852, AI912434, AI687127, AI783861, AI282504, AI862139, AI439762, AI089970, AI812107, AI520785, AI635478, AW020693, AI697137, AW152186, AI922215, AI537303, AW074993, AI349614, AW268253, AI349256, AI312152, AI480118, AW075084, AL036396, AI745713, AW084447, AI349937, AI357316, AI334884, AI307708, AI312325, AA911767, AI307520, AI334893, AI926790, AI689420, AI924971, AI282355, AI961589, AW192652, AI799158, AI349226, AI251221, AI572787, AW193236, AL048656, AI340603, AI804585, AI566003, AW268072, AI628850, AI683173, AI590227, AA572758, AI355849, AW087534, AI340519, AW075351, AI590120, AA579232, AI874109, AA420722, AL120853, AI364788, AI923989, AI798303, AI307543, AI499393, AW081255, AW074172, AI446248, AI590530, AI367210, AI783504, AI493567, AI453322, AW169132, AI348897, AI609331, AI475371, AI799199, |

| | | | | |
|--|--|--|--|---|
| | | | | AI247293, AI634707, AW129106, AI273839, AI671679, AW075519, AI886206, AI362081, AI432969, AW149869, AW148320, AI686808, AW088805, AL036631, AI909661, AI310575, AW403717, AW302965, AW071380, AI343112, AI349645, AW074869, AW029401, AI500706, AW302988, AI340533, AW301300, AI349598, AI345745, AI613017, AW075207, AI687362, AW269097, AI804983, AW029611, AI678989, AI669616, AI345735, AI560023, AI887151, AI859402, AI918449, AW087455, AI801592, AI310925, AL036403, AI270183, AI312399, AW088903, AI345737, AI611743, AF113694, I48978, A08916, I89947, A08913, A08910, AF090900, A08909, AL049452, I48979, I89931, X65873, AF153205, I49625, AL110197, AL050149, AF090896, I00734, AL122050, AL050024, X70685, AL049283, AL049382, AF177401, AL137523, AL133568, I09360, A93016, AL049314, AL137550, X96540, AL133560, X82434, AF113699, AL122098, AF113677, E00617, E00717, E00778, Y09972, AL050146, AL080124, I03321, AL133640, X52128, AL117460, AL122093, AL049464, AF113019, AL096744, X84990, AL133075, AL110225, AL096751, U91329, E06743, AF078844, Y16645, AL049430, AJ238278, AL133557, AL110221, AF090903, AF067728, A65341, A77033, A77035, AF087943, X80340, AL137271, E02349, AF106827, Z82022, A93350, AF090901, AJ012755, U00763, AL137283, AF097996, AL110296, I17767, AL050393, AJ000937, S68736, AL137521, U88966, A58524, A58523, AL050277, AF118064, AL049466, AL133093, AF017152, E15569, AL050108, A08912, AL117394, AF113013, AL137527, AF104032, AF017437, AF113690, AL137560, AL117583, AL050116, AL133016, AL137658, AF146568, S78214, AF113691, AL117435, U35846, AL133565, AL133031, AF106862, AF061943, AL122110, AF119337, AF090934, AL110196, AL133080, I33392, I42402, AL137459, AF183393, AL137529, AL117457, L31396, AL133072, AF079765, A03736, L31397, AL137480, S61953, AL080060, AF118094, U67958, AF069506, AF111851, AF090886, AF125949, AL137705, Y07905, X63574, AL122123, AB016226, AF090943, AL049938, AF118070, Y11254, AL133098, AJ242859, AF026816, AB019565, AF091084, AL137526, I66342, AL049300, AL137557, Y11587, AL137478, AL137648, AR011880, AL117585, E07108, I30339, I30334, AL080137, AL050138, U42766, AL133606, X98834, U72620, |
|--|--|--|--|---|

| | | | | |
|--|--|--|--|---|
| | | | | AF111112, E03348, I26207, AF113689, AL133558, AL080159, AF031147, AR059958, AF125948, AL133113, X93495, AL122121, X72889, AL137463, AL133104, AL050155, AL137538, AF026124, Z37987, AL080127, AF113676, A12297, AF158248, U80742, E07361, AF079763, AF065135, A08908, AL050172, AL137479, AL137665, M92439, AR038854, AR000496, U39656, AL133081, U87620, Y14314, AL110222, AL110280, E01573, E02319, AF100931, AL122049, AL133077, AL133014, AF185576, AL080126, AF057300, AF057299, E08263, E08264, Z13966, Y10655, A18777, AF028823, AF132676, AF061836, AF017790, I89934, AF126247, E05822, AL080234, A07647, AL137488, AB007812, I09499, AR053103, AL137539, X56039, AF008439, I80064, AL133637, AL137556, AL122111, X92070, and E01614. |
|--|--|--|--|---|

Having generally described the invention, the same will be more readily understood by reference to the following examples, which are provided by way of illustration and are not intended as limiting.

5

Examples

Example 1: Isolation of a Selected cDNA Clone From the Deposited Sample

Each cDNA clone in a cited ATCC deposit is contained in a plasmid vector.

10 Table 1 identifies the vectors used to construct the cDNA library from which each clone was isolated. In many cases, the vector used to construct the library is a phage vector from which a plasmid has been excised. The table immediately below correlates the related plasmid for each phage vector used in constructing the cDNA library. For example, where a particular clone is identified in Table 1 as being

15 isolated in the vector "Lambda Zap," the corresponding deposited clone is in "pBluescript."

| | <u>Vector Used to Construct Library</u> | <u>Corresponding Deposited</u> |
|----|---|--------------------------------|
| | <u>Plasmid</u> | |
| | Lambda Zap | pBluescript (pBS) |
| 20 | Uni-Zap XR | pBluescript (pBS) |
| | Zap Express | pBK |
| | lafmid BA | plafmid BA |
| | pSport1 | pSport1 |
| | pCMVSPORT 2.0 | pCMVSPORT 2.0 |
| 25 | pCMVSPORT 3.0 | pCMVSPORT 3.0 |
| | pCR [®] 2.1 | pCR [®] 2.1 |

Vectors Lambda Zap (U.S. Patent Nos. 5,128,256 and 5,286,636), Uni-Zap XR (U.S. Patent Nos. 5,128, 256 and 5,286,636), Zap Express (U.S. Patent Nos. 5,128,256 and 5,286,636), pBluescript (pBS) (Short, J. M. et al., Nucleic Acids Res. 16:7583-7600 (1988); Alting-Mees, M. A. and Short, J. M., Nucleic Acids Res. 17:9494 (1989)) and pBK (Alting-Mees, M. A. et al., Strategies 5:58-61 (1992)) are commercially available from Stratagene Cloning Systems, Inc., 11011 N. Torrey

30

- Pines Road, La Jolla, CA, 92037. pBS contains an ampicillin resistance gene and pBK contains a neomycin resistance gene. Both can be transformed into E. coli strain XL-1 Blue, also available from Stratagene. pBS comes in 4 forms SK+, SK-, KS+ and KS-. The S and K refers to the orientation of the polylinker to the T7 and T3 primer sequences which flank the polylinker region ("S" is for SacI and "K" is for KpnI which are the first sites on each respective end of the linker). "+" or "-" refer to the orientation of the fl origin of replication ("ori"), such that in one orientation, single stranded rescue initiated from the fl ori generates sense strand DNA and in the other, antisense.
- 10 Vectors pSport1, pCMVSPORT 2.0 and pCMVSPORT 3.0, were obtained from Life Technologies, Inc., P. O. Box 6009, Gaithersburg, MD 20897. All Sport vectors contain an ampicillin resistance gene and may be transformed into E. coli strain DH10B, also available from Life Technologies. (See, for instance, Gruber, C. E., et al., Focus 15:59 (1993).) Vector lacmid BA (Bento Soares, Columbia University,
- 15 NY) contains an ampicillin resistance gene and can be transformed into E. coli strain XL-1 Blue. Vector pCR[®]2.1, which is available from Invitrogen, 1600 Faraday Avenue, Carlsbad, CA 92008, contains an ampicillin resistance gene and may be transformed into E. coli strain DH10B, available from Life Technologies. (See, for instance, Clark, J. M., Nuc. Acids Res. 16:9677-9686 (1988) and Mead, D. et al.,
- 20 Bio/Technology 9: (1991).) Preferably, a polynucleotide of the present invention does not comprise the phage vector sequences identified for the particular clone in Table 1, as well as the corresponding plasmid vector sequences designated above.

The deposited material in the sample assigned the ATCC Deposit Number cited in Table 1 for any given cDNA clone also may contain one or more additional

25 plasmids, each comprising a cDNA clone different from that given clone. Thus, deposits sharing the same ATCC Deposit Number contain at least a plasmid for each cDNA clone identified in Table 1. Typically, each ATCC deposit sample cited in Table 1 comprises a mixture of approximately equal amounts (by weight) of about 50 plasmid DNAs, each containing a different cDNA clone; but such a deposit sample

30 may include plasmids for more or less than 50 cDNA clones, up to about 500 cDNA clones.

Two approaches can be used to isolate a particular clone from the deposited sample of plasmid DNAs cited for that clone in Table 1. First, a plasmid is directly isolated by screening the clones using a polynucleotide probe corresponding to SEQ ID NO:X.

- 5 Particularly, a specific polynucleotide with 30-40 nucleotides is synthesized using an Applied Biosystems DNA synthesizer according to the sequence reported. The oligonucleotide is labeled, for instance, with ^{32}P - γ -ATP using T4 polynucleotide kinase and purified according to routine methods. (E.g., Maniatis et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press, Cold Spring, NY (1982).)
- 10 The plasmid mixture is transformed into a suitable host, as indicated above (such as XL-1 Blue (Stratagene)) using techniques known to those of skill in the art, such as those provided by the vector supplier or in related publications or patents cited above. The transformants are plated on 1.5% agar plates (containing the appropriate selection agent, e.g., ampicillin) to a density of about 150 transformants (colonies) per plate.
- 15 These plates are screened using Nylon membranes according to routine methods for bacterial colony screening (e.g., Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd Edit., (1989), Cold Spring Harbor Laboratory Press, pages 1.93 to 1.104), or other techniques known to those of skill in the art.

- Alternatively, two primers of 17-20 nucleotides derived from both ends of the
- 20 SEQ ID NO:X (i.e., within the region of SEQ ID NO:X bounded by the 5' NT and the 3' NT of the clone defined in Table 1) are synthesized and used to amplify the desired cDNA using the deposited cDNA plasmid as a template. The polymerase chain reaction is carried out under routine conditions, for instance, in 25 μl of reaction mixture with 0.5 μg of the above cDNA template. A convenient reaction mixture is
- 25 1.5-5 mM MgCl_2 , 0.01% (w/v) gelatin, 20 μM each of dATP, dCTP, dGTP, dTTP, 25 pmol of each primer and 0.25 Unit of Taq polymerase. Thirty five cycles of PCR (denaturation at 94 degree C for 1 min; annealing at 55 degree C for 1 min; elongation at 72 degree C for 1 min) are performed with a Perkin-Elmer Cetus automated
- thermal cycler. The amplified product is analyzed by agarose gel electrophoresis and
- 30 the DNA band with expected molecular weight is excised and purified. The PCR product is verified to be the selected sequence by subcloning and sequencing the DNA product.

Several methods are available for the identification of the 5' or 3' non-coding portions of a gene which may not be present in the deposited clone. These methods include but are not limited to, filter probing, clone enrichment using specific probes, and protocols similar or identical to 5' and 3' "RACE" protocols which are well known in the art. For instance, a method similar to 5' RACE is available for generating the missing 5' end of a desired full-length transcript. (Fromont-Racine et al., Nucleic Acids Res. 21(7):1683-1684 (1993).)

Briefly, a specific RNA oligonucleotide is ligated to the 5' ends of a population of RNA presumably containing full-length gene RNA transcripts. A primer set containing a primer specific to the ligated RNA oligonucleotide and a primer specific to a known sequence of the gene of interest is used to PCR amplify the 5' portion of the desired full-length gene. This amplified product may then be sequenced and used to generate the full length gene.

This above method starts with total RNA isolated from the desired source, although poly-A+ RNA can be used. The RNA preparation can then be treated with phosphatase if necessary to eliminate 5' phosphate groups on degraded or damaged RNA which may interfere with the later RNA ligase step. The phosphatase should then be inactivated and the RNA treated with tobacco acid pyrophosphatase in order to remove the cap structure present at the 5' ends of messenger RNAs. This reaction leaves a 5' phosphate group at the 5' end of the cap cleaved RNA which can then be ligated to an RNA oligonucleotide using T4 RNA ligase.

This modified RNA preparation is used as a template for first strand cDNA synthesis using a gene specific oligonucleotide. The first strand synthesis reaction is used as a template for PCR amplification of the desired 5' end using a primer specific to the ligated RNA oligonucleotide and a primer specific to the known sequence of the gene of interest. The resultant product is then sequenced and analyzed to confirm that the 5' end sequence belongs to the desired gene.

Example 2: Isolation of Genomic Clones Corresponding to a Polynucleotide

A human genomic P1 library (Genomic Systems, Inc.) is screened by PCR using primers selected for the cDNA sequence corresponding to SEQ ID NO:X., according to the method described in Example 1. (See also, Sambrook.)

Example 3: Tissue Distribution of Polypeptide

Tissue distribution of mRNA expression of polynucleotides of the present invention is determined using protocols for Northern blot analysis, described by, among others, Sambrook et al. For example, a cDNA probe produced by the method described in Example 1 is labeled with P³² using the rediprime™ DNA labeling system (Amersham Life Science), according to manufacturer's instructions. After labeling, the probe is purified using CHROMA SPIN-100™ column (Clontech Laboratories, Inc.), according to manufacturer's protocol number PT1200-1. The purified labeled probe is then used to examine various human tissues for mRNA expression.

Multiple Tissue Northern (MTN) blots containing various human tissues (H) or human immune system tissues (IM) (Clontech) are examined with the labeled probe using ExpressHyb™ hybridization solution (Clontech) according to manufacturer's protocol number PT1190-1. Following hybridization and washing, the blots are mounted and exposed to film at -70 degree C overnight, and the films developed according to standard procedures.

Example 4: Chromosomal Mapping of the Polynucleotides

An oligonucleotide primer set is designed according to the sequence at the 5' end of SEQ ID NO:X. This primer preferably spans about 100 nucleotides. This primer set is then used in a polymerase chain reaction under the following set of conditions : 30 seconds, 95 degree C; 1 minute, 56 degree C; 1 minute, 70 degree C. This cycle is repeated 32 times followed by one 5 minute cycle at 70 degree C. Human, mouse, and hamster DNA is used as template in addition to a somatic cell hybrid panel containing individual chromosomes or chromosome fragments (Bios, Inc). The reactions is analyzed on either 8% polyacrylamide gels or 3.5 % agarose gels. Chromosome mapping is determined by the presence of an approximately 100 bp PCR fragment in the particular somatic cell hybrid.

Example 5: Bacterial Expression of a Polypeptide

A polynucleotide encoding a polypeptide of the present invention is amplified using PCR oligonucleotide primers corresponding to the 5' and 3' ends of the DNA sequence, as outlined in Example 1, to synthesize insertion fragments. The primers used to amplify the cDNA insert should preferably contain restriction sites, such as
5 BamHI and XbaI, at the 5' end of the primers in order to clone the amplified product into the expression vector. For example, BamHI and XbaI correspond to the restriction enzyme sites on the bacterial expression vector pQE-9. (Qiagen, Inc., Chatsworth, CA). This plasmid vector encodes antibiotic resistance (Amp^r), a bacterial origin of replication (ori), an IPTG-regulatable promoter/operator (P/O), a
10 ribosome binding site (RBS), a 6-histidine tag (6-His), and restriction enzyme cloning sites.

The pQE-9 vector is digested with BamHI and XbaI and the amplified fragment is ligated into the pQE-9 vector maintaining the reading frame initiated at the bacterial RBS. The ligation mixture is then used to transform the E. coli strain
15 M15/rep4 (Qiagen, Inc.) which contains multiple copies of the plasmid pREP4, which expresses the lacI repressor and also confers kanamycin resistance (Kan^r). Transformants are identified by their ability to grow on LB plates and ampicillin/kanamycin resistant colonies are selected. Plasmid DNA is isolated and confirmed by restriction analysis.

20 Clones containing the desired constructs are grown overnight (O/N) in liquid culture in LB media supplemented with both Amp (100 ug/ml) and Kan (25 ug/ml). The O/N culture is used to inoculate a large culture at a ratio of 1:100 to 1:250. The cells are grown to an optical density 600 (O.D.⁶⁰⁰) of between 0.4 and 0.6. IPTG (Isopropyl-B-D-thiogalactopyranoside) is then added to a final concentration of 1
25 mM. IPTG induces by inactivating the lacI repressor, clearing the P/O leading to increased gene expression.

Cells are grown for an extra 3 to 4 hours. Cells are then harvested by centrifugation (20 mins at 6000Xg). The cell pellet is solubilized in the chaotropic agent 6 Molar Guanidine HCl by stirring for 3-4 hours at 4 degree C. The cell debris
30 is removed by centrifugation, and the supernatant containing the polypeptide is loaded onto a nickel-nitrilo-tri-acetic acid ("Ni-NTA") affinity resin column (available from QIAGEN, Inc., *supra*). Proteins with a 6 x His tag bind to the Ni-NTA resin with

high affinity and can be purified in a simple one-step procedure (for details see: The QIAexpressionist (1995) QIAGEN, Inc., *supra*).

Briefly, the supernatant is loaded onto the column in 6 M guanidine-HCl, pH 8, the column is first washed with 10 volumes of 6 M guanidine-HCl, pH 8, then
5 washed with 10 volumes of 6 M guanidine-HCl pH 6, and finally the polypeptide is eluted with 6 M guanidine-HCl, pH 5.

The purified protein is then renatured by dialyzing it against phosphate-buffered saline (PBS) or 50 mM Na-acetate, pH 6 buffer plus 200 mM NaCl. Alternatively, the protein can be successfully refolded while immobilized on the Ni-
10 NTA column. The recommended conditions are as follows: renature using a linear 6M-1M urea gradient in 500 mM NaCl, 20% glycerol, 20 mM Tris/HCl pH 7.4, containing protease inhibitors. The renaturation should be performed over a period of 1.5 hours or more. After renaturation the proteins are eluted by the addition of 250 mM imidazole. Imidazole is removed by a final dialyzing step against PBS or 50
15 mM sodium acetate pH 6 buffer plus 200 mM NaCl. The purified protein is stored at 4 degree C or frozen at -80 degree C.

In addition to the above expression vector, the present invention further includes an expression vector comprising phage operator and promoter elements operatively linked to a polynucleotide of the present invention, called pHE4a. (ATCC
20 Accession Number 209645, deposited on February 25, 1998.) This vector contains: 1) a neomycinphosphotransferase gene as a selection marker, 2) an E. coli origin of replication, 3) a T5 phage promoter sequence, 4) two lac operator sequences, 5) a Shine-Delgarno sequence, and 6) the lactose operon repressor gene (*lacIq*). The origin of replication (*oriC*) is derived from pUC19 (LTI, Gaithersburg, MD). The
25 promoter sequence and operator sequences are made synthetically.

DNA can be inserted into the pHEa by restricting the vector with NdeI and XbaI, BamHI, XhoI, or Asp718, running the restricted product on a gel, and isolating the larger fragment (the stuffer fragment should be about 310 base pairs). The DNA insert is generated according to the PCR protocol described in Example 1, using PCR
30 primers having restriction sites for NdeI (5' primer) and XbaI, BamHI, XhoI, or Asp718 (3' primer). The PCR insert is gel purified and restricted with compatible enzymes. The insert and vector are ligated according to standard protocols.

The engineered vector could easily be substituted in the above protocol to express protein in a bacterial system.

Example 6: Purification of a Polypeptide from an Inclusion Body

5 The following alternative method can be used to purify a polypeptide expressed in *E. coli* when it is present in the form of inclusion bodies. Unless otherwise specified, all of the following steps are conducted at 4-10 degree C.

 Upon completion of the production phase of the *E. coli* fermentation, the cell culture is cooled to 4-10 degree C and the cells harvested by continuous
10 centrifugation at 15,000 rpm (Heraeus Sepatech). On the basis of the expected yield of protein per unit weight of cell paste and the amount of purified protein required, an appropriate amount of cell paste, by weight, is suspended in a buffer solution containing 100 mM Tris, 50 mM EDTA, pH 7.4. The cells are dispersed to a homogeneous suspension using a high shear mixer.

15 The cells are then lysed by passing the solution through a microfluidizer (Microfluidics, Corp. or APV Gaulin, Inc.) twice at 4000-6000 psi. The homogenate is then mixed with NaCl solution to a final concentration of 0.5 M NaCl, followed by centrifugation at 7000 xg for 15 min. The resultant pellet is washed again using 0.5M NaCl, 100 mM Tris, 50 mM EDTA, pH 7.4.

20 The resulting washed inclusion bodies are solubilized with 1.5 M guanidine hydrochloride (GuHCl) for 2-4 hours. After 7000 xg centrifugation for 15 min., the pellet is discarded and the polypeptide containing supernatant is incubated at 4 degree C overnight to allow further GuHCl extraction.

 Following high speed centrifugation (30,000 xg) to remove insoluble particles,
25 the GuHCl solubilized protein is refolded by quickly mixing the GuHCl extract with 20 volumes of buffer containing 50 mM sodium, pH 4.5, 150 mM NaCl, 2 mM EDTA by vigorous stirring. The refolded diluted protein solution is kept at 4 degree C without mixing for 12 hours prior to further purification steps.

 To clarify the refolded polypeptide solution, a previously prepared tangential
30 filtration unit equipped with 0.16 um membrane filter with appropriate surface area (e.g., Filtron), equilibrated with 40 mM sodium acetate, pH 6.0 is employed. The filtered sample is loaded onto a cation exchange resin (e.g., Poros HS-50, Perseptive

Biosystems). The column is washed with 40 mM sodium acetate, pH 6.0 and eluted with 250 mM, 500 mM, 1000 mM, and 1500 mM NaCl in the same buffer, in a stepwise manner. The absorbance at 280 nm of the effluent is continuously monitored. Fractions are collected and further analyzed by SDS-PAGE.

5 Fractions containing the polypeptide are then pooled and mixed with 4 volumes of water. The diluted sample is then loaded onto a previously prepared set of tandem columns of strong anion (Poros HQ-50, Perseptive Biosystems) and weak anion (Poros CM-20, Perseptive Biosystems) exchange resins. The columns are equilibrated with 40 mM sodium acetate, pH 6.0. Both columns are washed with 40
10 mM sodium acetate, pH 6.0, 200 mM NaCl. The CM-20 column is then eluted using a 10 column volume linear gradient ranging from 0.2 M NaCl, 50 mM sodium acetate, pH 6.0 to 1.0 M NaCl, 50 mM sodium acetate, pH 6.5. Fractions are collected under constant A_{280} monitoring of the effluent. Fractions containing the polypeptide (determined, for instance, by 16% SDS-PAGE) are then pooled.

15 The resultant polypeptide should exhibit greater than 95% purity after the above refolding and purification steps. No major contaminant bands should be observed from Commassie blue stained 16% SDS-PAGE gel when 5 ug of purified protein is loaded. The purified protein can also be tested for endotoxin/LPS contamination, and typically the LPS content is less than 0.1 ng/ml according to LAL
20 assays.

Example 7: Cloning and Expression of a Polypeptide in a Baculovirus

Expression System

In this example, the plasmid shuttle vector pA2 is used to insert a
25 polynucleotide into a baculovirus to express a polypeptide. This expression vector contains the strong polyhedrin promoter of the *Autographa californica* nuclear polyhedrosis virus (AcMNPV) followed by convenient restriction sites such as BamHI, Xba I and Asp718. The polyadenylation site of the simian virus 40 ("SV40") is used for efficient polyadenylation. For easy selection of recombinant virus, the
30 plasmid contains the beta-galactosidase gene from *E. coli* under control of a weak *Drosophila* promoter in the same orientation, followed by the polyadenylation signal of the polyhedrin gene. The inserted genes are flanked on both sides by viral

sequences for cell-mediated homologous recombination with wild-type viral DNA to generate a viable virus that express the cloned polynucleotide.

Many other baculovirus vectors can be used in place of the vector above, such as pAc373, pVL941, and pAcIM1, as one skilled in the art would readily appreciate, as long as the construct provides appropriately located signals for transcription, translation, secretion and the like, including a signal peptide and an in-frame AUG as required. Such vectors are described, for instance, in Luckow et al., *Virology* 170:31-39 (1989).

Specifically, the cDNA sequence contained in the deposited clone, including the AUG initiation codon and the naturally associated leader sequence identified in Table 1, is amplified using the PCR protocol described in Example 1. If the naturally occurring signal sequence is used to produce the secreted protein, the pA2 vector does not need a second signal peptide. Alternatively, the vector can be modified (pA2 GP) to include a baculovirus leader sequence, using the standard methods described in Summers et al., "A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures," Texas Agricultural Experimental Station Bulletin No. 1555 (1987).

The amplified fragment is isolated from a 1% agarose gel using a commercially available kit ("Geneclean," BIO 101 Inc., La Jolla, Ca.). The fragment then is digested with appropriate restriction enzymes and again purified on a 1% agarose gel.

The plasmid is digested with the corresponding restriction enzymes and optionally, can be dephosphorylated using calf intestinal phosphatase, using routine procedures known in the art. The DNA is then isolated from a 1% agarose gel using a commercially available kit ("Geneclean" BIO 101 Inc., La Jolla, Ca.).

The fragment and the dephosphorylated plasmid are ligated together with T4 DNA ligase. *E. coli* HB101 or other suitable *E. coli* hosts such as XL-1 Blue (Stratagene Cloning Systems, La Jolla, CA) cells are transformed with the ligation mixture and spread on culture plates. Bacteria containing the plasmid are identified by digesting DNA from individual colonies and analyzing the digestion product by gel electrophoresis. The sequence of the cloned fragment is confirmed by DNA sequencing.

Five ug of a plasmid containing the polynucleotide is co-transfected with 1.0 ug of a commercially available linearized baculovirus DNA ("BaculoGold™ baculovirus DNA", Pharmingen, San Diego, CA), using the lipofection method described by Felgner et al., Proc. Natl. Acad. Sci. USA 84:7413-7417 (1987). One ug
5 of BaculoGold™ virus DNA and 5 ug of the plasmid are mixed in a sterile well of a microtiter plate containing 50 ul of serum-free Grace's medium (Life Technologies Inc., Gaithersburg, MD). Afterwards, 10 ul Lipofectin plus 90 ul Grace's medium are added, mixed and incubated for 15 minutes at room temperature. Then the transfection mixture is added drop-wise to Sf9 insect cells (ATCC CRL 1711) seeded
10 in a 35 mm tissue culture plate with 1 ml Grace's medium without serum. The plate is then incubated for 5 hours at 27 degrees C. The transfection solution is then removed from the plate and 1 ml of Grace's insect medium supplemented with 10% fetal calf serum is added. Cultivation is then continued at 27 degrees C for four days.

After four days the supernatant is collected and a plaque assay is performed,
15 as described by Summers and Smith, *supra*. An agarose gel with "Blue Gal" (Life Technologies Inc., Gaithersburg) is used to allow easy identification and isolation of gal-expressing clones, which produce blue-stained plaques. (A detailed description of a "plaque assay" of this type can also be found in the user's guide for insect cell culture and baculovirology distributed by Life Technologies Inc., Gaithersburg, page
20 9-10.) After appropriate incubation, blue stained plaques are picked with the tip of a micropipettor (e.g., Eppendorf). The agar containing the recombinant viruses is then resuspended in a microcentrifuge tube containing 200 ul of Grace's medium and the suspension containing the recombinant baculovirus is used to infect Sf9 cells seeded in 35 mm dishes. Four days later the supernatants of these culture dishes are
25 harvested and then they are stored at 4 degree C.

To verify the expression of the polypeptide, Sf9 cells are grown in Grace's medium supplemented with 10% heat-inactivated FBS. The cells are infected with the recombinant baculovirus containing the polynucleotide at a multiplicity of infection ("MOI") of about 2. If radiolabeled proteins are desired, 6 hours later the
30 medium is removed and is replaced with SF900 II medium minus methionine and cysteine (available from Life Technologies Inc., Rockville, MD). After 42 hours, 5 uCi of ³⁵S-methionine and 5 uCi ³⁵S-cysteine (available from Amersham) are added.

The cells are further incubated for 16 hours and then are harvested by centrifugation. The proteins in the supernatant as well as the intracellular proteins are analyzed by SDS-PAGE followed by autoradiography (if radiolabeled).

Microsequencing of the amino acid sequence of the amino terminus of
5 purified protein may be used to determine the amino terminal sequence of the produced protein.

Example 8: Expression of a Polypeptide in Mammalian Cells

The polypeptide of the present invention can be expressed in a mammalian cell. A typical mammalian expression vector contains a promoter element, which
10 mediates the initiation of transcription of mRNA, a protein coding sequence, and signals required for the termination of transcription and polyadenylation of the transcript. Additional elements include enhancers, Kozak sequences and intervening sequences flanked by donor and acceptor sites for RNA splicing. Highly efficient transcription is achieved with the early and late promoters from SV40, the long
15 terminal repeats (LTRs) from Retroviruses, e.g., RSV, HTLVI, HIVI and the early promoter of the cytomegalovirus (CMV). However, cellular elements can also be used (e.g., the human actin promoter).

Suitable expression vectors for use in practicing the present invention include, for example, vectors such as pSVL and pMSG (Pharmacia, Uppsala, Sweden),
20 pRSVcat (ATCC 37152), pSV2dhfr (ATCC 37146), pBC12MI (ATCC 67109), pCMVSPORT 2.0, and pCMVSPORT 3.0. Mammalian host cells that could be used include, human Hela, 293, H9 and Jurkat cells, mouse NIH3T3 and C127 cells, Cos 1, Cos 7 and CV1, quail QC1-3 cells, mouse L cells and Chinese hamster ovary (CHO) cells.

25 Alternatively, the polypeptide can be expressed in stable cell lines containing the polynucleotide integrated into a chromosome. The co-transfection with a selectable marker such as dhfr, gpt, neomycin, hygromycin allows the identification and isolation of the transfected cells.

The transfected gene can also be amplified to express large amounts of the
30 encoded protein. The DHFR (dihydrofolate reductase) marker is useful in developing cell lines that carry several hundred or even several thousand copies of the gene of

interest. (See, e.g., Alt, F. W., et al., J. Biol. Chem. 253:1357-1370 (1978); Hamlin, J. L. and Ma, C., Biochem. et Biophys. Acta, 1097:107-143 (1990); Page, M. J. and Sydenham, M. A., Biotechnology 9:64-68 (1991).) Another useful selection marker is the enzyme glutamine synthase (GS) (Murphy et al., Biochem J. 227:277-279 (1991); Bebbington et al., Bio/Technology 10:169-175 (1992). Using these markers, the mammalian cells are grown in selective medium and the cells with the highest resistance are selected. These cell lines contain the amplified gene(s) integrated into a chromosome. Chinese hamster ovary (CHO) and NSO cells are often used for the production of proteins.

Derivatives of the plasmid pSV2-dhfr (ATCC Accession No. 37146), the expression vectors pC4 (ATCC Accession No. 209646) and pC6 (ATCC Accession No.209647) contain the strong promoter (LTR) of the Rous Sarcoma Virus (Cullen et al., Molecular and Cellular Biology, 438-447 (March, 1985)) plus a fragment of the CMV-enhancer (Boshart et al., Cell 41:521-530 (1985).) Multiple cloning sites, e.g., with the restriction enzyme cleavage sites BamHI, XbaI and Asp718, facilitate the cloning of the gene of interest. The vectors also contain the 3' intron, the polyadenylation and termination signal of the rat preproinsulin gene, and the mouse DHFR gene under control of the SV40 early promoter.

Specifically, the plasmid pC6, for example, is digested with appropriate restriction enzymes and then dephosphorylated using calf intestinal phosphates by procedures known in the art. The vector is then isolated from a 1% agarose gel.

A polynucleotide of the present invention is amplified according to the protocol outlined in Example 1. If the naturally occurring signal sequence is used to produce the secreted protein, the vector does not need a second signal peptide. Alternatively, if the naturally occurring signal sequence is not used, the vector can be modified to include a heterologous signal sequence. (See, e.g., WO 96/34891.)

The amplified fragment is isolated from a 1% agarose gel using a commercially available kit ("GeneClean," BIO 101 Inc., La Jolla, Ca.). The fragment then is digested with appropriate restriction enzymes and again purified on a 1% agarose gel.

The amplified fragment is then digested with the same restriction enzyme and purified on a 1% agarose gel. The isolated fragment and the dephosphorylated vector

are then ligated with T4 DNA ligase. *E. coli* HB101 or XL-1 Blue cells are then transformed and bacteria are identified that contain the fragment inserted into plasmid pC6 using, for instance, restriction enzyme analysis.

Chinese hamster ovary cells lacking an active DHFR gene is used for transfection. Five μ g of the expression plasmid pC6 a pC4 is cotransfected with 0.5 μ g of the plasmid pSVneo using lipofectin (Felgner et al., *supra*). The plasmid pSV2-neo contains a dominant selectable marker, the *neo* gene from Tn5 encoding an enzyme that confers resistance to a group of antibiotics including G418. The cells are seeded in alpha minus MEM supplemented with 1 mg/ml G418. After 2 days, the cells are trypsinized and seeded in hybridoma cloning plates (Greiner, Germany) in alpha minus MEM supplemented with 10, 25, or 50 ng/ml of methotrexate plus 1 mg/ml G418. After about 10-14 days single clones are trypsinized and then seeded in 6-well petri dishes or 10 ml flasks using different concentrations of methotrexate (50 nM, 100 nM, 200 nM, 400 nM, 800 nM). Clones growing at the highest concentrations of methotrexate are then transferred to new 6-well plates containing even higher concentrations of methotrexate (1 μ M, 2 μ M, 5 μ M, 10 mM, 20 mM). The same procedure is repeated until clones are obtained which grow at a concentration of 100 - 200 μ M. Expression of the desired gene product is analyzed, for instance, by SDS-PAGE and Western blot or by reversed phase HPLC analysis.

Example 9: Protein Fusions

The polypeptides of the present invention are preferably fused to other proteins. These fusion proteins can be used for a variety of applications. For example, fusion of the present polypeptides to His-tag, HA-tag, protein A, IgG domains, and maltose binding protein facilitates purification. (See Example 5; see also EP A 394,827; Traunecker, et al., *Nature* 331:84-86 (1988).) Similarly, fusion to IgG-1, IgG-3, and albumin increases the halflife time in vivo. Nuclear localization signals fused to the polypeptides of the present invention can target the protein to a specific subcellular localization, while covalent heterodimer or homodimers can increase or decrease the activity of a fusion protein. Fusion proteins can also create chimeric molecules having more than one function. Finally, fusion proteins can increase solubility and/or stability of the fused protein compared to the non-fused

protein. All of the types of fusion proteins described above can be made by modifying the following protocol, which outlines the fusion of a polypeptide to an IgG molecule, or the protocol described in Example 5.

5 Briefly, the human Fc portion of the IgG molecule can be PCR amplified, using primers that span the 5' and 3' ends of the sequence described below. These primers also should have convenient restriction enzyme sites that will facilitate cloning into an expression vector, preferably a mammalian expression vector.

10 For example, if pC4 (Accession No. 209646) is used, the human Fc portion can be ligated into the BamHI cloning site. Note that the 3' BamHI site should be destroyed. Next, the vector containing the human Fc portion is re-restricted with BamHI, linearizing the vector, and a polynucleotide of the present invention, isolated by the PCR protocol described in Example 1, is ligated into this BamHI site. Note that the polynucleotide is cloned without a stop codon, otherwise a fusion protein will not be produced.

15 If the naturally occurring signal sequence is used to produce the secreted protein, pC4 does not need a second signal peptide. Alternatively, if the naturally occurring signal sequence is not used, the vector can be modified to include a heterologous signal sequence. (See, e.g., WO 96/34891.)

20 Human IgG Fc region:

GGGATCCGGAGCCCAAATCTTCTGACAAAACCTCACACATGCCCACCGTGC
CCAGCACCTGAATTCGAGGGTGCACCGTCAGTCTTCCTCTTCCCCCAAAA
CCCAAGGACACCCTCATGATCTCCCGGACTCCTGAGGTCACATGCGTGGT
GGTGGACGTAAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGG
25 ACGGCGTGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTA
CAACAGCACGTACCGTGTGGTCAGCGTCCTCACCGTCCTGCACCAGGACT
GGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCCA
ACCCCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAAC
CACAGGTGTACACCCTGCCCCCATCCCGGGATGAGCTGACCAAGAACCAG
30 GTCAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCAAGCGACATCGCCGT
GGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACACTACAAGACCACGCCT
CCCGTGCTGGACTCCGACGGCTCCTTCTTCCTCTACAGCAAGCTCACCGTG

GACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCA
TGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGG
GTAAATGAGTGCACGGCCGCGACTCTAGAGGAT (SEQ ID NO:1)

5 **Example 10: Production of an Antibody from a Polypeptide**

The antibodies of the present invention can be prepared by a variety of methods. (See, Current Protocols, Chapter 2.) As one example of such methods, cells expressing a polypeptide of the present invention is administered to an animal to induce the production of sera containing polyclonal antibodies. In a preferred
10 method, a preparation of the secreted protein is prepared and purified to render it substantially free of natural contaminants. Such a preparation is then introduced into an animal in order to produce polyclonal antisera of greater specific activity.

In the most preferred method, the antibodies of the present invention are monoclonal antibodies (or protein binding fragments thereof). Such monoclonal
15 antibodies can be prepared using hybridoma technology. (Köhler et al., Nature 256:495 (1975); Köhler et al., Eur. J. Immunol. 6:511 (1976); Köhler et al., Eur. J. Immunol. 6:292 (1976); Hammerling et al., in: Monoclonal Antibodies and T-Cell Hybridomas, Elsevier, N.Y., pp. 563-681 (1981).) In general, such procedures involve immunizing an animal (preferably a mouse) with polypeptide or, more
20 preferably, with a secreted polypeptide-expressing cell. Such cells may be cultured in any suitable tissue culture medium; however, it is preferable to culture cells in Earle's modified Eagle's medium supplemented with 10% fetal bovine serum (inactivated at about 56 degrees C), and supplemented with about 10 g/l of nonessential amino acids, about 1,000 U/ml of penicillin, and about 100 ug/ml of streptomycin.

The splenocytes of such mice are extracted and fused with a suitable myeloma cell line. Any suitable myeloma cell line may be employed in accordance with the present invention; however, it is preferable to employ the parent myeloma cell line (SP2O), available from the ATCC. After fusion, the resulting hybridoma cells are selectively maintained in HAT medium, and then cloned by limiting dilution as
25 described by Wands et al. (Gastroenterology 80:225-232 (1981).) The hybridoma cells obtained through such a selection are then assayed to identify clones which secrete antibodies capable of binding the polypeptide.
30

Alternatively, additional antibodies capable of binding to the polypeptide can be produced in a two-step procedure using anti-idiotypic antibodies. Such a method makes use of the fact that antibodies are themselves antigens, and therefore, it is possible to obtain an antibody which binds to a second antibody. In accordance with this method, protein specific antibodies are used to immunize an animal, preferably a mouse. The splenocytes of such an animal are then used to produce hybridoma cells, and the hybridoma cells are screened to identify clones which produce an antibody whose ability to bind to the protein-specific antibody can be blocked by the polypeptide. Such antibodies comprise anti-idiotypic antibodies to the protein-specific antibody and can be used to immunize an animal to induce formation of further protein-specific antibodies.

It will be appreciated that Fab and F(ab')₂ and other fragments of the antibodies of the present invention may be used according to the methods disclosed herein. Such fragments are typically produced by proteolytic cleavage, using enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab')₂ fragments). Alternatively, secreted protein-binding fragments can be produced through the application of recombinant DNA technology or through synthetic chemistry.

For in vivo use of antibodies in humans, it may be preferable to use "humanized" chimeric monoclonal antibodies. Such antibodies can be produced using genetic constructs derived from hybridoma cells producing the monoclonal antibodies described above. Methods for producing chimeric antibodies are known in the art. (See, for review, Morrison, Science 229:1202 (1985); Oi et al., BioTechniques 4:214 (1986); Cabilly et al., U.S. Patent No. 4,816,567; Taniguchi et al., EP 171496; Morrison et al., EP 173494; Neuberger et al., WO 8601533; Robinson et al., WO 8702671; Boulianne et al., Nature 312:643 (1984); Neuberger et al., Nature 314:268 (1985).)

Example 11: Production Of Secreted Protein For High-Throughput Screening

Assays

The following protocol produces a supernatant containing a polypeptide to be tested. This supernatant can then be used in the Screening Assays described herein.

First, dilute Poly-D-Lysine (644 587 Boehringer-Mannheim) stock solution (1mg/ml in PBS) 1:20 in PBS (w/o calcium or magnesium 17-516F Biowhittaker) for a working solution of 50ug/ml. Add 200 ul of this solution to each well (24 well plates) and incubate at RT for 20 minutes. Be sure to distribute the solution over each well (note: a 12-channel pipetter may be used with tips on every other channel).

Aspirate off the Poly-D-Lysine solution and rinse with 1ml PBS (Phosphate Buffered Saline). The PBS should remain in the well until just prior to plating the cells and plates may be poly-lysine coated in advance for up to two weeks.

Plate 293T cells (do not carry cells past P+20) at 2×10^5 cells/well in .5ml DMEM(Dulbecco's Modified Eagle Medium)(with 4.5 G/L glucose and L-glutamine (12-604F Biowhittaker))/10% heat inactivated FBS(14-503F Biowhittaker)/1x Penstrep(17-602E Biowhittaker). Let the cells grow overnight.

The next day, mix together in a sterile solution basin: 300 ul Lipofectamine (18324-012 Gibco/BRL) and 5ml Optimem I (31985070 Gibco/BRL)/96-well plate.

With a small volume multi-channel pipetter, aliquot approximately 2ug of an expression vector containing a polynucleotide insert, produced by the methods described in Examples 8 or 9, into an appropriately labeled 96-well round bottom plate. With a multi-channel pipetter, add 50ul of the Lipofectamine/Optimem I mixture to each well. Pipette up and down gently to mix. Incubate at RT 15-45 minutes. After about 20 minutes, use a multi-channel pipetter to add 150ul Optimem I to each well. As a control, one plate of vector DNA lacking an insert should be transfected with each set of transfections.

Preferably, the transfection should be performed by tag-teaming the following tasks. By tag-teaming, hands on time is cut in half, and the cells do not spend too much time on PBS. First, person A aspirates off the media from four 24-well plates of cells, and then person B rinses each well with .5-1ml PBS. Person A then aspirates off PBS rinse, and person B, using a 12-channel pipetter with tips on every other channel, adds the 200ul of DNA/Lipofectamine/Optimem I complex to the odd wells first, then to the even wells, to each row on the 24-well plates. Incubate at 37 degrees C for 6 hours.

While cells are incubating, prepare appropriate media, either 1%BSA in DMEM with 1x penstrep, or CHO-5 media (116.6 mg/L of CaCl₂ (anhyd); 0.00130

mg/L $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$; 0.050 mg/L of $\text{Fe}(\text{NO}_3)_3 \cdot 9\text{H}_2\text{O}$; 0.417 mg/L of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$;
 311.80 mg/L of KCl; 28.64 mg/L of MgCl_2 ; 48.84 mg/L of MgSO_4 ; 6995.50 mg/L of
 NaCl; 2400.0 mg/L of NaHCO_3 ; 62.50 mg/L of $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$; 71.02 mg/L of
 Na_2HPO_4 ; .4320 mg/L of $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$; .002 mg/L of Arachidonic Acid ; 1.022 mg/L
 5 of Cholesterol; .070 mg/L of DL-alpha-Tocopherol-Acetate; 0.0520 mg/L of Linoleic
 Acid; 0.010 mg/L of Linolenic Acid; 0.010 mg/L of Myristic Acid; 0.010 mg/L of
 Oleic Acid; 0.010 mg/L of Palmitric Acid; 0.010 mg/L of Palmitic Acid; 100 mg/L of
 Pluronic F-68; 0.010 mg/L of Stearic Acid; 2.20 mg/L of Tween 80; 4551 mg/L of D-
 Glucose; 130.85 mg/ml of L- Alanine; 147.50 mg/ml of L-Arginine-HCL; 7.50 mg/ml
 10 of L-Asparagine- H_2O ; 6.65 mg/ml of L-Aspartic Acid; 29.56 mg/ml of L-Cystine-
 2HCL- H_2O ; 31.29 mg/ml of L-Cystine-2HCL; 7.35 mg/ml of L-Glutamic Acid; 365.0
 mg/ml of L-Glutamine; 18.75 mg/ml of Glycine; 52.48 mg/ml of L-Histidine-HCL-
 H_2O ; 106.97 mg/ml of L-Isoleucine; 111.45 mg/ml of L-Leucine; 163.75 mg/ml of L-
 Lysine HCL; 32.34 mg/ml of L-Methionine; 68.48 mg/ml of L-Phenylalanine; 40.0
 15 mg/ml of L-Proline; 26.25 mg/ml of L-Serine; 101.05 mg/ml of L-Threonine; 19.22
 mg/ml of L-Tryptophan; 91.79 mg/ml of L-Tyrosine-2Na- $2\text{H}_2\text{O}$; 99.65 mg/ml of L-
 Valine; 0.0035 mg/L of Biotin; 3.24 mg/L of D-Ca Pantothenate; 11.78 mg/L of
 Choline Chloride; 4.65 mg/L of Folic Acid; 15.60 mg/L of i-Inositol; 3.02 mg/L of
 Niacinamide; 3.00 mg/L of Pyridoxal HCL; 0.031 mg/L of Pyridoxine HCL; 0.319
 20 mg/L of Riboflavin; 3.17 mg/L of Thiamine HCL; 0.365 mg/L of Thymidine; and
 0:680 mg/L of Vitamin B₁₂; 25 mM of HEPES Buffer; 2.39 mg/L of Na
 Hypoxanthine; 0.105 mg/L of Lipoic Acid; 0.081 mg/L of Sodium Putrescine-2HCL;
 55.0 mg/L of Sodium Pyruvate; 0.0067 mg/L of Sodium Selenite; 20uM of
 Ethanolamine; 0.122 mg/L of Ferric Citrate; 41.70 mg/L of Methyl-B-Cyclodextrin
 25 complexed with Linoleic Acid; 33.33 mg/L of Methyl-B-Cyclodextrin complexed
 with Oleic Acid; and 10 mg/L of Methyl-B-Cyclodextrin complexed with Retinal)
 with 2mm glutamine and 1x penstrep. (BSA (81-068-3 Bayer) 100gm dissolved in 1L
 DMEM for a 10% BSA stock solution). Filter the media and collect 50 ul for
 endotoxin assay in 15ml polystyrene conical.

30 The transfection reaction is terminated, preferably by tag-teaming, at the end
 of the incubation period. Person A aspirates off the transfection media, while person

B adds 1.5ml appropriate media to each well. Incubate at 37 degrees C for 45 or 72 hours depending on the media used: 1%BSA for 45 hours or CHO-5 for 72 hours.

On day four, using a 300ul multichannel pipetter, aliquot 600ul in one 1ml deep well plate and the remaining supernatant into a 2ml deep well. The supernatants
5 from each well can then be used in the assays described in Examples 13-20.

It is specifically understood that when activity is obtained in any of the assays described below using a supernatant, the activity originates from either the polypeptide directly (e.g., as a secreted protein) or by the polypeptide inducing expression of other proteins, which are then secreted into the supernatant. Thus, the
10 invention further provides a method of identifying the protein in the supernatant characterized by an activity in a particular assay.

Example 12: Construction of GAS Reporter Construct

One signal transduction pathway involved in the differentiation and
15 proliferation of cells is called the Jaks-STATs pathway. Activated proteins in the Jaks-STATs pathway bind to gamma activation site "GAS" elements or interferon-sensitive responsive element ("ISRE"), located in the promoter of many genes. The binding of a protein to these elements alter the expression of the associated gene.

GAS and ISRE elements are recognized by a class of transcription factors
20 called Signal Transducers and Activators of Transcription, or "STATs." There are six members of the STATs family. Stat1 and Stat3 are present in many cell types, as is Stat2 (as response to IFN-alpha is widespread). Stat4 is more restricted and is not in many cell types though it has been found in T helper class I, cells after treatment with IL-12. Stat5 was originally called mammary growth factor, but has been found at
25 higher concentrations in other cells including myeloid cells. It can be activated in tissue culture cells by many cytokines.

The STATs are activated to translocate from the cytoplasm to the nucleus upon tyrosine phosphorylation by a set of kinases known as the Janus Kinase ("Jaks") family. Jaks represent a distinct family of soluble tyrosine kinases and include Tyk2,
30 Jak1, Jak2, and Jak3. These kinases display significant sequence similarity and are generally catalytically inactive in resting cells.

The Jaks are activated by a wide range of receptors summarized in the Table below. (Adapted from review by Schidler and Darnell, *Ann. Rev. Biochem.* 64:621-51 (1995).) A cytokine receptor family, capable of activating Jaks, is divided into two groups: (a) Class 1 includes receptors for IL-2, IL-3, IL-4, IL-6, IL-7, IL-9, IL-11, IL-12, IL-15, Epo, PRL, GH, G-CSF, GM-CSF, LIF, CNTF, and thrombopoietin; and (b) Class 2 includes IFN-a, IFN-g, and IL-10. The Class 1 receptors share a conserved cysteine motif (a set of four conserved cysteines and one tryptophan) and a WSXWS motif (a membrane proximal region encoding Trp-Ser-Xxx-Trp-Ser (SEQ ID NO:2)).

Thus, on binding of a ligand to a receptor, Jaks are activated, which in turn activate STATs, which then translocate and bind to GAS elements. This entire process is encompassed in the Jaks-STATs signal transduction pathway.

Therefore, activation of the Jaks-STATs pathway, reflected by the binding of the GAS or the ISRE element, can be used to indicate proteins involved in the proliferation and differentiation of cells. For example, growth factors and cytokines are known to activate the Jaks-STATs pathway. (See Table below.) Thus, by using GAS elements linked to reporter molecules, activators of the Jaks-STATs pathway can be identified.

| | <u>Ligand</u> | <u>tyk2</u> | <u>JAKs</u> <u>Jak1</u> | <u>Jak2</u> | <u>Jak3</u> | <u>STATS</u> | <u>GAS(elements) or ISRE</u> |
|----|----------------------------------|-------------|----------------------------|-------------|-------------|--------------|------------------------------|
| | <u>IFN family</u> | | | | | | |
| 5 | IFN-a/B | + | + | - | - | 1,2,3 | ISRE |
| | IFN-g | | + | + | - | 1 | GAS (IRF1>Lys6>IFP) |
| | Il-10 | + | ? | ? | - | 1,3 | |
| | <u>gp130 family</u> | | | | | | |
| 10 | IL-6 (Pleiotrophic) | + | + | + | ? | 1,3 | GAS (IRF1>Lys6>IFP) |
| | Il-11(Pleiotrophic) | ? | + | ? | ? | 1,3 | |
| | OnM(Pleiotrophic) | ? | + | + | ? | 1,3 | |
| | LIF(Pleiotrophic) | ? | + | + | ? | 1,3 | |
| | CNTF(Pleiotrophic) | -/+ | + | + | ? | 1,3 | |
| 15 | G-CSF(Pleiotrophic) | ? | + | ? | ? | 1,3 | |
| | IL-12(Pleiotrophic) | + | - | + | + | 1,3 | |
| | <u>g-C family</u> | | | | | | |
| | IL-2 (lymphocytes) | - | + | - | + | 1,3,5 | GAS |
| 20 | IL-4 (lymph/myeloid) | - | + | - | + | 6 | GAS (IRF1 = IFP >>Ly6)(IgH) |
| | IL-7 (lymphocytes) | - | + | - | + | 5 | GAS |
| | IL-9 (lymphocytes) | - | + | - | + | 5 | GAS |
| | IL-13 (lymphocyte) | - | + | ? | ? | 6 | GAS |
| | IL-15 | ? | + | ? | + | 5 | GAS |
| 25 | <u>gp140 family</u> | | | | | | |
| | IL-3 (myeloid) | - | - | + | - | 5 | GAS (IRF1>IFP>>Ly6) |
| | IL-5 (myeloid) | - | - | + | - | 5 | GAS |
| | GM-CSF (myeloid) | - | - | + | - | 5 | GAS |
| 30 | <u>Growth hormone family</u> | | | | | | |
| | GH | ? | - | + | - | 5 | |
| | PRL | ? | +/- | + | - | 1,3,5 | |
| | EPO | ? | - | + | - | 5 | GAS(B-CAS>IRF1=IFP>>Ly6) |
| 35 | <u>Receptor Tyrosine Kinases</u> | | | | | | |
| | EGF | ? | + | + | - | 1,3 | GAS (IRF1) |
| | PDGF | ? | + | + | - | 1,3 | |
| | CSF-1 | ? | + | + | - | 1,3 | GAS (not IRF1) |
| 40 | | | | | | | |

To construct a synthetic GAS containing promoter element, which is used in the Biological Assays described in Examples 13-14, a PCR based strategy is employed to generate a GAS-SV40 promoter sequence. The 5' primer contains four tandem copies of the GAS binding site found in the IRF1 promoter and previously demonstrated to bind STATs upon induction with a range of cytokines (Rothman et al., Immunity 1:457-468 (1994).), although other GAS or ISRE elements can be used instead. The 5' primer also contains 18bp of sequence complementary to the SV40 early promoter sequence and is flanked with an XhoI site. The sequence of the 5' primer is:

5':GCGCCTCGAGATTTCCTCGAAATCTAGATTTCCTCGAAATGATTTCCTCGAAATGATTTCCTCGAAATATCTGCCATCTCAATTAG:3' (SEQ ID NO:3)

The downstream primer is complementary to the SV40 promoter and is flanked with a Hind III site: 5':GCGGCAAGCTTTTGGCAAAGCCTAGGC:3' (SEQ ID NO:4)

PCR amplification is performed using the SV40 promoter template present in the B-gal:promoter plasmid obtained from Clontech. The resulting PCR fragment is digested with XhoI/Hind III and subcloned into BLSK2-. (Stratagene.) Sequencing with forward and reverse primers confirms that the insert contains the following sequence:

5':CTCGAGATTTCCTCGAAATCTAGATTTCCTCGAAATGATTTCCTCGAAATGATTTCCTCGAAATATCTGCCATCTCAATTAGTCAGCAACCATAGTCCCGCCCTAACTCCGCCCATCCCGCCCCTAACTCCGCCCAGTTCCGCCCATTCTCCGCCCCATGGCTGACTAATTTTTTTTATTTATGCAGAGGCCGAGGCCGCCTCGGCCTCTGAGCTATTCCAGAAGTAGTGAGGAGGCTTTTTTGGAGGCCTAGGCTTTTGGAAAAAGCTT:3' (SEQ ID NO:5)

With this GAS promoter element linked to the SV40 promoter, a GAS:SEAP2 reporter construct is next engineered. Here, the reporter molecule is a secreted alkaline phosphatase, or "SEAP." Clearly, however, any reporter molecule can be instead of SEAP, in this or in any of the other Examples. Well known reporter molecules that can be used instead of SEAP include chloramphenicol acetyltransferase (CAT), luciferase, alkaline phosphatase, B-galactosidase, green fluorescent protein (GFP), or any protein detectable by an antibody.

The above sequence confirmed synthetic GAS-SV40 promoter element is subcloned into the pSEAP-Promoter vector obtained from Clontech using HindIII and XhoI, effectively replacing the SV40 promoter with the amplified GAS:SV40 promoter element, to create the GAS-SEAP vector. However, this vector does not contain a neomycin resistance gene, and therefore, is not preferred for mammalian expression systems.

Thus, in order to generate mammalian stable cell lines expressing the GAS-SEAP reporter, the GAS-SEAP cassette is removed from the GAS-SEAP vector using SalI and NotI, and inserted into a backbone vector containing the neomycin resistance gene, such as pGFP-1 (Clontech), using these restriction sites in the multiple cloning site, to create the GAS-SEAP/Neo vector. Once this vector is transfected into mammalian cells, this vector can then be used as a reporter molecule for GAS binding as described in Examples 13-14.

Other constructs can be made using the above description and replacing GAS with a different promoter sequence. For example, construction of reporter molecules containing NFK-B and EGR promoter sequences are described in Examples 15 and 16. However, many other promoters can be substituted using the protocols described in these Examples. For instance, SRE, IL-2, NFAT, or Osteocalcin promoters can be substituted, alone or in combination (e.g., GAS/NF-KB/EGR, GAS/NF-KB, IL-2/NFAT, or NF-KB/GAS). Similarly, other cell lines can be used to test reporter construct activity, such as HELA (epithelial), HUVEC (endothelial), Reh (B-cell), Saos-2 (osteoblast), HUVAC (aortic), or Cardiomyocyte.

Example 13: High-Throughput Screening Assay for T-cell Activity.

The following protocol is used to assess T-cell activity by identifying factors, and determining whether supernate containing a polypeptide of the invention proliferates and/or differentiates T-cells. T-cell activity is assessed using the GAS/SEAP/Neo construct produced in Example 12. Thus, factors that increase SEAP activity indicate the ability to activate the Jaks-STATS signal transduction pathway. The T-cell used in this assay is Jurkat T-cells (ATCC Accession No. TIB-152), although Molt-3 cells (ATCC Accession No. CRL-1552) and Molt-4 cells (ATCC Accession No. CRL-1582) cells can also be used.

Jurkat T-cells are lymphoblastic CD4+ Th1 helper cells. In order to generate stable cell lines, approximately 2 million Jurkat cells are transfected with the GAS-SEAP/neo vector using DMRIE-C (Life Technologies)(transfection procedure described below). The transfected cells are seeded to a density of approximately
5 20,000 cells per well and transfectants resistant to 1 mg/ml gentamicin selected. Resistant colonies are expanded and then tested for their response to increasing concentrations of interferon gamma. The dose response of a selected clone is demonstrated.

Specifically, the following protocol will yield sufficient cells for 75 wells
10 containing 200 ul of cells. Thus, it is either scaled up, or performed in multiple to generate sufficient cells for multiple 96 well plates. Jurkat cells are maintained in RPMI + 10% serum with 1%Pen-Strep. Combine 2.5 mls of OPTI-MEM (Life Technologies) with 10 ug of plasmid DNA in a T25 flask. Add 2.5 ml OPTI-MEM containing 50 ul of DMRIE-C and incubate at room temperature for 15-45 mins.

15 During the incubation period, count cell concentration, spin down the required number of cells (10^7 per transfection), and resuspend in OPTI-MEM to a final concentration of 10^7 cells/ml. Then add 1ml of 1×10^7 cells in OPTI-MEM to T25 flask and incubate at 37 degrees C for 6 hrs. After the incubation, add 10 ml of RPMI + 15% serum.

20 The Jurkat:GAS-SEAP stable reporter lines are maintained in RPMI + 10% serum, 1 mg/ml Gentamicin, and 1% Pen-Strep. These cells are treated with supernatants containing polypeptides of the invention and/or induced polypeptides of the invention as produced by the protocol described in Example 11.

On the day of treatment with the supernatant, the cells should be washed and
25 resuspended in fresh RPMI + 10% serum to a density of 500,000 cells per ml. The exact number of cells required will depend on the number of supernatants being screened. For one 96 well plate, approximately 10 million cells (for 10 plates, 100 million cells) are required.

Transfer the cells to a triangular reservoir boat, in order to dispense the cells
30 into a 96 well dish, using a 12 channel pipette. Using a 12 channel pipette, transfer 200 ul of cells into each well (therefore adding 100, 000 cells per well).

After all the plates have been seeded, 50 ul of the supernatants are transferred directly from the 96 well plate containing the supernatants into each well using a 12 channel pipette. In addition, a dose of exogenous interferon gamma (0.1, 1.0, 10 ng) is added to wells H9, H10, and H11 to serve as additional positive controls for the
5 assay.

The 96 well dishes containing Jurkat cells treated with supernatants are placed in an incubator for 48 hrs (note: this time is variable between 48-72 hrs). 35 ul samples from each well are then transferred to an opaque 96 well plate using a 12 channel pipette. The opaque plates should be covered (using sellophene covers) and
10 stored at -20 degrees C until SEAP assays are performed according to Example 17. The plates containing the remaining treated cells are placed at 4 degrees C and serve as a source of material for repeating the assay on a specific well if desired.

As a positive control, 100 Unit/ml interferon gamma can be used which is known to activate Jurkat T cells. Over 30 fold induction is typically observed in the
15 positive control wells.

The above protocol may be used in the generation of both transient, as well as, stable transfected cells, which would be apparent to those of skill in the art.

Example 14: High-Throughput Screening Assay Identifying Myeloid Activity

20 The following protocol is used to assess myeloid activity by determining whether polypeptides of the invention proliferates and/or differentiates myeloid cells. Myeloid cell activity is assessed using the GAS/SEAP/Neo construct produced in Example 12. Thus, factors that increase SEAP activity indicate the ability to activate the Jaks-STATS signal transduction pathway. The myeloid cell used in this assay is
25 U937, a pre-monocyte cell line, although TF-1, HL60, or KG1 can be used.

To transiently transfect U937 cells with the GAS/SEAP/Neo construct produced in Example 12, a DEAE-Dextran method (Kharbanda et. al., 1994, Cell Growth & Differentiation, 5:259-265) is used. First, harvest 2×10^7 U937 cells and wash with PBS. The U937 cells are usually grown in RPMI 1640 medium containing
30 10% heat-inactivated fetal bovine serum (FBS) supplemented with 100 units/ml penicillin and 100 mg/ml streptomycin.

Next, suspend the cells in 1 ml of 20 mM Tris-HCl (pH 7.4) buffer containing 0.5 mg/ml DEAE-Dextran, 8 ug GAS-SEAP2 plasmid DNA, 140 mM NaCl, 5 mM KCl, 375 uM Na₂HPO₄·7H₂O, 1 mM MgCl₂, and 675 uM CaCl₂. Incubate at 37 degrees C for 45 min.

- 5 Wash the cells with RPMI 1640 medium containing 10% FBS and then resuspend in 10 ml complete medium and incubate at 37 degrees C for 36 hr.

The GAS-SEAP/U937 stable cells are obtained by growing the cells in 400 ug/ml G418. The G418-free medium is used for routine growth but every one to two months, the cells should be re-grown in 400 ug/ml G418 for couple of passages.

- 10 These cells are tested by harvesting 1×10^8 cells (this is enough for ten 96-well plates assay) and wash with PBS. Suspend the cells in 200 ml above described growth medium, with a final density of 5×10^5 cells/ml. Plate 200 ul cells per well in the 96-well plate (or 1×10^5 cells/well).

Add 50 ul of the supernatant prepared by the protocol described in Example

- 15 11. Incubate at 37 degrees C for 48 to 72 hr. As a positive control, 100 Unit/ml interferon gamma can be used which is known to activate U937 cells. Over 30 fold induction is typically observed in the positive control wells. SEAP assay the supernatant according to the protocol described in Example 17.

20 **Example 15: High-Throughput Screening Assay Identifying Neuronal Activity.**

- When cells undergo differentiation and proliferation, a group of genes are activated through many different signal transduction pathways. One of these genes, EGR1 (early growth response gene 1), is induced in various tissues and cell types upon activation. The promoter of EGR1 is responsible for such induction. Using the
- 25 EGR1 promoter linked to reporter molecules, activation of cells can be assessed.

- Particularly, the following protocol is used to assess neuronal activity in PC12 cell lines. PC12 cells (rat pheochromocytoma cells) are known to proliferate and/or differentiate by activation with a number of mitogens, such as TPA (tetradecanoyl phorbol acetate), NGF (nerve growth factor), and EGF (epidermal growth factor).
- 30 The EGR1 gene expression is activated during this treatment. Thus, by stably transfecting PC12 cells with a construct containing an EGR promoter linked to SEAP reporter, activation of PC12 cells can be assessed.

The EGR/SEAP reporter construct can be assembled by the following protocol. The EGR-1 promoter sequence (-633 to +1)(Sakamoto K et al., Oncogene 6:867-871 (1991)) can be PCR amplified from human genomic DNA using the following primers:

- 5 5' GCGCTCGAGGGATGACAGCGATAGAACCCCGG -3' (SEQ ID NO:6)
 5' GCGAAGCTTCGCGACTCCCCGGATCCGCCTC-3' (SEQ ID NO:7)

Using the GAS:SEAP/Neo vector produced in Example 12, EGR1 amplified product can then be inserted into this vector. Linearize the GAS:SEAP/Neo vector using restriction enzymes XhoI/HindIII, removing the GAS/SV40 stuffer. Restrict the
10 EGR1 amplified product with these same enzymes. Ligate the vector and the EGR1 promoter.

To prepare 96 well-plates for cell culture, two mls of a coating solution (1:30 dilution of collagen type I (Upstate Biotech Inc. Cat#08-115) in 30% ethanol (filter sterilized)) is added per one 10 cm plate or 50 ml per well of the 96-well plate, and
15 allowed to air dry for 2 hr.

PC12 cells are routinely grown in RPMI-1640 medium (Bio Whittaker) containing 10% horse serum (JRH BIOSCIENCES, Cat. # 12449-78P), 5% heat-inactivated fetal bovine serum (FBS) supplemented with 100 units/ml penicillin and 100 ug/ml streptomycin on a precoated 10 cm tissue culture dish. One to four split is
20 done every three to four days. Cells are removed from the plates by scraping and resuspended with pipetting up and down for more than 15 times.

Transfect the EGR/SEAP/Neo construct into PC12 using the Lipofectamine protocol described in Example 11. EGR-SEAP/PC12 stable cells are obtained by growing the cells in 300 ug/ml G418. The G418-free medium is used for routine
25 growth but every one to two months, the cells should be re-grown in 300 ug/ml G418 for couple of passages.

To assay for neuronal activity, a 10 cm plate with cells around 70 to 80% confluent is screened by removing the old medium. Wash the cells once with PBS (Phosphate buffered saline). Then starve the cells in low serum medium (RPMI-1640
30 containing 1% horse serum and 0.5% FBS with antibiotics) overnight.

The next morning, remove the medium and wash the cells with PBS. Scrape off the cells from the plate, suspend the cells well in 2 ml low serum medium. Count

the cell number and add more low serum medium to reach final cell density as 5×10^5 cells/ml.

Add 200 μ l of the cell suspension to each well of 96-well plate (equivalent to 1×10^5 cells/well). Add 50 μ l supernatant produced by Example 11, 37°C for 48 to 72 hr. As a positive control, a growth factor known to activate PC12 cells through EGR can be used, such as 50 ng/ μ l of Neuronal Growth Factor (NGF). Over fifty-fold induction of SEAP is typically seen in the positive control wells. SEAP assay the supernatant according to Example 17.

10 **Example 16: High-Throughput Screening Assay for T-cell Activity**

NF-KB (Nuclear Factor KB) is a transcription factor activated by a wide variety of agents including the inflammatory cytokines IL-1 and TNF, CD30 and CD40, lymphotoxin-alpha and lymphotoxin-beta, by exposure to LPS or thrombin, and by expression of certain viral gene products. As a transcription factor, NF-KB regulates the expression of genes involved in immune cell activation, control of apoptosis (NF- KB appears to shield cells from apoptosis), B and T-cell development, anti-viral and antimicrobial responses, and multiple stress responses.

In non-stimulated conditions, NF- KB is retained in the cytoplasm with I-KB (Inhibitor KB). However, upon stimulation, I- KB is phosphorylated and degraded, causing NF- KB to shuttle to the nucleus, thereby activating transcription of target genes. Target genes activated by NF- KB include IL-2, IL-6, GM-CSF, ICAM-1 and class 1 MHC.

Due to its central role and ability to respond to a range of stimuli, reporter constructs utilizing the NF-KB promoter element are used to screen the supernatants produced in Example 11. Activators or inhibitors of NF-KB would be useful in treating diseases. For example, inhibitors of NF-KB could be used to treat those diseases related to the acute or chronic activation of NF-KB, such as rheumatoid arthritis.

To construct a vector containing the NF-KB promoter element, a PCR based strategy is employed. The upstream primer contains four tandem copies of the NF-KB binding site (GGGGACTTTC) (SEQ ID NO:8), 18 bp of sequence

complementary to the 5' end of the SV40 early promoter sequence, and is flanked with an XhoI site:

5':GCGGCCTCGAGGGGACTTTCCCGGGGACTTTCCGGGGACTTTCCGGGAC
TTTCCATCCTGCCATCTCAATTAG:3' (SEQ ID NO:9)

- 5 The downstream primer is complementary to the 3' end of the SV40 promoter and is flanked with a Hind III site:

5':GCGGCAAGCTTTTTGCAAAGCCTAGGC:3' (SEQ ID NO:4)

- PCR amplification is performed using the SV40 promoter template present in the pB-gal:promoter plasmid obtained from Clontech. The resulting PCR fragment is
10 digested with XhoI and Hind III and subcloned into BLSK2-. (Stratagene)
Sequencing with the T7 and T3 primers confirms the insert contains the following sequence:

- 5':CTCGAGGGGACTTTCCCGGGGACTTTCCGGGGACTTTCCGGGACTTTCC
15 ATCTGCCATCTCAATTAGTCAGCAACCATAGTCCCGCCCCTAACTCCGCCC
ATCCCGCCCCTAACTCCGCCCAGTTCCGCCCATTCTCCGCCCCATGGCTGA
CTAATTTTTTTTATTTATGCAGAGGCCGAGGCCGCCTCGGCCTCTGAGCTA
TTCCAGAAGTAGTGAGGAGGCTTTTTTGGAGGCCTAGGCTTTTGCAAAAA
GCTT:3' (SEQ ID NO:10)

20

Next, replace the SV40 minimal promoter element present in the pSEAP2-promoter plasmid (Clontech) with this NF-KB/SV40 fragment using XhoI and HindIII. However, this vector does not contain a neomycin resistance gene, and therefore, is not preferred for mammalian expression systems.

- 25 In order to generate stable mammalian cell lines, the NF-KB/SV40/SEAP cassette is removed from the above NF-KB/SEAP vector using restriction enzymes SalI and NotI, and inserted into a vector containing neomycin resistance. Particularly, the NF-KB/SV40/SEAP cassette was inserted into pGFP-1 (Clontech), replacing the GFP gene, after restricting pGFP-1 with SalI and NotI.

- 30 Once NF-KB/SV40/SEAP/Neo vector is created, stable Jurkat T-cells are created and maintained according to the protocol described in Example 13. Similarly, the method for assaying supernatants with these stable Jurkat T-cells is also described

in Example 13. As a positive control, exogenous TNF alpha (0.1, 1, 10 ng) is added to wells H9, H10, and H11, with a 5-10 fold activation typically observed.

Example 17: Assay for SEAP Activity

5 As a reporter molecule for the assays described in Examples 13-16, SEAP activity is assayed using the Tropix Phospho-light Kit (Cat. BP-400) according to the following general procedure. The Tropix Phospho-light Kit supplies the Dilution, Assay, and Reaction Buffers used below.

10 Prime a dispenser with the 2.5x Dilution Buffer and dispense 15 ul of 2.5x dilution buffer into Optiplates containing 35 ul of a supernatant. Seal the plates with a plastic sealer and incubate at 65 degree C for 30 min. Separate the Optiplates to avoid uneven heating.

15 Cool the samples to room temperature for 15 minutes. Empty the dispenser and prime with the Assay Buffer. Add 50 ml Assay Buffer and incubate at room temperature 5 min. Empty the dispenser and prime with the Reaction Buffer (see the table below). Add 50 ul Reaction Buffer and incubate at room temperature for 20 minutes. Since the intensity of the chemiluminescent signal is time dependent, and it takes about 10 minutes to read 5 plates on luminometer, one should treat 5 plates at each time and start the second set 10 minutes later.

20 Read the relative light unit in the luminometer. Set H12 as blank, and print the results. An increase in chemiluminescence indicates reporter activity.

Reaction Buffer Formulation:

| # of plates | Rxn buffer diluent (ml) | CSPD (ml) |
|-------------|-------------------------|-----------|
| 10 | 60 | 3 |
| 11 | 65 | 3.25 |
| 12 | 70 | 3.5 |
| 13 | 75 | 3.75 |
| 14 | 80 | 4 |
| 15 | 85 | 4.25 |
| 16 | 90 | 4.5 |
| 17 | 95 | 4.75 |
| 18 | 100 | 5 |
| 19 | 105 | 5.25 |
| 20 | 110 | 5.5 |
| 21 | 115 | 5.75 |
| 22 | 120 | 6 |
| 23 | 125 | 6.25 |
| 24 | 130 | 6.5 |

| | | |
|----|-----|-------|
| 25 | 135 | 6.75 |
| 26 | 140 | 7 |
| 27 | 145 | 7.25 |
| 28 | 150 | 7.5 |
| 29 | 155 | 7.75 |
| 30 | 160 | 8 |
| 31 | 165 | 8.25 |
| 32 | 170 | 8.5 |
| 33 | 175 | 8.75 |
| 34 | 180 | 9 |
| 35 | 185 | 9.25 |
| 36 | 190 | 9.5 |
| 37 | 195 | 9.75 |
| 38 | 200 | 10 |
| 39 | 205 | 10.25 |
| 40 | 210 | 10.5 |
| 41 | 215 | 10.75 |
| 42 | 220 | 11 |
| 43 | 225 | 11.25 |
| 44 | 230 | 11.5 |
| 45 | 235 | 11.75 |
| 46 | 240 | 12 |
| 47 | 245 | 12.25 |
| 48 | 250 | 12.5 |
| 49 | 255 | 12.75 |
| 50 | 260 | 13 |

Example 18: High-Throughput Screening Assay Identifying Changes in Small Molecule Concentration and Membrane Permeability

Binding of a ligand to a receptor is known to alter intracellular levels of small molecules, such as calcium, potassium, sodium, and pH, as well as alter membrane potential. These alterations can be measured in an assay to identify supernatants which bind to receptors of a particular cell. Although the following protocol describes an assay for calcium, this protocol can easily be modified to detect changes in potassium, sodium, pH, membrane potential, or any other small molecule which is detectable by a fluorescent probe.

The following assay uses Fluorometric Imaging Plate Reader ("FLIPR") to measure changes in fluorescent molecules (Molecular Probes) that bind small molecules. Clearly, any fluorescent molecule detecting a small molecule can be used instead of the calcium fluorescent molecule, fluo-4 (Molecular Probes, Inc.; catalog no. F-14202), used here.

For adherent cells, seed the cells at 10,000 -20,000 cells/well in a Co-star black 96-well plate with clear bottom. The plate is incubated in a CO₂ incubator for

20 hours. The adherent cells are washed two times in Biotek washer with 200 ul of HBSS (Hank's Balanced Salt Solution) leaving 100 ul of buffer after the final wash.

A stock solution of 1 mg/ml fluo-4 is made in 10% pluronic acid DMSO. To load the cells with fluo-4, 50 ul of 12 ug/ml fluo-4 is added to each well. The plate is incubated at 37 degrees C in a CO₂ incubator for 60 min. The plate is washed four times in the Biotek washer with HBSS leaving 100 ul of buffer.

For non-adherent cells, the cells are spun down from culture media. Cells are re-suspended to 2-5x10⁶ cells/ml with HBSS in a 50-ml conical tube. 4 ul of 1 mg/ml fluo-4 solution in 10% pluronic acid DMSO is added to each ml of cell suspension. The tube is then placed in a 37 degrees C water bath for 30-60 min. The cells are washed twice with HBSS, resuspended to 1x10⁶ cells/ml, and dispensed into a microplate, 100 ul/well. The plate is centrifuged at 1000 rpm for 5 min. The plate is then washed once in Denley CellWash with 200 ul, followed by an aspiration step to 100 ul final volume.

For a non-cell based assay, each well contains a fluorescent molecule, such as fluo-4. The supernatant is added to the well, and a change in fluorescence is detected.

To measure the fluorescence of intracellular calcium, the FLIPR is set for the following parameters: (1) System gain is 300-800 mW; (2) Exposure time is 0.4 second; (3) Camera F/stop is F/2; (4) Excitation is 488 nm; (5) Emission is 530 nm; and (6) Sample addition is 50 ul. Increased emission at 530 nm indicates an extracellular signaling event which has resulted in an increase in the intracellular Ca⁺⁺ concentration.

Example 19: High-Throughput Screening Assay Identifying Tyrosine Kinase Activity

The Protein Tyrosine Kinases (PTK) represent a diverse group of transmembrane and cytoplasmic kinases. Within the Receptor Protein Tyrosine Kinase (RPTK) group are receptors for a range of mitogenic and metabolic growth factors including the PDGF, FGF, EGF, NGF, HGF and Insulin receptor subfamilies. In addition there are a large family of RPTKs for which the corresponding ligand is

unknown. Ligands for RPTKs include mainly secreted small proteins, but also membrane-bound and extracellular matrix proteins.

Activation of RPTK by ligands involves ligand-mediated receptor dimerization, resulting in transphosphorylation of the receptor subunits and activation of the cytoplasmic tyrosine kinases. The cytoplasmic tyrosine kinases include receptor associated tyrosine kinases of the src-family (e.g., src, yes, lck, lyn, fyn) and non-receptor linked and cytosolic protein tyrosine kinases, such as the Jak family, members of which mediate signal transduction triggered by the cytokine superfamily of receptors (e.g., the Interleukins, Interferons, GM-CSF, and Leptin).

Because of the wide range of known factors capable of stimulating tyrosine kinase activity, the identification of novel human secreted proteins capable of activating tyrosine kinase signal transduction pathways are of interest. Therefore, the following protocol is designed to identify those novel human secreted proteins capable of activating the tyrosine kinase signal transduction pathways.

Seed target cells (e.g., primary keratinocytes) at a density of approximately 25,000 cells per well in a 96 well Loprodyne Silent Screen Plates purchased from Nalge Nunc (Naperville, IL). The plates are sterilized with two 30 minute rinses with 100% ethanol, rinsed with water and dried overnight. Some plates are coated for 2 hr with 100 ml of cell culture grade type I collagen (50 mg/ml), gelatin (2%) or polylysine (50 mg/ml), all of which can be purchased from Sigma Chemicals (St. Louis, MO) or 10% Matrigel purchased from Becton Dickinson (Bedford, MA), or calf serum, rinsed with PBS and stored at 4 degree C. Cell growth on these plates is assayed by seeding 5,000 cells/well in growth medium and indirect quantitation of cell number through use of alamarBlue as described by the manufacturer Alamar Biosciences, Inc. (Sacramento, CA) after 48 hr. Falcon plate covers #3071 from Becton Dickinson (Bedford, MA) are used to cover the Loprodyne Silent Screen Plates. Falcon Microtest III cell culture plates can also be used in some proliferation experiments.

To prepare extracts, A431 cells are seeded onto the nylon membranes of Loprodyne plates (20,000/200ml/well) and cultured overnight in complete medium. Cells are quiesced by incubation in serum-free basal medium for 24 hr. After 5-20 minutes treatment with EGF (60ng/ml) or 50 ul of the supernatant produced in

Example 11, the medium was removed and 100 ml of extraction buffer ((20 mM HEPES pH 7.5, 0.15 M NaCl, 1% Triton X-100, 0.1% SDS, 2 mM Na₃VO₄, 2 mM Na₄P₂O₇ and a cocktail of protease inhibitors (# 1836170) obtained from Boehringer Mannheim (Indianapolis, IN) is added to each well and the plate is
5 shaken on a rotating shaker for 5 minutes at 4 degrees C. The plate is then placed in a vacuum transfer manifold and the extract filtered through the 0.45 mm membrane bottoms of each well using house vacuum. Extracts are collected in a 96-well catch/assay plate in the bottom of the vacuum manifold and immediately placed on ice. To obtain extracts clarified by centrifugation, the content of each well, after
10 detergent solubilization for 5 minutes, is removed and centrifuged for 15 minutes at 4 degrees C at 16,000 x g.

Test the filtered extracts for levels of tyrosine kinase activity. Although many methods of detecting tyrosine kinase activity are known, one method is described here.

15 Generally, the tyrosine kinase activity of a supernatant is evaluated by determining its ability to phosphorylate a tyrosine residue on a specific substrate (a biotinylated peptide). Biotinylated peptides that can be used for this purpose include PSK1 (corresponding to amino acids 6-20 of the cell division kinase cdc2-p34) and PSK2 (corresponding to amino acids 1-17 of gastrin). Both peptides are substrates for
20 a range of tyrosine kinases and are available from Boehringer Mannheim.

The tyrosine kinase reaction is set up by adding the following components in order. First, add 10ul of 5uM Biotinylated Peptide, then 10ul ATP/Mg₂⁺ (5mM ATP/50mM MgCl₂), then 10ul of 5x Assay Buffer (40mM imidazole hydrochloride, pH7.3, 40 mM beta-glycerophosphate, 1mM EGTA, 100mM MgCl₂, 5 mM MnCl₂,
25 0.5 mg/ml BSA), then 5ul of Sodium Vanadate(1mM), and then 5ul of water. Mix the components gently and preincubate the reaction mix at 30 degrees C for 2 min. Initial the reaction by adding 10ul of the control enzyme or the filtered supernatant.

The tyrosine kinase assay reaction is then terminated by adding 10 ul of 120mM EDTA and place the reactions on ice.

30 Tyrosine kinase activity is determined by transferring 50 ul aliquot of reaction mixture to a microtiter plate (MTP) module and incubating at 37 degrees C for 20

min. This allows the streptavidin coated 96 well plate to associate with the biotinylated peptide. Wash the MTP module with 300ul/well of PBS four times. Next add 75 ul of anti-phosphotyrosine antibody conjugated to horse radish peroxidase(anti-P-Tyr-POD(0.5u/ml)) to each well and incubate at 37 degrees C for one hour. Wash the well as above.

Next add 100ul of peroxidase substrate solution (Boehringer Mannheim) and incubate at room temperature for at least 5 mins (up to 30 min). Measure the absorbance of the sample at 405 nm by using ELISA reader. The level of bound peroxidase activity is quantitated using an ELISA reader and reflects the level of tyrosine kinase activity.

Example 20: High-Throughput Screening Assay Identifying Phosphorylation Activity

As a potential alternative and/or compliment to the assay of protein tyrosine kinase activity described in Example 19, an assay which detects activation (phosphorylation) of major intracellular signal transduction intermediates can also be used. For example, as described below one particular assay can detect tyrosine phosphorylation of the Erk-1 and Erk-2 kinases. However, phosphorylation of other molecules, such as Raf, JNK, p38 MAP, Map kinase kinase (MEK), MEK kinase, Src, Muscle specific kinase (MuSK), IRAK, Tec, and Janus, as well as any other phosphoserine, phosphotyrosine, or phosphothreonine molecule, can be detected by substituting these molecules for Erk-1 or Erk-2 in the following assay.

Specifically, assay plates are made by coating the wells of a 96-well ELISA plate with 0.1ml of protein G (1ug/ml) for 2 hr at room temp, (RT). The plates are then rinsed with PBS and blocked with 3% BSA/PBS for 1 hr at RT. The protein G plates are then treated with 2 commercial monoclonal antibodies (100ng/well) against Erk-1 and Erk-2 (1 hr at RT) (Santa Cruz Biotechnology). (To detect other molecules, this step can easily be modified by substituting a monoclonal antibody detecting any of the above described molecules.) After 3-5 rinses with PBS, the plates are stored at 4 degrees C until use.

A431 cells are seeded at 20,000/well in a 96-well Loprodyne filterplate and

cultured overnight in growth medium. The cells are then starved for 48 hr in basal medium (DMEM) and then treated with EGF (6ng/well) or 50 ul of the supernatants obtained in Example 11 for 5-20 minutes. The cells are then solubilized and extracts filtered directly into the assay plate.

- 5 After incubation with the extract for 1 hr at RT, the wells are again rinsed. As a positive control, a commercial preparation of MAP kinase (10ng/well) is used in place of A431 extract. Plates are then treated with a commercial polyclonal (rabbit) antibody (1ug/ml) which specifically recognizes the phosphorylated epitope of the Erk-1 and Erk-2 kinases (1 hr at RT). This antibody is biotinylated by standard
10 procedures. The bound polyclonal antibody is then quantitated by successive incubations with Europium-streptavidin and Europium fluorescence enhancing reagent in the Wallac DELFIA instrument (time-resolved fluorescence). An increased fluorescent signal over background indicates a phosphorylation.

15 **Example 21: Method of Determining Alterations in a Gene Corresponding to a Polynucleotide**

- RNA isolated from entire families or individual patients presenting with a phenotype of interest (such as a disease) is be isolated. cDNA is then generated from these RNA samples using protocols known in the art. (See, Sambrook.) The cDNA
20 is then used as a template for PCR, employing primers surrounding regions of interest in SEQ ID NO:X. Suggested PCR conditions consist of 35 cycles at 95 degrees C for 30 seconds; 60-120 seconds at 52-58 degrees C; and 60-120 seconds at 70 degrees C, using buffer solutions described in Sidransky et al., Science 252:706 (1991).

- PCR products are then sequenced using primers labeled at their 5' end with T4
25 polynucleotide kinase, employing SequiTherm Polymerase. (Epicentre Technologies). The intron-exon borders of selected exons is also determined and genomic PCR products analyzed to confirm the results. PCR products harboring suspected mutations is then cloned and sequenced to validate the results of the direct sequencing.

- 30 PCR products is cloned into T-tailed vectors as described in Holton et al., Nucleic Acids Research, 19:1156 (1991) and sequenced with T7 polymerase (United

States Biochemical). Affected individuals are identified by mutations not present in unaffected individuals.

Genomic rearrangements are also observed as a method of determining alterations in a gene corresponding to a polynucleotide. Genomic clones isolated according to Example 2 are nick-translated with digoxigenindeoxy-uridine 5'-triphosphate (Boehringer Mannheim), and FISH performed as described in Johnson et al., Methods Cell Biol. 35:73-99 (1991). Hybridization with the labeled probe is carried out using a vast excess of human cot-1 DNA for specific hybridization to the corresponding genomic locus.

Chromosomes are counterstained with 4,6-diamino-2-phenylidole and propidium iodide, producing a combination of C- and R-bands. Aligned images for precise mapping are obtained using a triple-band filter set (Chroma Technology, Brattleboro, VT) in combination with a cooled charge-coupled device camera (Photometrics, Tucson, AZ) and variable excitation wavelength filters. (Johnson et al., Genet. Anal. Tech. Appl., 8:75 (1991).) Image collection, analysis and chromosomal fractional length measurements are performed using the ISee Graphical Program System. (Inovision Corporation, Durham, NC.) Chromosome alterations of the genomic region hybridized by the probe are identified as insertions, deletions, and translocations. These alterations are used as a diagnostic marker for an associated disease.

Example 22: Method of Detecting Abnormal Levels of a Polypeptide in a Biological Sample

A polypeptide of the present invention can be detected in a biological sample, and if an increased or decreased level of the polypeptide is detected, this polypeptide is a marker for a particular phenotype. Methods of detection are numerous, and thus, it is understood that one skilled in the art can modify the following assay to fit their particular needs.

For example, antibody-sandwich ELISAs are used to detect polypeptides in a sample, preferably a biological sample. Wells of a microtiter plate are coated with specific antibodies, at a final concentration of 0.2 to 10 ug/ml. The antibodies are either monoclonal or polyclonal and are produced by the method described in

Example 10. The wells are blocked so that non-specific binding of the polypeptide to the well is reduced.

The coated wells are then incubated for > 2 hours at RT with a sample containing the polypeptide. Preferably, serial dilutions of the sample should be used to validate results. The plates are then washed three times with deionized or distilled water to remove unbounded polypeptide.

Next, 50 ul of specific antibody-alkaline phosphatase conjugate, at a concentration of 25-400 ng, is added and incubated for 2 hours at room temperature. The plates are again washed three times with deionized or distilled water to remove unbounded conjugate.

Add 75 ul of 4-methylumbelliferyl phosphate (MUP) or p-nitrophenyl phosphate (NPP) substrate solution to each well and incubate 1 hour at room temperature. Measure the reaction by a microtiter plate reader. Prepare a standard curve, using serial dilutions of a control sample, and plot polypeptide concentration on the X-axis (log scale) and fluorescence or absorbance of the Y-axis (linear scale). Interpolate the concentration of the polypeptide in the sample using the standard curve.

Example 23: Formulation

The invention also provides methods of treatment and/or prevention diseases, disorders, and/or conditions (such as, for example, any one or more of the diseases or disorders disclosed herein) by administration to a subject of an effective amount of a Therapeutic. By therapeutic is meant a polynucleotides or polypeptides of the invention (including fragments and variants), agonists or antagonists thereof, and/or antibodies thereto, in combination with a pharmaceutically acceptable carrier type (e.g., a sterile carrier).

The Therapeutic will be formulated and dosed in a fashion consistent with good medical practice, taking into account the clinical condition of the individual patient (especially the side effects of treatment with the Therapeutic alone), the site of delivery, the method of administration, the scheduling of administration, and other factors known to practitioners. The "effective amount" for purposes herein is thus determined by such considerations.

As a general proposition, the total pharmaceutically effective amount of the Therapeutic administered parenterally per dose will be in the range of about 1 ug/kg/day to 10 mg/kg/day of patient body weight, although, as noted above, this will be subject to therapeutic discretion. More preferably, this dose is at least 0.01 mg/kg/day, and most preferably for humans between about 0.01 and 1 mg/kg/day for the hormone. If given continuously, the Therapeutic is typically administered at a dose rate of about 1 ug/kg/hour to about 50 ug/kg/hour, either by 1-4 injections per day or by continuous subcutaneous infusions, for example, using a mini-pump. An intravenous bag solution may also be employed. The length of treatment needed to observe changes and the interval following treatment for responses to occur appears to vary depending on the desired effect.

Therapeutics can be are administered orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, gels, drops or transdermal patch), buccally, or as an oral or nasal spray. "Pharmaceutically acceptable carrier" refers to a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any. The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

Therapeutics of the invention are also suitably administered by sustained-release systems. Suitable examples of sustained-release Therapeutics are administered orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, gels, drops or transdermal patch), buccally, or as an oral or nasal spray. "Pharmaceutically acceptable carrier" refers to a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

Therapeutics of the invention are also suitably administered by sustained-release systems. Suitable examples of sustained-release Therapeutics include suitable polymeric materials (such as, for example, semi-permeable polymer matrices in the form of shaped articles, e.g., films, or microcapsules), suitable hydrophobic materials

(for example as an emulsion in an acceptable oil) or ion exchange resins, and sparingly soluble derivatives (such as, for example, a sparingly soluble salt).

Sustained-release matrices include polylactides (U.S. Pat. No. 3,773,919, EP 58,481), copolymers of L-glutamic acid and gamma-ethyl-L-glutamate (Sidman et al.,
5 Biopolymers 22:547-556 (1983)), poly (2- hydroxyethyl methacrylate) (Langer et al., J. Biomed. Mater. Res. 15:167-277 (1981), and Langer, Chem. Tech. 12:98-105 (1982)), ethylene vinyl acetate (Langer et al., Id.) or poly-D- (-)-3-hydroxybutyric acid (EP 133,988).

Sustained-release Therapeutics also include liposomally entrapped
10 Therapeutics of the invention (see generally, Langer, *Science* 249:1527-1533 (1990); Treat et al., in *Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 317 -327 and 353-365 (1989)). Liposomes containing the Therapeutic are prepared by methods known per se: DE 3,218,121; Epstein et al., Proc. Natl. Acad. Sci. (USA) 82:3688-3692 (1985); Hwang
15 et al., Proc. Natl. Acad. Sci.(USA) 77:4030-4034 (1980); EP 52,322; EP 36,676; EP 88,046; EP 143,949; EP 142,641; Japanese Pat. Appl. 83-118008; U.S. Pat. Nos. 4,485,045 and 4,544,545; and EP 102,324. Ordinarily, the liposomes are of the small (about 200-800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol. percent cholesterol, the selected proportion being adjusted for the
20 optimal Therapeutic.

— In yet an additional embodiment, the Therapeutics of the invention are delivered by way of a pump (see Langer, *supra*; Sefton, CRC Crit. Ref. Biomed. Eng. 14:201 (1987); Buchwald et al., *Surgery* 88:507 (1980); Saudek et al., *N. Engl. J. Med.* 321:574 (1989)).

25 Other controlled release systems are discussed in the review by Langer (*Science* 249:1527-1533 (1990)).

For parenteral administration, in one embodiment, the Therapeutic is formulated generally by mixing it at the desired degree of purity, in a unit dosage injectable form (solution, suspension, or emulsion), with a pharmaceutically
30 acceptable carrier, i.e., one that is non-toxic to recipients at the dosages and concentrations employed and is compatible with other ingredients of the formulation.

For example, the formulation preferably does not include oxidizing agents and other compounds that are known to be deleterious to the Therapeutic.

Generally, the formulations are prepared by contacting the Therapeutic uniformly and intimately with liquid carriers or finely divided solid carriers or both.

5 Then, if necessary, the product is shaped into the desired formulation. Preferably the carrier is a parenteral carrier, more preferably a solution that is isotonic with the blood of the recipient. Examples of such carrier vehicles include water, saline, Ringer's solution, and dextrose solution. Non-aqueous vehicles such as fixed oils and ethyl oleate are also useful herein, as well as liposomes.

10 The carrier suitably contains minor amounts of additives such as substances that enhance isotonicity and chemical stability. Such materials are non-toxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, succinate, acetic acid, and other organic acids or their salts; antioxidants such as ascorbic acid; low molecular weight (less than about ten
15 residues) polypeptides, e.g., polyarginine or tripeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids, such as glycine, glutamic acid, aspartic acid, or arginine; monosaccharides, disaccharides, and other carbohydrates including cellulose or its derivatives, glucose, manose, or dextrans; chelating agents such as EDTA; sugar
20 alcohols such as mannitol or sorbitol; counterions such as sodium; and/or nonionic surfactants such as polysorbates, poloxamers, or PEG.

The Therapeutic is typically formulated in such vehicles at a concentration of about 0.1 mg/ml to 100 mg/ml, preferably 1-10 mg/ml, at a pH of about 3 to 8. It will be understood that the use of certain of the foregoing excipients, carriers, or
25 stabilizers will result in the formation of polypeptide salts.

Any pharmaceutical used for therapeutic administration can be sterile. Sterility is readily accomplished by filtration through sterile filtration membranes (e.g., 0.2 micron membranes). Therapeutics generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a
30 stopper pierceable by a hypodermic injection needle.

Therapeutics ordinarily will be stored in unit or multi-dose containers, for example, sealed ampoules or vials, as an aqueous solution or as a lyophilized

formulation for reconstitution. As an example of a lyophilized formulation, 10-ml vials are filled with 5 ml of sterile-filtered 1% (w/v) aqueous Therapeutic solution, and the resulting mixture is lyophilized. The infusion solution is prepared by reconstituting the lyophilized Therapeutic using bacteriostatic Water-for-Injection.

5 The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the Therapeutics of the invention. Associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use
10 or sale for human administration. In addition, the Therapeutics may be employed in conjunction with other therapeutic compounds.

 The Therapeutics of the invention may be administered alone or in combination with adjuvants. Adjuvants that may be administered with the Therapeutics of the invention include, but are not limited to, alum, alum plus
15 deoxycholate (ImmunoAg), MTP-PE (Biocine Corp.), QS21 (Genentech, Inc.), BCG, and MPL. In a specific embodiment, Therapeutics of the invention are administered in combination with alum. In another specific embodiment, Therapeutics of the invention are administered in combination with QS-21. Further adjuvants that may be administered with the Therapeutics of the invention include, but are not limited to,
20 Monophosphoryl lipid immunomodulator, AduVax 100a, QS-21, QS-18, CRL1005, Aluminum salts, MF-59, and Virosomal adjuvant technology. Vaccines that may be administered with the Therapeutics of the invention include, but are not limited to, vaccines directed toward protection against MMR (measles, mumps, rubella), polio, varicella, tetanus/diphtheria, hepatitis A, hepatitis B, haemophilus influenzae B,
25 whooping cough, pneumonia, influenza, Lyme's Disease, rotavirus, cholera, yellow fever, Japanese encephalitis, poliomyelitis, rabies, typhoid fever, and pertussis. Combinations may be administered either concomitantly, e.g., as an admixture, separately but simultaneously or concurrently; or sequentially. This includes presentations in which the combined agents are administered together as a therapeutic
30 mixture, and also procedures in which the combined agents are administered separately but simultaneously, e.g., as through separate intravenous lines into the same individual. Administration "in combination" further includes the separate

administration of one of the compounds or agents given first, followed by the second.

The Therapeutics of the invention may be administered alone or in combination with other therapeutic agents. Therapeutic agents that may be administered in combination with the Therapeutics of the invention, include but not
5 limited to, other members of the TNF family, chemotherapeutic agents, antibiotics, steroidal and non-steroidal anti-inflammatories, conventional immunotherapeutic agents, cytokines and/or growth factors. Combinations may be administered either concomitantly, e.g., as an admixture, separately but simultaneously or concurrently; or sequentially. This includes presentations in which the combined agents are
10 administered together as a therapeutic mixture, and also procedures in which the combined agents are administered separately but simultaneously, e.g., as through separate intravenous lines into the same individual. Administration "in combination" further includes the separate administration of one of the compounds or agents given first, followed by the second.

15 In one embodiment, the Therapeutics of the invention are administered in combination with members of the TNF family. TNF, TNF-related or TNF-like molecules that may be administered with the Therapeutics of the invention include, but are not limited to, soluble forms of TNF-alpha, lymphotoxin-alpha (LT-alpha, also known as TNF-beta), LT-beta (found in complex heterotrimer LT-alpha2-beta),
20 OPGL, FasL, CD27L, CD30L, CD40L, 4-1BBL, DcR3, OX40L, TNF-gamma (International Publication No. WO 96/14328), AIM-I (International Publication No. WO 97/33899), endokine-alpha (International Publication No. WO 98/07880), TR6 (International Publication No. WO 98/30694), OPG, and neutrokin-alpha (International Publication No. WO 98/18921, OX40, and nerve growth factor (NGF),
25 and soluble forms of Fas, CD30, CD27, CD40 and 4-1BB, TR2 (International Publication No. WO 96/34095), DR3 (International Publication No. WO 97/33904), DR4 (International Publication No. WO 98/32856), TR5 (International Publication No. WO 98/30693), TR6 (International Publication No. WO 98/30694), TR7 (International Publication No. WO 98/41629), TRANK, TR9 (International
30 Publication No. WO 98/56892), TR10 (International Publication No. WO 98/54202), 312C2 (International Publication No. WO 98/06842), and TR12, and soluble forms CD154, CD70, and CD153.

In certain embodiments, Therapeutics of the invention are administered in combination with antiretroviral agents, nucleoside reverse transcriptase inhibitors, non-nucleoside reverse transcriptase inhibitors, and/or protease inhibitors. Nucleoside reverse transcriptase inhibitors that may be administered in combination with the Therapeutics of the invention, include, but are not limited to, RETROVIR™ (zidovudine/AZT), VIDEX™ (didanosine/ddI), HIVID™ (zalcitabine/ddC), ZERIT™ (stavudine/d4T), EPIVIR™ (lamivudine/3TC), and COMBIVIR™ (zidovudine/lamivudine). Non-nucleoside reverse transcriptase inhibitors that may be administered in combination with the Therapeutics of the invention, include, but are not limited to, VIRAMUNE™ (nevirapine), RESCRIPTOR™ (delavirdine), and SUSTIVA™ (efavirenz). Protease inhibitors that may be administered in combination with the Therapeutics of the invention, include, but are not limited to, CRIVAN™ (indinavir), NORVIR™ (ritonavir), INVIRASE™ (saquinavir), and VIRACEPT™ (nelfinavir). In a specific embodiment, antiretroviral agents, nucleoside reverse transcriptase inhibitors, non-nucleoside reverse transcriptase inhibitors, and/or protease inhibitors may be used in any combination with Therapeutics of the invention to treat AIDS and/or to prevent or treat HIV infection.

In other embodiments, Therapeutics of the invention may be administered in combination with anti-opportunistic infection agents. Anti-opportunistic agents that may be administered in combination with the Therapeutics of the invention, include, but are not limited to, TRIMETHOPRIM-SULFAMETHOXAZOLE™, DAPSONE™, PENTAMIDINE™, ATOVAQUONE™, ISONIAZID™, RIFAMPIN™, PYRAZINAMIDE™, ETHAMBUTOL™, RIFABUTIN™, CLARITHROMYCIN™, AZITHROMYCIN™, GANCICLOVIR™, FOSCARNET™, CIDOFOVIR™, FLUCONAZOLE™, ITRACONAZOLE™, KETOCONAZOLE™, ACYCLOVIR™, FAMCICOLVIR™, PYRIMETHAMINE™, LEUCOVORIN™, NEUPOGEN™ (filgrastim/G-CSF), and LEUKINE™ (sargramostim/GM-CSF). In a specific embodiment, Therapeutics of the invention are used in any combination with TRIMETHOPRIM-SULFAMETHOXAZOLE™, DAPSONE™, PENTAMIDINE™, and/or ATOVAQUONE™ to prophylactically treat or prevent an opportunistic *Pneumocystis carinii* pneumonia infection. In

another specific embodiment, Therapeutics of the invention are used in any combination with ISONIAZID™, RIFAMPIN™, PYRAZINAMIDE™, and/or ETHAMBUTOL™ to prophylactically treat or prevent an opportunistic *Mycobacterium avium* complex infection. In another specific embodiment,

5 Therapeutics of the invention are used in any combination with RIFABUTIN™, CLARITHROMYCIN™, and/or AZITHROMYCIN™ to prophylactically treat or prevent an opportunistic *Mycobacterium tuberculosis* infection. In another specific embodiment, Therapeutics of the invention are used in any combination with GANCICLOVIR™, FOSCARNET™, and/or CIDOFOVIR™ to prophylactically treat

10 or prevent an opportunistic cytomegalovirus infection. In another specific embodiment, Therapeutics of the invention are used in any combination with FLUCONAZOLE™, ITRACONAZOLE™, and/or KETOCONAZOLE™ to prophylactically treat or prevent an opportunistic fungal infection. In another specific embodiment, Therapeutics of the invention are used in any combination with

15 ACYCLOVIR™ and/or FAMCICOLVIR™ to prophylactically treat or prevent an opportunistic herpes simplex virus type I and/or type II infection. In another specific embodiment, Therapeutics of the invention are used in any combination with PYRIMETHAMINE™ and/or LEUCOVORIN™ to prophylactically treat or prevent an opportunistic *Toxoplasma gondii* infection. In another specific embodiment,

20 Therapeutics of the invention are used in any combination with LEUCOVORIN™ and/or NEUPOGEN™ to prophylactically treat or prevent an opportunistic bacterial infection.

In a further embodiment, the Therapeutics of the invention are administered in combination with an antiviral agent. Antiviral agents that may be administered

25 with the Therapeutics of the invention include, but are not limited to, acyclovir, ribavirin, amantadine, and remantidine.

In a further embodiment, the Therapeutics of the invention are administered in combination with an antibiotic agent. Antibiotic agents that may be administered with the Therapeutics of the invention include, but are not limited to, amoxicillin,

30 beta-lactamases, aminoglycosides, beta-lactam (glycopeptide), beta-lactamases, Clindamycin, chloramphenicol, cephalosporins, ciprofloxacin, ciprofloxacin,

erythromycin, fluoroquinolones, macrolides, metronidazole, penicillins, quinolones, rifampin, streptomycin, sulfonamide, tetracyclines, trimethoprim, trimethoprim-sulfamthoxazole, and vancomycin.

Conventional nonspecific immunosuppressive agents, that may be administered in combination with the Therapeutics of the invention include, but are not limited to, steroids, cyclosporine, cyclosporine analogs, cyclophosphamide methylprednisone, prednisone, azathioprine, FK-506, 15-deoxyspergualin, and other immunosuppressive agents that act by suppressing the function of responding T cells.

In specific embodiments, Therapeutics of the invention are administered in combination with immunosuppressants. Immunosuppressants preparations that may be administered with the Therapeutics of the invention include, but are not limited to, ORTHOCLONE™ (OKT3), SANDIMMUNE™/NEORAL™/SANGDYA™ (cyclosporin), PROGRAF™ (tacrolimus), CELLCEPT™ (mycophenolate), Azathioprine, glucocorticosteroids, and RAPAMUNE™ (sirolimus). In a specific embodiment, immunosuppressants may be used to prevent rejection of organ or bone marrow transplantation.

In an additional embodiment, Therapeutics of the invention are administered alone or in combination with one or more intravenous immune globulin preparations. Intravenous immune globulin preparations that may be administered with the Therapeutics of the invention include, but not limited to, GAMMAR™, IVEEGAM™, SANDOGLOBULIN™, GAMMAGARD S/D™, and GAMIMUNE™. In a specific embodiment, Therapeutics of the invention are administered in combination with intravenous immune globulin preparations in transplantation therapy (e.g., bone marrow transplant).

In an additional embodiment, the Therapeutics of the invention are administered alone or in combination with an anti-inflammatory agent. Anti-inflammatory agents that may be administered with the Therapeutics of the invention include, but are not limited to, glucocorticoids and the nonsteroidal anti-inflammatories, aminoarylcarboxylic acid derivatives, arylacetic acid derivatives, arylbutyric acid derivatives, arylcarboxylic acids, arylpropionic acid derivatives, pyrazoles, pyrazolones, salicylic acid derivatives, thiazinecarboxamides, e-

acetamidocaproic acid, S-adenosylmethionine, 3-amino-4-hydroxybutyric acid, amixetrine, bendazac, benzydamine, bucolome, difenpiramide, ditazol, emorfazone, guaiazulene, nabumetone, nimesulide, orgotein, oxaceprol, paranyline, perisoxal, pifoxime, proquazone, proxazole, and tenidap.

5 In another embodiment, compositions of the invention are administered in combination with a chemotherapeutic agent. Chemotherapeutic agents that may be administered with the Therapeutics of the invention include, but are not limited to, antibiotic derivatives (e.g., doxorubicin, bleomycin, daunorubicin, and dactinomycin); antiestrogens (e.g., tamoxifen); antimetabolites (e.g., fluorouracil, 5-
10 FU, methotrexate, floxuridine, interferon alpha-2b, glutamic acid, plicamycin, mercaptopurine, and 6-thioguanine); cytotoxic agents (e.g., carmustine, BCNU, lomustine, CCNU, cytosine arabinoside, cyclophosphamide, estramustine, hydroxyurea, procarbazine, mitomycin, busulfan, cis-platin, and vincristine sulfate); hormones (e.g., medroxyprogesterone, estramustine phosphate sodium, ethinyl
15 estradiol, estradiol, megestrol acetate, methyltestosterone, diethylstilbestrol diphosphate, chlorotrianisene, and testolactone); nitrogen mustard derivatives (e.g., mephalen, chorambucil, mechlorethamine (nitrogen mustard) and thiotepa); steroids and combinations (e.g., bethamethasone sodium phosphate); and others (e.g., dicarbazine, asparaginase, mitotane, vincristine sulfate, vinblastine sulfate, and
20 etoposide).

In a specific embodiment, Therapeutics of the invention are administered in combination with CHOP (cyclophosphamide, doxorubicin, vincristine, and prednisone) or any combination of the components of CHOP. In another embodiment, Therapeutics of the invention are administered in combination with
25 Rituximab. In a further embodiment, Therapeutics of the invention are administered with Rituxmab and CHOP, or Rituxmab and any combination of the components of CHOP.

In an additional embodiment, the Therapeutics of the invention are administered in combination with cytokines. Cytokines that may be administered
30 with the Therapeutics of the invention include, but are not limited to, IL2, IL3, IL4, IL5, IL6, IL7, IL10, IL12, IL13, IL15, anti-CD40, CD40L, IFN-gamma and TNF-alpha. In another embodiment, Therapeutics of the invention may be administered

with any interleukin, including, but not limited to, IL-1alpha, IL-1beta, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IL-16, IL-17, IL-18, IL-19, IL-20, and IL-21.

In an additional embodiment, the Therapeutics of the invention are administered in combination with angiogenic proteins. Angiogenic proteins that may be administered with the Therapeutics of the invention include, but are not limited to, Glioma Derived Growth Factor (GDGF), as disclosed in European Patent Number EP-399816; Platelet Derived Growth Factor-A (PDGF-A), as disclosed in European Patent Number EP-682110; Platelet Derived Growth Factor-B (PDGF-B), as disclosed in European Patent Number EP-282317; Placental Growth Factor (PlGF), as disclosed in International Publication Number WO 92/06194; Placental Growth Factor-2 (PlGF-2), as disclosed in Hauser et al., Growth Factors, 4:259-268 (1993); Vascular Endothelial Growth Factor (VEGF), as disclosed in International Publication Number WO 90/13649; Vascular Endothelial Growth Factor-A (VEGF-A), as disclosed in European Patent Number EP-506477; Vascular Endothelial Growth Factor-2 (VEGF-2), as disclosed in International Publication Number WO 96/39515; Vascular Endothelial Growth Factor B (VEGF-3); Vascular Endothelial Growth Factor B-186 (VEGF-B186), as disclosed in International Publication Number WO 96/26736; Vascular Endothelial Growth Factor-D (VEGF-D), as disclosed in International Publication Number WO 98/02543; Vascular Endothelial Growth Factor-D (VEGF-D), as disclosed in International Publication Number WO 98/07832; and Vascular Endothelial Growth Factor-E (VEGF-E), as disclosed in German Patent Number DE19639601. The above mentioned references are incorporated herein by reference herein.

In an additional embodiment, the Therapeutics of the invention are administered in combination with hematopoietic growth factors. Hematopoietic growth factors that may be administered with the Therapeutics of the invention include, but are not limited to, LEUKINE™ (SARGRAMOSTIM™) and NEUPOGEN™ (FILGRASTIM™).

In an additional embodiment, the Therapeutics of the invention are administered in combination with Fibroblast Growth Factors. Fibroblast Growth Factors that may be administered with the Therapeutics of the invention include, but

are not limited to, FGF-1, FGF-2, FGF-3, FGF-4, FGF-5, FGF-6, FGF-7, FGF-8, FGF-9, FGF-10, FGF-11, FGF-12, FGF-13, FGF-14, and FGF-15.

In additional embodiments, the Therapeutics of the invention are administered in combination with other therapeutic or prophylactic regimens, such as, for example,
5 radiation therapy.

Example 24: Method of Treating Decreased Levels of the Polypeptide

The present invention relates to a method for treating an individual in need of an increased level of a polypeptide of the invention in the body comprising
10 administering to such an individual a composition comprising a therapeutically effective amount of an agonist of the invention (including polypeptides of the invention). Moreover, it will be appreciated that conditions caused by a decrease in the standard or normal expression level of a secreted protein in an individual can be treated by administering the polypeptide of the present invention, preferably in the
15 secreted form. Thus, the invention also provides a method of treatment of an individual in need of an increased level of the polypeptide comprising administering to such an individual a Therapeutic comprising an amount of the polypeptide to increase the activity level of the polypeptide in such an individual.

For example, a patient with decreased levels of a polypeptide receives a daily
20 dose 0.1-100 ug/kg of the polypeptide for six consecutive days. Preferably, the polypeptide is in the secreted form. The exact details of the dosing scheme, based on administration and formulation, are provided in Example 23.

Example 25: Method of Treating Increased Levels of the Polypeptide

25 The present invention also relates to a method of treating an individual in need of a decreased level of a polypeptide of the invention in the body comprising administering to such an individual a composition comprising a therapeutically effective amount of an antagonist of the invention (including polypeptides and antibodies of the invention).

30 In one example, antisense technology is used to inhibit production of a polypeptide of the present invention. This technology is one example of a method of

decreasing levels of a polypeptide, preferably a secreted form, due to a variety of etiologies, such as cancer. For example, a patient diagnosed with abnormally increased levels of a polypeptide is administered intravenously antisense polynucleotides at 0.5, 1.0, 1.5, 2.0 and 3.0 mg/kg day for 21 days. This treatment is repeated after a 7-day rest period if the treatment was well tolerated. The formulation of the antisense polynucleotide is provided in Example 23.

Example 26: Method of Treatment Using Gene Therapy-Ex Vivo

One method of gene therapy transplants fibroblasts, which are capable of expressing a polypeptide, onto a patient. Generally, fibroblasts are obtained from a subject by skin biopsy. The resulting tissue is placed in tissue-culture medium and separated into small pieces. Small chunks of the tissue are placed on a wet surface of a tissue culture flask, approximately ten pieces are placed in each flask. The flask is turned upside down, closed tight and left at room temperature over night. After 24 hours at room temperature, the flask is inverted and the chunks of tissue remain fixed to the bottom of the flask and fresh media (e.g., Ham's F12 media, with 10% FBS, penicillin and streptomycin) is added. The flasks are then incubated at 37 degree C for approximately one week.

At this time, fresh media is added and subsequently changed every several days. After an additional two weeks in culture, a monolayer of fibroblasts emerge. The monolayer is trypsinized and scaled into larger flasks.

pMV-7 (Kirschmeier, P.T. et al., DNA, 7:219-25 (1988)), flanked by the long terminal repeats of the Moloney murine sarcoma virus, is digested with EcoRI and HindIII and subsequently treated with calf intestinal phosphatase. The linear vector is fractionated on agarose gel and purified, using glass beads.

The cDNA encoding a polypeptide of the present invention can be amplified using PCR primers which correspond to the 5' and 3' end sequences respectively as set forth in Example 1 using primers and having appropriate restriction sites and initiation/stop codons, if necessary. Preferably, the 5' primer contains an EcoRI site and the 3' primer includes a HindIII site. Equal quantities of the Moloney murine sarcoma virus linear backbone and the amplified EcoRI and-HindIII fragment are added together, in the presence of T4 DNA ligase. The resulting mixture is

maintained under conditions appropriate for ligation of the two fragments. The ligation mixture is then used to transform bacteria HB101, which are then plated onto agar containing kanamycin for the purpose of confirming that the vector has the gene of interest properly inserted.

5 The amphotropic pA317 or GP+am12 packaging cells are grown in tissue culture to confluent density in Dulbecco's Modified Eagles Medium (DMEM) with 10% calf serum (CS), penicillin and streptomycin. The MSV vector containing the gene is then added to the media and the packaging cells transduced with the vector. The packaging cells now produce infectious viral particles containing the gene (the
10 packaging cells are now referred to as producer cells).

 Fresh media is added to the transduced producer cells, and subsequently, the media is harvested from a 10 cm plate of confluent producer cells. The spent media, containing the infectious viral particles, is filtered through a millipore filter to remove detached producer cells and this media is then used to infect fibroblast cells. Media is
15 removed from a sub-confluent plate of fibroblasts and quickly replaced with the media from the producer cells. This media is removed and replaced with fresh media. If the titer of virus is high, then virtually all fibroblasts will be infected and no selection is required. If the titer is very low, then it is necessary to use a retroviral vector that has a selectable marker, such as neo or his. Once the fibroblasts have been
20 efficiently infected, the fibroblasts are analyzed to determine whether protein is produced.

 The engineered fibroblasts are then transplanted onto the host, either alone or after having been grown to confluence on cytodex 3 microcarrier beads.

25 **Example 27: Gene Therapy Using Endogenous Genes Corresponding To Polynucleotides of the Invention**

 Another method of gene therapy according to the present invention involves operably associating the endogenous polynucleotide sequence of the invention with a promoter via homologous recombination as described, for example, in U.S. Patent
30 NO: 5,641,670, issued June 24, 1997; International Publication NO: WO 96/29411, published September 26, 1996; International Publication NO: WO 94/12650, published August 4, 1994; Koller et al., *Proc. Natl. Acad. Sci. USA*, 86:8932-8935

(1989); and Zijlstra et al., *Nature*, 342:435-438 (1989). This method involves the activation of a gene which is present in the target cells, but which is not expressed in the cells, or is expressed at a lower level than desired.

Polynucleotide constructs are made which contain a promoter and targeting sequences, which are homologous to the 5' non-coding sequence of endogenous polynucleotide sequence, flanking the promoter. The targeting sequence will be sufficiently near the 5' end of the polynucleotide sequence so the promoter will be operably linked to the endogenous sequence upon homologous recombination. The promoter and the targeting sequences can be amplified using PCR. Preferably, the amplified promoter contains distinct restriction enzyme sites on the 5' and 3' ends. Preferably, the 3' end of the first targeting sequence contains the same restriction enzyme site as the 5' end of the amplified promoter and the 5' end of the second targeting sequence contains the same restriction site as the 3' end of the amplified promoter.

The amplified promoter and the amplified targeting sequences are digested with the appropriate restriction enzymes and subsequently treated with calf intestinal phosphatase. The digested promoter and digested targeting sequences are added together in the presence of T4 DNA ligase. The resulting mixture is maintained under conditions appropriate for ligation of the two fragments. The construct is size fractionated on an agarose gel then purified by phenol extraction and ethanol precipitation.

In this Example, the polynucleotide constructs are administered as naked polynucleotides via electroporation. However, the polynucleotide constructs may also be administered with transfection-facilitating agents, such as liposomes, viral sequences, viral particles, precipitating agents, etc. Such methods of delivery are known in the art.

Once the cells are transfected, homologous recombination will take place which results in the promoter being operably linked to the endogenous polynucleotide sequence. This results in the expression of polynucleotide corresponding to the polynucleotide in the cell. Expression may be detected by immunological staining, or any other method known in the art.

Fibroblasts are obtained from a subject by skin biopsy. The resulting tissue is placed in DMEM + 10% fetal calf serum. Exponentially growing or early stationary phase fibroblasts are trypsinized and rinsed from the plastic surface with nutrient medium. An aliquot of the cell suspension is removed for counting, and the remaining
5 cells are subjected to centrifugation. The supernatant is aspirated and the pellet is resuspended in 5 ml of electroporation buffer (20 mM HEPES pH 7.3, 137 mM NaCl, 5 mM KCl, 0.7 mM Na₂ HPO₄, 6 mM dextrose). The cells are recentrifuged, the supernatant aspirated, and the cells resuspended in electroporation buffer containing 1 mg/ml acetylated bovine serum albumin. The final cell suspension contains
10 approximately 3×10^6 cells/ml. Electroporation should be performed immediately following resuspension.

Plasmid DNA is prepared according to standard techniques. For example, to construct a plasmid for targeting to the locus corresponding to the polynucleotide of the invention, plasmid pUC18 (MBI Fermentas, Amherst, NY) is digested with
15 HindIII. The CMV promoter is amplified by PCR with an XbaI site on the 5' end and a BamHI site on the 3' end. Two non-coding sequences are amplified via PCR: one non-coding sequence (fragment 1) is amplified with a HindIII site at the 5' end and an Xba site at the 3' end; the other non-coding sequence (fragment 2) is amplified with a BamHI site at the 5' end and a HindIII site at the 3' end. The CMV promoter and the
20 fragments (1 and 2) are digested with the appropriate enzymes (CMV promoter - XbaI and BamHI; fragment 1 - XbaI; fragment 2 - BamHI) and ligated together. The resulting ligation product is digested with HindIII, and ligated with the HindIII-digested pUC18 plasmid.

Plasmid DNA is added to a sterile cuvette with a 0.4 cm electrode gap
25 (Bio-Rad). The final DNA concentration is generally at least 120 µg/ml. 0.5 ml of the cell suspension (containing approximately 1.5×10^6 cells) is then added to the cuvette, and the cell suspension and DNA solutions are gently mixed. Electroporation is performed with a Gene-Pulser apparatus (Bio-Rad). Capacitance and voltage are set at 960 µF and 250-300 V, respectively. As voltage increases, cell survival decreases, but
30 the percentage of surviving cells that stably incorporate the introduced DNA into their genome increases dramatically. Given these parameters, a pulse time of approximately 14-20 mSec should be observed.

Electroporated cells are maintained at room temperature for approximately 5 min, and the contents of the cuvette are then gently removed with a sterile transfer pipette. The cells are added directly to 10 ml of prewarmed nutrient media (DMEM with 15% calf serum) in a 10 cm dish and incubated at 37 degree C. The following
5 day, the media is aspirated and replaced with 10 ml of fresh media and incubated for a further 16-24 hours.

The engineered fibroblasts are then injected into the host, either alone or after having been grown to confluence on cytodex 3 microcarrier beads. The fibroblasts now produce the protein product. The fibroblasts can then be introduced into a
10 patient as described above.

Example 28: Method of Treatment Using Gene Therapy - In Vivo

Another aspect of the present invention is using *in vivo* gene therapy methods to treat disorders, diseases and conditions. The gene therapy method relates to the
15 introduction of naked nucleic acid (DNA, RNA, and antisense DNA or RNA) sequences into an animal to increase or decrease the expression of the polypeptide. The polynucleotide of the present invention may be operatively linked to a promoter or any other genetic elements necessary for the expression of the polypeptide by the target tissue. Such gene therapy and delivery techniques and methods are known in
20 the art, see, for example, WO90/11092, WO98/11779; U.S. Patent NO. 5693622, 5705151; 5580859; Tabata et al., Cardiovasc. Res. 35(3):470-479 (1997); Chao et al., Pharmacol. Res. 35(6):517-522 (1997); Wolff, Neuromuscul. Disord. 7(5):314-318 (1997); Schwartz et al., Gene Ther. 3(5):405-411 (1996); Tsurumi et al., Circulation 94(12):3281-3290 (1996) (incorporated herein by reference).

25 The polynucleotide constructs may be delivered by any method that delivers injectable materials to the cells of an animal, such as, injection into the interstitial space of tissues (heart, muscle, skin, lung, liver, intestine and the like). The polynucleotide constructs can be delivered in a pharmaceutically acceptable liquid or aqueous carrier.

30 The term "naked" polynucleotide, DNA or RNA, refers to sequences that are free from any delivery vehicle that acts to assist, promote, or facilitate entry into the cell, including viral sequences, viral particles, liposome formulations, lipofectin or

precipitating agents and the like. However, the polynucleotides of the present invention may also be delivered in liposome formulations (such as those taught in Felgner P.L. et al. (1995) Ann. NY Acad. Sci. 772:126-139 and Abdallah B. et al. (1995) Biol. Cell 85(1):1-7) which can be prepared by methods well known to those skilled in the art.

The polynucleotide vector constructs used in the gene therapy method are preferably constructs that will not integrate into the host genome nor will they contain sequences that allow for replication. Any strong promoter known to those skilled in the art can be used for driving the expression of DNA. Unlike other gene therapies techniques, one major advantage of introducing naked nucleic acid sequences into target cells is the transitory nature of the polynucleotide synthesis in the cells. Studies have shown that non-replicating DNA sequences can be introduced into cells to provide production of the desired polypeptide for periods of up to six months.

The polynucleotide construct can be delivered to the interstitial space of tissues within the an animal, including of muscle, skin, brain, lung, liver, spleen, bone marrow, thymus, heart, lymph, blood, bone, cartilage, pancreas, kidney, gall bladder, stomach, intestine, testis, ovary, uterus, rectum, nervous system, eye, gland, and connective tissue. Interstitial space of the tissues comprises the intercellular fluid, mucopolysaccharide matrix among the reticular fibers of organ tissues, elastic fibers in the walls of vessels or chambers, collagen fibers of fibrous tissues, or that same matrix within connective tissue ensheathing muscle cells or in the lacunae of bone. It is similarly the space occupied by the plasma of the circulation and the lymph fluid of the lymphatic channels. Delivery to the interstitial space of muscle tissue is preferred for the reasons discussed below. They may be conveniently delivered by injection into the tissues comprising these cells. They are preferably delivered to and expressed in persistent, non-dividing cells which are differentiated, although delivery and expression may be achieved in non-differentiated or less completely differentiated cells, such as, for example, stem cells of blood or skin fibroblasts. *In vivo* muscle cells are particularly competent in their ability to take up and express polynucleotides.

For the naked polynucleotide injection, an effective dosage amount of DNA or RNA will be in the range of from about 0.05 g/kg body weight to about 50 mg/kg

body weight. Preferably the dosage will be from about 0.005 mg/kg to about 20 mg/kg and more preferably from about 0.05 mg/kg to about 5 mg/kg. Of course, as the artisan of ordinary skill will appreciate, this dosage will vary according to the tissue site of injection. The appropriate and effective dosage of nucleic acid sequence
5 can readily be determined by those of ordinary skill in the art and may depend on the condition being treated and the route of administration. The preferred route of administration is by the parenteral route of injection into the interstitial space of tissues. However, other parenteral routes may also be used, such as, inhalation of an aerosol formulation particularly for delivery to lungs or bronchial tissues, throat or
10 mucous membranes of the nose. In addition, naked polynucleotide constructs can be delivered to arteries during angioplasty by the catheter used in the procedure.

The dose response effects of injected polynucleotide in muscle *in vivo* is determined as follows. Suitable template DNA for production of mRNA coding for polypeptide of the present invention is prepared in accordance with a standard
15 recombinant DNA methodology. The template DNA, which may be either circular or linear, is either used as naked DNA or complexed with liposomes. The quadriceps muscles of mice are then injected with various amounts of the template DNA.

Five to six week old female and male Balb/C mice are anesthetized by intraperitoneal injection with 0.3 ml of 2.5% Avertin. A 1.5 cm incision is made on
20 the anterior thigh, and the quadriceps muscle is directly visualized. The template DNA is injected in 0.1 ml of carrier in a 1 cc syringe through a 27 gauge needle over one minute, approximately 0.5 cm from the distal insertion site of the muscle into the knee and about 0.2 cm deep. A suture is placed over the injection site for future localization, and the skin is closed with stainless steel clips.

25 After an appropriate incubation time (e.g., 7 days) muscle extracts are prepared by excising the entire quadriceps. Every fifth 15 um cross-section of the individual quadriceps muscles is histochemically stained for protein expression. A time course for protein expression may be done in a similar fashion except that quadriceps from different mice are harvested at different times. Persistence of DNA
30 in muscle following injection may be determined by Southern blot analysis after preparing total cellular DNA and HIRT supernatants from injected and control mice. The results of the above experimentation in mice can be use to extrapolate proper

dosages and other treatment parameters in humans and other animals using naked DNA.

Example 29: Transgenic Animals.

5 The polypeptides of the invention can also be expressed in transgenic animals. Animals of any species, including, but not limited to, mice, rats, rabbits, hamsters, guinea pigs, pigs, micro-pigs, goats, sheep, cows and non-human primates, *e.g.*, baboons, monkeys, and chimpanzees may be used to generate transgenic animals. In a specific embodiment, techniques described herein or otherwise known in the art, are
10 used to express polypeptides of the invention in humans, as part of a gene therapy protocol.

 Any technique known in the art may be used to introduce the transgene (*i.e.*, polynucleotides of the invention) into animals to produce the founder lines of transgenic animals. Such techniques include, but are not limited to, pronuclear
15 microinjection (Paterson et al., Appl. Microbiol. Biotechnol. 40:691-698 (1994); Carver et al., Biotechnology (NY) 11:1263-1270 (1993); Wright et al., Biotechnology (NY) 9:830-834 (1991); and Hoppe et al., U.S. Pat. No. 4,873,191 (1989)); retrovirus mediated gene transfer into germ lines (Van der Putten et al., Proc. Natl. Acad. Sci., USA 82:6148-6152 (1985)), blastocysts or embryos; gene targeting in embryonic
20 stem cells (Thompson et al., Cell 56:313-321 (1989)); electroporation of cells or embryos (Lo, 1983, Mol Cell. Biol. 3:1803-1814 (1983)); introduction of the polynucleotides of the invention using a gene gun (see, *e.g.*, Ulmer et al., Science 259:1745 (1993); introducing nucleic acid constructs into embryonic pluripotent stem cells and transferring the stem cells back into the blastocyst; and sperm-
25 mediated gene transfer (Lavitrano et al., Cell 57:717-723 (1989); etc. For a review of such techniques, see Gordon, "Transgenic Animals," Intl. Rev. Cytol. 115:171-229 (1989), which is incorporated by reference herein in its entirety.

 Any technique known in the art may be used to produce transgenic clones containing polynucleotides of the invention, for example, nuclear transfer into
30 enucleated oocytes of nuclei from cultured embryonic, fetal, or adult cells induced to quiescence (Campell et al., Nature 380:64-66 (1996); Wilmut et al., Nature 385:810-813 (1997)).

The present invention provides for transgenic animals that carry the transgene in all their cells, as well as animals which carry the transgene in some, but not all their cells, *i.e.*, mosaic animals or chimeric. The transgene may be integrated as a single transgene or as multiple copies such as in concatamers, *e.g.*, head-to-head tandems or
5 head-to-tail tandems. The transgene may also be selectively introduced into and activated in a particular cell type by following, for example, the teaching of Lasko et al. (Lasko et al., Proc. Natl. Acad. Sci. USA 89:6232-6236 (1992)). The regulatory sequences required for such a cell-type specific activation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art. When
10 it is desired that the polynucleotide transgene be integrated into the chromosomal site of the endogenous gene, gene targeting is preferred. Briefly, when such a technique is to be utilized, vectors containing some nucleotide sequences homologous to the endogenous gene are designed for the purpose of integrating, via homologous recombination with chromosomal sequences, into and disrupting the function of the
15 nucleotide sequence of the endogenous gene. The transgene may also be selectively introduced into a particular cell type, thus inactivating the endogenous gene in only that cell type, by following, for example, the teaching of Gu et al. (Gu et al., Science 265:103-106 (1994)). The regulatory sequences required for such a cell-type specific inactivation will depend upon the particular cell type of interest, and will be apparent
20 to those of skill in the art.

Once transgenic animals have been generated, the expression of the recombinant gene may be assayed utilizing standard techniques. Initial screening may be accomplished by Southern blot analysis or PCR techniques to analyze animal tissues to verify that integration of the transgene has taken place. The level of mRNA
25 expression of the transgene in the tissues of the transgenic animals may also be assessed using techniques which include, but are not limited to, Northern blot analysis of tissue samples obtained from the animal, *in situ* hybridization analysis, and reverse transcriptase-PCR (rt-PCR). Samples of transgenic gene-expressing tissue may also be evaluated immunocytochemically or immunohistochemically using antibodies
30 specific for the transgene product.

Once the founder animals are produced, they may be bred, inbred, outbred, or crossbred to produce colonies of the particular animal. Examples of such breeding

strategies include, but are not limited to: outbreeding of founder animals with more than one integration site in order to establish separate lines; inbreeding of separate lines in order to produce compound transgenics that express the transgene at higher levels because of the effects of additive expression of each transgene; crossing of
5 heterozygous transgenic animals to produce animals homozygous for a given integration site in order to both augment expression and eliminate the need for screening of animals by DNA analysis; crossing of separate homozygous lines to produce compound heterozygous or homozygous lines; and breeding to place the transgene on a distinct background that is appropriate for an experimental model of
10 interest.

Transgenic animals of the invention have uses which include, but are not limited to, animal model systems useful in elaborating the biological function of polypeptides of the present invention, studying diseases, disorders, and/or conditions associated with aberrant expression, and in screening for compounds effective in
15 ameliorating such diseases, disorders, and/or conditions.

Example 30: Knock-Out Animals.

Endogenous gene expression can also be reduced by inactivating or "knocking out" the gene and/or its promoter using targeted homologous recombination. (*E.g.*,
20 see Smithies et al., *Nature* 317:230-234 (1985); Thomas & Capecchi, *Cell* 51:503-512 (1987); Thompson et al., *Cell* 5:313-321 (1989); each of which is incorporated by reference herein in its entirety). For example, a mutant, non-functional polynucleotide of the invention (or a completely unrelated DNA sequence) flanked by DNA homologous to the endogenous polynucleotide sequence (either the coding
25 regions or regulatory regions of the gene) can be used, with or without a selectable marker and/or a negative selectable marker, to transfect cells that express polypeptides of the invention *in vivo*. In another embodiment, techniques known in the art are used to generate knockouts in cells that contain, but do not express the gene of interest. Insertion of the DNA construct, via targeted homologous recombination,
30 results in inactivation of the targeted gene. Such approaches are particularly suited in research and agricultural fields where modifications to embryonic stem cells can be used to generate animal offspring with an inactive targeted gene (*e.g.*, see Thomas &

Capecchi 1987 and Thompson 1989, *supra*). However this approach can be routinely adapted for use in humans provided the recombinant DNA constructs are directly administered or targeted to the required site *in vivo* using appropriate viral vectors that will be apparent to those of skill in the art.

5 In further embodiments of the invention, cells that are genetically engineered to express the polypeptides of the invention, or alternatively, that are genetically engineered not to express the polypeptides of the invention (e.g., knockouts) are administered to a patient *in vivo*. Such cells may be obtained from the patient (i.e., animal, including human) or an MHC compatible donor and can include, but are not
10 limited to fibroblasts, bone marrow cells, blood cells (e.g., lymphocytes), adipocytes, muscle cells, endothelial cells etc. The cells are genetically engineered *in vitro* using recombinant DNA techniques to introduce the coding sequence of polypeptides of the invention into the cells, or alternatively, to disrupt the coding sequence and/or endogenous regulatory sequence associated with the polypeptides of the invention,
15 e.g., by transduction (using viral vectors, and preferably vectors that integrate the transgene into the cell genome) or transfection procedures, including, but not limited to, the use of plasmids, cosmids, YACs, naked DNA, electroporation, liposomes, etc. The coding sequence of the polypeptides of the invention can be placed under the control of a strong constitutive or inducible promoter or promoter/enhancer to achieve
20 expression, and preferably secretion, of the polypeptides of the invention. The engineered cells which express and preferably secrete the polypeptides of the invention can be introduced into the patient systemically, e.g., in the circulation, or intraperitoneally.

 Alternatively, the cells can be incorporated into a matrix and implanted in the
25 body, e.g., genetically engineered fibroblasts can be implanted as part of a skin graft; genetically engineered endothelial cells can be implanted as part of a lymphatic or vascular graft. (See, for example, Anderson et al. U.S. Patent No. 5,399,349; and Mulligan & Wilson, U.S. Patent No. 5,460,959 each of which is incorporated by reference herein in its entirety).

30 When the cells to be administered are non-autologous or non-MHC compatible cells, they can be administered using well known techniques which prevent the development of a host immune response against the introduced cells. For

example, the cells may be introduced in an encapsulated form which, while allowing for an exchange of components with the immediate extracellular environment, does not allow the introduced cells to be recognized by the host immune system.

Transgenic and "knock-out" animals of the invention have uses which include, but are not limited to, animal model systems useful in elaborating the biological function of polypeptides of the present invention, studying diseases, disorders, and/or conditions associated with aberrant expression, and in screening for compounds effective in ameliorating such diseases, disorders, and/or conditions.

10 **Example 31: Production of an Antibody**

a) Hybridoma Technology

The antibodies of the present invention can be prepared by a variety of methods. (See, Current Protocols, Chapter 2.) As one example of such methods, cells expressing polypeptide(s) of the invention are administered to an animal to induce the production of sera containing polyclonal antibodies. In a preferred method, a preparation of polypeptide(s) of the invention is prepared and purified to render it substantially free of natural contaminants. Such a preparation is then introduced into an animal in order to produce polyclonal antisera of greater specific activity.

Monoclonal antibodies specific for polypeptide(s) of the invention are prepared using hybridoma technology. (Kohler et al., Nature 256:495 (1975); Kohler et al., Eur. J. Immunol. 6:511 (1976); Kohler et al., Eur. J. Immunol. 6:292 (1976); Hammerling et al., in: Monoclonal Antibodies and T-Cell Hybridomas, Elsevier, N.Y., pp. 563-681 (1981)). In general, an animal (preferably a mouse) is immunized with polypeptide(s) of the invention, or, more preferably, with a secreted polypeptide-expressing cell. Such polypeptide-expressing cells are cultured in any suitable tissue culture medium, preferably in Earle's modified Eagle's medium supplemented with 10% fetal bovine serum (inactivated at about 56°C), and supplemented with about 10 g/l of nonessential amino acids, about 1,000 U/ml of penicillin, and about 100 µg/ml of streptomycin.

The splenocytes of such mice are extracted and fused with a suitable myeloma cell line. Any suitable myeloma cell line may be employed in accordance with the present invention; however, it is preferable to employ the parent myeloma cell line

(SP2O), available from the ATCC. After fusion, the resulting hybridoma cells are selectively maintained in HAT medium, and then cloned by limiting dilution as described by Wands et al. (Gastroenterology 80:225-232 (1981)). The hybridoma cells obtained through such a selection are then assayed to identify clones which
5 secrete antibodies capable of binding the polypeptide(s) of the invention.

Alternatively, additional antibodies capable of binding polypeptide(s) of the invention can be produced in a two-step procedure using anti-idiotypic antibodies. Such a method makes use of the fact that antibodies are themselves antigens, and therefore, it is possible to obtain an antibody which binds to a second antibody. In
10 accordance with this method, protein specific antibodies are used to immunize an animal, preferably a mouse. The splenocytes of such an animal are then used to produce hybridoma cells, and the hybridoma cells are screened to identify clones which produce an antibody whose ability to bind to the polypeptide(s) of the invention protein-specific antibody can be blocked by polypeptide(s) of the invention.

15 Such antibodies comprise anti-idiotypic antibodies to the polypeptide(s) of the invention protein-specific antibody and are used to immunize an animal to induce formation of further polypeptide(s) of the invention protein-specific antibodies.

For in vivo use of antibodies in humans, an antibody is "humanized". Such antibodies can be produced using genetic constructs derived from hybridoma cells
20 producing the monoclonal antibodies described above. Methods for producing chimeric and humanized antibodies are known in the art and are discussed herein. (See, for review, Morrison, Science 229:1202 (1985); Oi et al., BioTechniques 4:214 (1986); Cabilly et al., U.S. Patent No. 4,816,567; Taniguchi et al., EP 171496; Morrison et al., EP 173494; Neuberger et al., WO 8601533; Robinson et al., WO
25 8702671; Boulianne et al., Nature 312:643 (1984); Neuberger et al., Nature 314:268 (1985).)

b) Isolation Of Antibody Fragments Directed

polypeptide(s) of the invention From A Library Of scFvs

30 Naturally occurring V-genes isolated from human PBLs are constructed into a library of antibody fragments which contain reactivities against polypeptide(s) of the

invention to which the donor may or may not have been exposed (see e.g., U.S. Patent 5,885,793 incorporated herein by reference in its entirety).

Rescue of the Library. A library of scFvs is constructed from the RNA of human PBLs as described in PCT publication WO 92/01047. To rescue phage displaying antibody fragments, approximately 109 E. coli harboring the phagemid are used to inoculate 50 ml of 2xTY containing 1% glucose and 100 µg/ml of ampicillin (2xTY-AMP-GLU) and grown to an O.D. of 0.8 with shaking. Five ml of this culture is used to inoculate 50 ml of 2xTY-AMP-GLU, 2 x 10⁸ TU of delta gene 3 helper (M13 delta gene III, see PCT publication WO 92/01047) are added and the culture incubated at 37°C for 45 minutes without shaking and then at 37°C for 45 minutes with shaking. The culture is centrifuged at 4000 r.p.m. for 10 min. and the pellet resuspended in 2 liters of 2xTY containing 100 µg/ml ampicillin and 50 ug/ml kanamycin and grown overnight. Phage are prepared as described in PCT publication WO 92/01047.

M13 delta gene III is prepared as follows: M13 delta gene III helper phage does not encode gene III protein, hence the phage(mid) displaying antibody fragments have a greater avidity of binding to antigen. Infectious M13 delta gene III particles are made by growing the helper phage in cells harboring a pUC19 derivative supplying the wild type gene III protein during phage morphogenesis. The culture is incubated for 1 hour at 37° C without shaking and then for a further hour at 37°C with shaking. Cells are spun down (IEC-Centra 8,400 r.p.m. for 10 min), resuspended in 300 ml 2xTY broth containing 100 µg ampicillin/ml and 25 µg kanamycin/ml (2xTY-AMP-KAN) and grown overnight, shaking at 37°C. Phage particles are purified and concentrated from the culture medium by two PEG-precipitations (Sambrook et al., 1990), resuspended in 2 ml PBS and passed through a 0.45 µm filter (Minisart NML; Sartorius) to give a final concentration of approximately 10¹³ transducing units/ml (ampicillin-resistant clones).

Panning of the Library. Immunotubes (Nunc) are coated overnight in PBS with 4 ml of either 100 µg/ml or 10 µg/ml of a polypeptide of the present invention. Tubes are blocked with 2% Marvel-PBS for 2 hours at 37°C and then washed 3 times in PBS. Approximately 10¹³ TU of phage is applied to the tube and incubated for 30 minutes at room temperature tumbling on an over and under turntable and then left to

stand for another 1.5 hours. Tubes are washed 10 times with PBS 0.1% Tween-20 and 10 times with PBS. Phage are eluted by adding 1 ml of 100 mM triethylamine and rotating 15 minutes on an under and over turntable after which the solution is immediately neutralized with 0.5 ml of 1.0M Tris-HCl, pH 7.4. Phage are then used to infect 10 ml of mid-log E. coli TG1 by incubating eluted phage with bacteria for 30 minutes at 37°C. The E. coli are then plated on TYE plates containing 1% glucose and 100 µg/ml ampicillin. The resulting bacterial library is then rescued with delta gene 3 helper phage as described above to prepare phage for a subsequent round of selection. This process is then repeated for a total of 4 rounds of affinity purification with tube-washing increased to 20 times with PBS, 0.1% Tween-20 and 20 times with PBS for rounds 3 and 4.

Characterization of Binders. Eluted phage from the 3rd and 4th rounds of selection are used to infect E. coli HB 2151 and soluble scFv is produced (Marks, et al., 1991) from single colonies for assay. ELISAs are performed with microtitre plates coated with either 10 pg/ml of the polypeptide of the present invention in 50 mM bicarbonate pH 9.6. Clones positive in ELISA are further characterized by PCR fingerprinting (see, e.g., PCT publication WO 92/01047) and then by sequencing. These ELISA positive clones may also be further characterized by techniques known in the art, such as, for example, epitope mapping, binding affinity, receptor signal transduction, ability to block or competitively inhibit antibody/antigen binding, and competitive agonistic or antagonistic activity.

Example 32: Assays Detecting Stimulation or Inhibition of B cell Proliferation and Differentiation

Generation of functional humoral immune responses requires both soluble and cognate signaling between B-lineage cells and their microenvironment. Signals may impart a positive stimulus that allows a B-lineage cell to continue its programmed development, or a negative stimulus that instructs the cell to arrest its current developmental pathway. To date, numerous stimulatory and inhibitory signals have been found to influence B cell responsiveness including IL-2, IL-4, IL-5, IL-6, IL-7, IL10, IL-13, IL-14 and IL-15. Interestingly, these signals are by themselves weak effectors but can,

in combination with various co-stimulatory proteins, induce activation, proliferation, differentiation, homing, tolerance and death among B cell populations.

One of the best studied classes of B-cell co-stimulatory proteins is the TNF-superfamily. Within this family CD40, CD27, and CD30 along with their respective
5 ligands CD154, CD70, and CD153 have been found to regulate a variety of immune responses. Assays which allow for the detection and/or observation of the proliferation and differentiation of these B-cell populations and their precursors are valuable tools in determining the effects various proteins may have on these B-cell populations in terms of proliferation and differentiation. Listed below are two assays designed to allow for the
10 detection of the differentiation, proliferation, or inhibition of B-cell populations and their precursors.

In Vitro Assay- Purified polypeptides of the invention, or truncated forms thereof, is assessed for its ability to induce activation, proliferation, differentiation or inhibition and/or death in B-cell populations and their precursors. The activity of the
15 polypeptides of the invention on purified human tonsillar B cells, measured qualitatively over the dose range from 0.1 to 10,000 ng/mL, is assessed in a standard B-lymphocyte co-stimulation assay in which purified tonsillar B cells are cultured in the presence of either formalin-fixed Staphylococcus aureus Cowan I (SAC) or immobilized anti-human IgM antibody as the priming agent. Second signals such as
20 IL-2 and IL-15 synergize with SAC and IgM crosslinking to elicit B cell proliferation as measured by tritiated-thymidine incorporation. Novel synergizing agents can be readily identified using this assay. The assay involves isolating human tonsillar B cells by magnetic bead (MACS) depletion of CD3-positive cells. The resulting cell population is greater than 95% B cells as assessed by expression of CD45R(B220).

25 Various dilutions of each sample are placed into individual wells of a 96-well plate to which are added 10^5 B-cells suspended in culture medium (RPMI 1640 containing 10% FBS, 5×10^{-5} M 2ME, 100U/ml penicillin, 10ug/ml streptomycin, and 10^{-5} dilution of SAC) in a total volume of 150ul. Proliferation or inhibition is quantitated by a 20h pulse (1uCi/well) with 3H-thymidine (6.7 Ci/mM) beginning 72h post factor addition. The
30 positive and negative controls are IL2 and medium respectively.

In Vivo Assay- BALB/c mice are injected (i.p.) twice per day with buffer only, or 2 mg/Kg of a polypeptide of the invention, or truncated forms thereof. Mice

receive this treatment for 4 consecutive days, at which time they are sacrificed and various tissues and serum collected for analyses. Comparison of H&E sections from normal spleens and spleens treated with polypeptides of the invention identify the results of the activity of the polypeptides on spleen cells, such as the diffusion of peri-arterial lymphatic sheaths, and/or significant increases in the nucleated cellularity of the red pulp regions, which may indicate the activation of the differentiation and proliferation of B-cell populations. Immunohistochemical studies using a B cell marker, anti-CD45R(B220), are used to determine whether any physiological changes to splenic cells, such as splenic disorganization, are due to increased B-cell representation within loosely defined B-cell zones that infiltrate established T-cell regions.

Flow cytometric analyses of the spleens from mice treated with polypeptide is used to indicate whether the polypeptide specifically increases the proportion of ThB+, CD45R(B220)dull B cells over that which is observed in control mice.

Likewise, a predicted consequence of increased mature B-cell representation in vivo is a relative increase in serum Ig titers. Accordingly, serum IgM and IgA levels are compared between buffer and polypeptide-treated mice.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides of the invention (e.g., gene therapy), agonists, and/or antagonists of polynucleotides or polypeptides of the invention.

Example 33: T Cell Proliferation Assay

Proliferation assay for Resting PBLs.

A CD3-induced proliferation assay is performed on PBMCs and is measured by the uptake of ³H-thymidine. The assay is performed as follows. Ninety-six well plates are coated with 100 microliters per well of mAb to CD3 (HIT3a, Pharmingen) or isotype-matched control mAb (B33.1) overnight at 4 °C (1 microgram/ml in .05M bicarbonate buffer, pH 9.5), then washed three times with PBS. PBMC are isolated by F/H gradient centrifugation from human peripheral blood and added to quadruplicate wells (5 x 10⁴/well) of mAb coated plates in RPMI containing 10% FCS and P/S in the presence of varying concentrations of TNF Delta and/or TNF Epsilon protein (total volume 200 microliters). Relevant protein buffer and medium alone are controls.

After 48 hr. culture at 37 °C, plates are spun for 2 min. at 1000 rpm and 100 microliters of supernatant is removed and stored -20 °C for measurement of IL-2 (or other cytokines) if effect on proliferation is observed. Wells are supplemented with 100 microliters of medium containing 0.5 microcuries of ³H-thymidine and cultured at 37 °C for 18-24 hr. Wells are harvested and

5 incorporation of ³H-thymidine used as a measure of proliferation. Anti-CD3 alone is the positive control for proliferation. IL-2 (100 U/ml) is also used as a control which enhances proliferation. Control antibody which does not induce proliferation of T cells is used as the negative controls for the effects of TNF Delta and/or TNF Epsilon proteins.

Alternatively, a proliferation assay on resting PBL (peripheral blood

10 lymphocytes) is measured by the up-take of ³H-thymidine. The assay is performed as follows. PBMC are isolated by Ficoll (LSM, ICN Biotechnologies, Aurora, Ohio) gradient centrifugation from human peripheral blood, and are cultured overnight in 10% (Fetal Calf Serum, Biofluids, Rockville, MD)/RPMI (Gibco BRL, Gaithersburg, MD). This overnight incubation period allows the adherent cells to attach to the

15 plastic, which results in a lower background in the assay as there are fewer cells that can act as antigen presenting cells or that might be producing growth factors. The following day the non-adherent cells are collected, washed and used in the proliferation assay. The assay is performed in a 96 well plate using 2 x 10⁴ cells/well in a final volume of 200 microliters. The supernatants (e.g., CHO or 293T

20 supernatants) expressing the protein of interest are tested at a 30% final dilution, therefore 60ul are added to 140ul of 10% FCS/RPMI containing the cells. Control supernatants are used at the same final dilution and express the following proteins: vector (negative control), IL-2 (*), IFN γ , TNF α , IL-10 and TR2. In addition to the control supernatants, recombinant human IL-2 (R & D Systems, Minneapolis, MN)

25 at a final concentration of 100ng/ml is also used. After 24 hours of culture, each well is pulsed with 1uCi of ³H-thymidine (Nen, Boston, MA). Cells are then harvested 20 hours following pulsing and incorporation of ³H-thymidine is used as a measure of proliferation. Results are expressed as an average of triplicate samples plus or minus standard error.

30 (*) The amount of the control cytokines IL-2, IFN γ , TNF α and IL-10 produced in each transfection varies between 300pg to 5ng/ml.

Costimulation assay.

A costimulation assay on resting PBL (peripheral blood lymphocytes) is performed in the presence of immobilized antibodies to CD3 and CD28. The use of antibodies specific for the invariant regions of CD3 mimic the induction of T cell activation that would occur through stimulation of the T cell receptor by an antigen. Cross-linking of the TCR (first signal) in the absence of a costimulatory signal (second signal) causes very low induction of proliferation and will eventually result in a state of "anergy", which is characterized by the absence of growth and inability to produce cytokines. The addition of a costimulatory signal such as an antibody to CD28, which mimics the action of the costimulatory molecule. B7-1 expressed on activated APCs, results in enhancement of T cell responses including cell survival and production of IL-2. Therefore this type of assay allows to detect both positive and negative effects caused by addition of supernatants expressing the proteins of interest on T cell proliferation.

The assay is performed as follows. Ninety-six well plates are coated with 100ng/ml anti-CD3 and 5ug/ml anti-CD28 (Pharmingen, San Diego, CA) in a final volume of 100ul and incubated overnight at 4C. Plates are washed twice with PBS before use. PBMC are isolated by Ficoll (LSM, ICN Biotechnologies, Aurora, Ohio) gradient centrifugation from human peripheral blood, and are cultured overnight in 10% FCS(Fetal Calf Serum, Biofluids, Rockville, MD)/RPMI (Gibco BRL, Gaithersburg, MD). This overnight incubation period allows the adherent cells to attach to the plastic, which results in a lower background in the assay as there are fewer cells that can act as antigen presenting cells or that might be producing growth factors. The following day the non adherent cells are collected, washed and used in the proliferation assay. The assay is performed in a 96 well plate using 2×10^4 cells/well in a final volume of 200ul. The supernatants (e.g., CHO or 293T supernatants) expressing the protein of interest are tested at a 30% final dilution, therefore 60ul are added to 140ul of 10% FCS/RPMI containing the cells. Control supernatants are used at the same final dilution and express the following proteins: vector only (negative control), IL-2, IFN γ , TNF α , IL-10 and TR2. In addition to the control supernatants recombinant human IL-2 (R & D Systems, Minneapolis, MN) at a final concentration of 10ng/ml is also used. After 24 hours of culture, each well is

pulsed with 1uCi of ^3H -thymidine (Nen, Boston, MA). Cells are then harvested 20 hours following pulsing and incorporation of ^3H -thymidine is used as a measure of proliferation. Results are expressed as an average of triplicate samples plus or minus standard error.

5 Costimulation assay: IFN γ and IL-2 ELISA

The assay is performed as follows. Twenty-four well plates are coated with either 300ng/ml or 600ng/ml anti-CD3 and 5ug/ml anti-CD28 (Pharmingen, San Diego, CA) in a final volume of 500ul and incubated overnight at 4C. Plates are washed twice with PBS before use. PBMC are isolated by Ficoll (LSM, ICN Biotechnologies, Aurora, Ohio) gradient centrifugation from human peripheral blood, and are cultured overnight in 10% FCS(Fetal Calf Serum, Biofluids, Rockville, MD)/RPMI (Gibco BRL, Gaithersburg, MD). This overnight incubation period allows the adherent cells to attach to the plastic, which results in a lower background in the assay as there are fewer cells that can act as antigen presenting cells or that might be producing growth factors. The following day the non adherent cells are collected, washed and used in the costimulation assay. The assay is performed in the pre-coated twenty-four well plate using 1×10^5 cells/well in a final volume of 900ul. The supernatants (293T supernatants) expressing the protein of interest are tested at a 30% final dilution, therefore 300ul are added to 600ul of 10% FCS/RPMI containing the cells. Control supernatants are used at the same final dilution and express the following proteins: vector only(negative control), IL-2, IFN γ , IL-12 and IL-18. In addition to the control supernatants recombinant human IL-2 (all cytokines were purchased from R & D Systems, Minneapolis, MN) at a final concentration of 10ng/ml, IL-12 at a final concentration of 1ng/ml and IL-18 at a final concentration of 50ng/ml are also used. Controls and unknown samples are tested in duplicate. Supernatant samples (250ul) are collected 2 days and 5 days after the beginning of the assay. ELISAs to test for IFN γ and IL-2 secretion are performed using kits purchased from R & D Systems, (Minneapolis, MN). Results are expressed as an average of duplicate samples plus or minus standard error.

30

Proliferation assay for preactivated-resting T cells.

A proliferation assay on preactivated-resting T cells is performed on cells that are previously activated with the lectin phytohemagglutinin (PHA). Lectins are polymeric plant proteins that can bind to residues on T cell surface glycoproteins including the TCR and act as polyclonal activators. PBLs treated with PHA and then
5 cultured in the presence of low doses of IL-2 resemble effector T cells. These cells are generally more sensitive to further activation induced by growth factors such as IL-2. This is due to the expression of high affinity IL-2 receptors that allows this population to respond to amounts of IL-2 that are 100 fold lower than what would have an effect on a naïve T cell. Therefore the use of this type of cells might enable
10 to detect the effect of very low doses of an unknown growth factor, that would not be sufficient to induce proliferation on resting (naïve) T cells.

The assay is performed as follows. PBMC are isolated by F/H gradient centrifugation from human peripheral blood, and are cultured in 10% FCS (Fetal Calf Serum, Biofluids, Rockville, MD)/RPMI (Gibco BRL, Gaithersburg, MD) in the
15 presence of 2 µg/ml PHA (Sigma, Saint Louis, MO) for three days. The cells are then washed in PBS and cultured in 10% FCS/RPMI in the presence of 5 ng/ml of human recombinant IL-2 (R & D Systems, Minneapolis, MN) for 3 days. The cells are washed and rested in starvation medium (1% FCS/RPMI) for 16 hours prior to the beginning of the proliferation assay. An aliquot of the cells is analyzed by FACS to
20 determine the percentage of T cells (CD3 positive cells) present; this usually ranges between 93-97% depending on the donor. The assay is performed in a 96 well plate using 2×10^4 cells/well in a final volume of 200 µl. The supernatants (e.g., CHO or 293T supernatants) expressing the protein of interest are tested at a 30% final dilution, therefore 60 µl are added to 140 µl of 10% FCS/RPMI containing the cells. Control
25 supernatants are used at the same final dilution and express the following proteins: vector (negative control), IL-2, IFN γ , TNF α , IL-10 and TR2. In addition to the control supernatants recombinant human IL-2 at a final concentration of 10 ng/ml is also used. After 24 hours of culture, each well is pulsed with 1 µCi of ^3H -thymidine (Nen, Boston, MA). Cells are then harvested 20 hours following pulsing
30 and incorporation of ^3H -thymidine is used as a measure of proliferation. Results are expressed as an average of triplicate samples plus or minus standard error.

The studies described in this example test activity of polypeptides of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides of the invention (e.g., gene therapy), agonists, and/or antagonists of polynucleotides or polypeptides of the invention.

5

Example 34: Effect of Polypeptides of the Invention on the Expression of MHC Class II, Costimulatory and Adhesion Molecules and Cell Differentiation of Monocytes and Monocyte-Derived Human Dendritic Cells

10 Dendritic cells are generated by the expansion of proliferating precursors found in the peripheral blood: adherent PBMC or elutriated monocytic fractions are cultured for 7-10 days with GM-CSF (50 ng/ml) and IL-4 (20 ng/ml). These dendritic cells have the characteristic phenotype of immature cells (expression of CD1, CD80, CD86, CD40 and MHC class II antigens). Treatment with activating factors, such as TNF- α , causes a rapid
15 change in surface phenotype (increased expression of MHC class I and II, costimulatory and adhesion molecules, downregulation of FCyRII, upregulation of CD83). These changes correlate with increased antigen-presenting capacity and with functional maturation of the dendritic cells.

FACS analysis of surface antigens is performed as follows. Cells are treated 1-3
20 days with increasing concentrations of polypeptides of the invention or LPS (positive control), washed with PBS containing 1% BSA and 0.02 mM sodium azide, and then incubated with 1:20 dilution of appropriate FITC- or PE-labeled monoclonal antibodies for 30 minutes at 4 degrees C. After an additional wash, the labeled cells are analyzed by flow cytometry on a FACScan (Becton Dickinson).

25

Effect on the production of cytokines. Cytokines generated by dendritic cells, in particular IL-12, are important in the initiation of T-cell dependent immune responses. IL-12 strongly influences the development of Th1 helper T-cell immune response, and induces cytotoxic T and NK cell function. An ELISA is used to
30 measure the IL-12 release as follows. Dendritic cells (10^6 /ml) are treated with increasing concentrations of polypeptides of the invention for 24 hours. LPS (100 ng/ml) is added to the cell culture as positive control. Supernatants from the cell

cultures are then collected and analyzed for IL-12 content using commercial ELISA kit (e.g., R & D Systems (Minneapolis, MN)). The standard protocols provided with the kits are used.

5 Effect on the expression of MHC Class II, costimulatory and adhesion molecules. Three major families of cell surface antigens can be identified on monocytes: adhesion molecules, molecules involved in antigen presentation, and Fc receptor. Modulation of the expression of MHC class II antigens and other costimulatory molecules, such as B7 and ICAM-1, may result in changes in the
10 antigen presenting capacity of monocytes and ability to induce T cell activation. Increase expression of Fc receptors may correlate with improved monocyte cytotoxic activity, cytokine release and phagocytosis.

FACS analysis is used to examine the surface antigens as follows. Monocytes are treated 1-5 days with increasing concentrations of polypeptides of the invention or
15 LPS (positive control), washed with PBS containing 1% BSA and 0.02 mM sodium azide, and then incubated with 1:20 dilution of appropriate FITC- or PE-labeled monoclonal antibodies for 30 minutes at 4 degreesC. After an additional wash, the labeled cells are analyzed by flow cytometry on a FACScan (Becton Dickinson).

20 Monocyte activation and/or increased survival. Assays for molecules that activate (or alternatively, inactivate) monocytes and/or increase monocyte survival (or alternatively, decrease monocyte survival) are known in the art and may routinely be applied to determine whether a molecule of the invention functions as an inhibitor or activator of monocytes. Polypeptides, agonists, or antagonists of the invention can be
25 screened using the three assays described below. For each of these assays, Peripheral blood mononuclear cells (PBMC) are purified from single donor leukopacks (American Red Cross, Baltimore, MD) by centrifugation through a Histopaque gradient (Sigma). Monocytes are isolated from PBMC by counterflow centrifugal elutriation.

30

Monocyte Survival Assay. Human peripheral blood monocytes progressively lose viability when cultured in absence of serum or other stimuli. Their death results

from internally regulated process (apoptosis). Addition to the culture of activating factors, such as TNF-alpha dramatically improves cell survival and prevents DNA fragmentation. Propidium iodide (PI) staining is used to measure apoptosis as follows. Monocytes are cultured for 48 hours in polypropylene tubes in serum-free medium (positive control), in the presence of 100 ng/ml TNF-alpha (negative control), and in the presence of varying concentrations of the compound to be tested. Cells are suspended at a concentration of 2×10^6 /ml in PBS containing PI at a final concentration of 5 μ g/ml, and then incubated at room temperature for 5 minutes before FACSscan analysis. PI uptake has been demonstrated to correlate with DNA fragmentation in this experimental paradigm.

Effect on cytokine release. An important function of monocytes/macrophages is their regulatory activity on other cellular populations of the immune system through the release of cytokines after stimulation. An ELISA to measure cytokine release is performed as follows. Human monocytes are incubated at a density of 5×10^5 cells/ml with increasing concentrations of the a polypeptide of the invention and under the same conditions, but in the absence of the polypeptide. For IL-12 production, the cells are primed overnight with IFN (100 U/ml) in presence of a polypeptide of the invention. LPS (10 ng/ml) is then added. Conditioned media are collected after 24h and kept frozen until use. Measurement of TNF-alpha, IL-10, MCP-1 and IL-8 is then performed using a commercially available ELISA kit (e.g., R & D Systems (Minneapolis, MN)) and applying the standard protocols provided with the kit.

Oxidative burst. Purified monocytes are plated in 96-w plate at 2×10^5 cell/well. Increasing concentrations of polypeptides of the invention are added to the wells in a total volume of 0.2 ml culture medium (RPMI 1640 + 10% FCS, glutamine and antibiotics). After 3 days incubation, the plates are centrifuged and the medium is removed from the wells. To the macrophage monolayers, 0.2 ml per well of phenol red solution (140 mM NaCl, 10 mM potassium phosphate buffer pH 7.0, 5.5 mM dextrose, 0.56 mM phenol red and 19 U/ml of HRPO) is added, together with the stimulant (200 nM PMA). The plates are incubated at 37°C for 2 hours and the reaction is stopped by adding 20 μ l 1N NaOH per well. The absorbance is read at 610

nm. To calculate the amount of H₂O₂ produced by the macrophages, a standard curve of a H₂O₂ solution of known molarity is performed for each experiment.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polypeptides, polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 35: Biological Effects of Polypeptides of the Invention

Astrocyte and Neuronal Assays

Recombinant polypeptides of the invention, expressed in *Escherichia coli* and purified as described above, can be tested for activity in promoting the survival, neurite outgrowth, or phenotypic differentiation of cortical neuronal cells and for inducing the proliferation of glial fibrillary acidic protein immunopositive cells, astrocytes. The selection of cortical cells for the bioassay is based on the prevalent expression of FGF-1 and FGF-2 in cortical structures and on the previously reported enhancement of cortical neuronal survival resulting from FGF-2 treatment. A thymidine incorporation assay, for example, can be used to elucidate a polypeptide of the invention's activity on these cells.

Moreover, previous reports describing the biological effects of FGF-2 (basic FGF) on cortical or hippocampal neurons *in vitro* have demonstrated increases in both neuron survival and neurite outgrowth (Walicke et al., "Fibroblast growth factor promotes survival of dissociated hippocampal neurons and enhances neurite extension." *Proc. Natl. Acad. Sci. USA* 83:3012-3016. (1986), assay herein incorporated by reference in its entirety). However, reports from experiments done on PC-12 cells suggest that these two responses are not necessarily synonymous and may depend on not only which FGF is being tested but also on which receptor(s) are expressed on the target cells. Using the primary cortical neuronal culture paradigm, the ability of a polypeptide of the invention to induce neurite outgrowth can be compared to the response achieved with FGF-2 using, for example, a thymidine incorporation assay.

Fibroblast and endothelial cell assays

Human lung fibroblasts are obtained from Clonetics (San Diego, CA) and maintained in growth media from Clonetics. Dermal microvascular endothelial cells are obtained from Cell Applications (San Diego, CA). For proliferation assays, the human lung fibroblasts and dermal microvascular endothelial cells can be cultured at 5,000 cells/well in a 96-well plate for one day in growth medium. The cells are then incubated for one day in 0.1% BSA basal medium. After replacing the medium with fresh 0.1% BSA medium, the cells are incubated with the test proteins for 3 days. Alamar Blue (Alamar Biosciences, Sacramento, CA) is added to each well to a final concentration of 10%. The cells are incubated for 4 hr. Cell viability is measured by reading in a CytoFluor fluorescence reader. For the PGE₂ assays, the human lung fibroblasts are cultured at 5,000 cells/well in a 96-well plate for one day. After a medium change to 0.1% BSA basal medium, the cells are incubated with FGF-2 or polypeptides of the invention with or without IL-1 α for 24 hours. The supernatants are collected and assayed for PGE₂ by EIA kit (Cayman, Ann Arbor, MI). For the IL-6 assays, the human lung fibroblasts are cultured at 5,000 cells/well in a 96-well plate for one day. After a medium change to 0.1% BSA basal medium, the cells are incubated with FGF-2 or with or without polypeptides of the invention IL-1 α for 24 hours. The supernatants are collected and assayed for IL-6 by ELISA kit (Endogen, Cambridge, MA).

Human lung fibroblasts are cultured with FGF-2 or polypeptides of the invention for 3 days in basal medium before the addition of Alamar Blue to assess effects on growth of the fibroblasts. FGF-2 should show a stimulation at 10 - 2500 ng/ml which can be used to compare stimulation with polypeptides of the invention.

Parkinson Models.

The loss of motor function in Parkinson's disease is attributed to a deficiency of striatal dopamine resulting from the degeneration of the nigrostriatal dopaminergic projection neurons. An animal model for Parkinson's that has been extensively characterized involves the systemic administration of 1-methyl-4 phenyl 1,2,3,6-tetrahydropyridine (MPTP). In the CNS, MPTP is taken-up by astrocytes and catabolized by monoamine oxidase B to 1-methyl-4-phenyl pyridine (MPP⁺) and released.

Subsequently, MPP^+ is actively accumulated in dopaminergic neurons by the high-affinity reuptake transporter for dopamine. MPP^+ is then concentrated in mitochondria by the electrochemical gradient and selectively inhibits nicotinamide adenine disphosphate: ubiquinone oxidoreductionase (complex I), thereby interfering with electron transport and
5 eventually generating oxygen radicals.

It has been demonstrated in tissue culture paradigms that FGF-2 (basic FGF) has trophic activity towards nigral dopaminergic neurons (Ferrari et al., Dev. Biol. 1989). Recently, Dr. Unsicker's group has demonstrated that administering FGF-2 in gel foam implants in the striatum results in the near complete protection of nigral dopaminergic
10 neurons from the toxicity associated with MPTP exposure (Otto and Unsicker, J. Neuroscience, 1990).

Based on the data with FGF-2, polypeptides of the invention can be evaluated to determine whether it has an action similar to that of FGF-2 in enhancing dopaminergic neuronal survival *in vitro* and it can also be tested *in vivo* for protection of dopaminergic
15 neurons in the striatum from the damage associated with MPTP treatment. The potential effect of a polypeptide of the invention is first examined *in vitro* in a dopaminergic neuronal cell culture paradigm. The cultures are prepared by dissecting the midbrain floor plate from gestation day 14 Wistar rat embryos. The tissue is dissociated with trypsin and seeded at a density of 200,000 cells/cm² on polyorthinine-laminin coated glass coverslips.
20 The cells are maintained in Dulbecco's Modified Eagle's medium and F12 medium containing hormonal supplements (N1). The cultures are fixed with paraformaldehyde after 8 days *in vitro* and are processed for tyrosine hydroxylase, a specific marker for dopaminergic neurons, immunohistochemical staining. Dissociated cell cultures are prepared from embryonic rats. The culture medium is changed every third day and the
25 factors are also added at that time.

Since the dopaminergic neurons are isolated from animals at gestation day 14, a developmental time which is past the stage when the dopaminergic precursor cells are proliferating, an increase in the number of tyrosine hydroxylase immunopositive neurons would represent an increase in the number of dopaminergic neurons surviving *in vitro*.
30 Therefore, if a polypeptide of the invention acts to prolong the survival of dopaminergic neurons, it would suggest that the polypeptide may be involved in Parkinson's Disease.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

5

Example 36: The Effect of Polypeptides of the Invention on the Growth of Vascular Endothelial Cells

On day 1, human umbilical vein endothelial cells (HUVEC) are seeded at 2.5×10^4 cells/35 mm dish density in M199 medium containing 4% fetal bovine serum (FBS), 16 units/ml heparin, and 50 units/ml endothelial cell growth supplements (ECGS, Biotechnology, Inc.). On day 2, the medium is replaced with M199 containing 10% FBS, 8 units/ml heparin. A polypeptide having the amino acid sequence of SEQ ID NO:Y, and positive controls, such as VEGF and basic FGF (bFGF) are added, at varying concentrations. On days 4 and 6, the medium is replaced. On day 8, cell number is determined with a Coulter Counter.

An increase in the number of HUVEC cells indicates that the polypeptide of the invention may proliferate vascular endothelial cells.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 37: Stimulatory Effect of Polypeptides of the Invention on the Proliferation of Vascular Endothelial Cells

For evaluation of mitogenic activity of growth factors, the colorimetric MTS (3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)2H-tetrazolium) assay with the electron coupling reagent PMS (phenazine methosulfate) was performed (CellTiter 96 AQ, Promega). Cells are seeded in a 96-well plate (5,000 cells/well) in 0.1 mL serum-supplemented medium and are allowed to attach overnight. After serum-starvation for 12 hours in 0.5% FBS, conditions (bFGF, VEGF₁₆₅ or a

polypeptide of the invention in 0.5% FBS) with or without Heparin (8 U/ml) are added to wells for 48 hours. 20 mg of MTS/PMS mixture (1:0.05) are added per well and allowed to incubate for 1 hour at 37°C before measuring the absorbance at 490 nm in an ELISA plate reader. Background absorbance from control wells (some media, no cells) is subtracted, and seven wells are performed in parallel for each condition. See, Leak *et al.* *In Vitro Cell. Dev. Biol.* 30A:512-518 (1994).

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 38: Inhibition of PDGF-induced Vascular Smooth Muscle Cell Proliferation Stimulatory Effect

HAoSMC proliferation can be measured, for example, by BrdUrd incorporation. Briefly, subconfluent, quiescent cells grown on the 4-chamber slides are transfected with CRP or FITC-labeled AT2-3LP. Then, the cells are pulsed with 10% calf serum and 6 mg/ml BrdUrd. After 24 h, immunocytochemistry is performed by using BrdUrd Staining Kit (Zymed Laboratories). In brief, the cells are incubated with the biotinylated mouse anti-BrdUrd antibody at 4 degrees C for 2 h after being exposed to denaturing solution and then incubated with the streptavidin-peroxidase and diaminobenzidine. After counterstaining with hematoxylin, the cells are mounted for microscopic examination, and the BrdUrd-positive cells are counted. The BrdUrd index is calculated as a percent of the BrdUrd-positive cells to the total cell number. In addition, the simultaneous detection of the BrdUrd staining (nucleus) and the FITC uptake (cytoplasm) is performed for individual cells by the concomitant use of bright field illumination and dark field-UV fluorescent illumination. See, Hayashida et al., *J. Biol. Chem.* 6:271(36):21985-21992 (1996).

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 39: Stimulation of Endothelial Migration

5 This example will be used to explore the possibility that a polypeptide of the invention may stimulate lymphatic endothelial cell migration.

Endothelial cell migration assays are performed using a 48 well microchemotaxis chamber (Neuroprobe Inc., Cabin John, MD; Falk, W., et al., J. Immunological Methods 1980;33:239-247). Polyvinylpyrrolidone-free polycarbonate filters with a pore size of 8 μ m (Nucleopore Corp. Cambridge, MA) are coated with 0.1% gelatin for at least 6 hours at room temperature and dried under sterile air. Test substances are diluted to appropriate concentrations in M199 supplemented with 0.25% bovine serum albumin (BSA), and 25 μ l of the final dilution is placed in the lower chamber of the modified Boyden apparatus. Subconfluent, early passage (2-6) HUVEC or BMEC cultures are washed and trypsinized for the minimum time required to achieve cell detachment. After placing the filter between lower and upper chamber, 2.5×10^5 cells suspended in 50 μ l M199 containing 1% FBS are seeded in the upper compartment. The apparatus is then incubated for 5 hours at 37°C in a humidified chamber with 5% CO₂ to allow cell migration. After the incubation period, the filter is removed and the upper side of the filter with the non-migrated cells is scraped with a rubber policeman. The filters are fixed with methanol and stained with a Giemsa solution (Diff-Quick, Baxter, McGraw Park, IL). Migration is quantified by counting cells of three random high-power fields (40x) in each well, and all groups are performed in quadruplicate.

25 The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.


Example 40: Stimulation of Nitric Oxide Production by Endothelial Cells

30 Nitric oxide released by the vascular endothelium is believed to be a mediator of vascular endothelium relaxation. Thus, activity of a polypeptide of the invention can be

assayed by determining nitric oxide production by endothelial cells in response to the polypeptide.

Nitric oxide is measured in 96-well plates of confluent microvascular endothelial cells after 24 hours starvation and a subsequent 4 hr exposure to various levels of a positive control (such as VEGF-1) and the polypeptide of the invention. Nitric oxide in the medium is determined by use of the Griess reagent to measure total nitrite after reduction of nitric oxide-derived nitrate by nitrate reductase. The effect of the polypeptide of the invention on nitric oxide release is examined on HUVEC.

Briefly, NO release from cultured HUVEC monolayer is measured with a NO-specific polarographic electrode connected to a NO meter (Iso-NO, World Precision Instruments Inc.) (1049). Calibration of the NO elements is performed according to the following equation:



The standard calibration curve is obtained by adding graded concentrations of KNO₂ (0, 5, 10, 25, 50, 100, 250, and 500 nmol/L) into the calibration solution containing KI and H₂SO₄. The specificity of the Iso-NO electrode to NO is previously determined by measurement of NO from authentic NO gas (1050). The culture medium is removed and HUVECs are washed twice with Dulbecco's phosphate buffered saline. The cells are then bathed in 5 ml of filtered Krebs-Henseleit solution in 6-well plates, and the cell plates are kept on a slide warmer (Lab Line Instruments Inc.) To maintain the temperature at 37°C. The NO sensor probe is inserted vertically into the wells, keeping the tip of the electrode 2 mm under the surface of the solution, before addition of the different conditions. S-nitroso acetyl penicillamin (SNAP) is used as a positive control. The amount of released NO is expressed as picomoles per 1x10⁶ endothelial cells. All values reported are means of four to six measurements in each group (number of cell culture wells). See, Leak *et al. Biochem. and Biophys. Res. Comm.* 217:96-105 (1995).

The studies described in this example tested activity of polypeptides of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 41: Effect of Polypeptides of the Invention on Cord Formation in Angiogenesis

Another step in angiogenesis is cord formation, marked by differentiation of endothelial cells. This bioassay measures the ability of microvascular endothelial cells to form capillary-like structures (hollow structures) when cultured *in vitro*.

CADMEC (microvascular endothelial cells) are purchased from Cell Applications, Inc. as proliferating (passage 2) cells and are cultured in Cell Applications' CADMEC Growth Medium and used at passage 5. For the *in vitro* angiogenesis assay, the wells of a 48-well cell culture plate are coated with Cell Applications' Attachment Factor Medium (200 ml/well) for 30 min. at 37°C. CADMEC are seeded onto the coated wells at 7,500 cells/well and cultured overnight in Growth Medium. The Growth Medium is then replaced with 300 mg Cell Applications' Chord Formation Medium containing control buffer or a polypeptide of the invention (0.1 to 100 ng/ml) and the cells are cultured for an additional 48 hr. The numbers and lengths of the capillary-like chords are quantitated through use of the Boeckeler VIA-170 video image analyzer. All assays are done in triplicate.

Commercial (R&D) VEGF (50 ng/ml) is used as a positive control. b-esteradiol (1 ng/ml) is used as a negative control. The appropriate buffer (without protein) is also utilized as a control.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 42: Angiogenic Effect on Chick Chorioallantoic Membrane

Chick chorioallantoic membrane (CAM) is a well-established system to examine angiogenesis. Blood vessel formation on CAM is easily visible and quantifiable. The ability of polypeptides of the invention to stimulate angiogenesis in CAM can be examined.

Fertilized eggs of the White Leghorn chick (*Gallus gallus*) and the Japanese quail (*Coturnix coturnix*) are incubated at 37.8°C and 80% humidity. Differentiated CAM of 16-day-old chick and 13-day-old quail embryos is studied with the following methods.

On Day 4 of development, a window is made into the egg shell of chick eggs. The embryos are checked for normal development and the eggs sealed with cello tape. They are further incubated until Day 13. Thermanox coverslips (Nunc, Naperville, IL) are cut into disks of about 5 mm in diameter. Sterile and salt-free growth factors are dissolved in distilled water and about 3.3 mg/ 5 ml are pipetted on the disks. After air-drying, the inverted disks are applied on CAM. After 3 days, the specimens are fixed in 3% glutaraldehyde and 2% formaldehyde and rinsed in 0.12 M sodium cacodylate buffer. They are photographed with a stereo microscope [Wild M8] and embedded for semi- and ultrathin sectioning as described above. Controls are performed with carrier disks alone.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 43: Angiogenesis Assay Using a Matrigel Implant in Mouse

In vivo angiogenesis assay of a polypeptide of the invention measures the ability of an existing capillary network to form new vessels in an implanted capsule of murine extracellular matrix material (Matrigel). The protein is mixed with the liquid Matrigel at 4 degree C and the mixture is then injected subcutaneously in mice where it solidifies. After 7 days, the solid "plug" of Matrigel is removed and examined for the presence of new blood vessels. Matrigel is purchased from Becton Dickinson Labware/Collaborative Biomedical Products.

When thawed at 4 degree C the Matrigel material is a liquid. The Matrigel is mixed with a polypeptide of the invention at 150 ng/ml at 4 degrees C and drawn into cold 3 ml syringes. Female C57Bl/6 mice approximately 8 weeks old are injected with the mixture of Matrigel and experimental protein at 2 sites at the midventral aspect of the abdomen (0.5 ml/site). After 7 days, the mice are sacrificed by cervical dislocation, the

Matrigel plugs are removed and cleaned (i.e., all clinging membranes and fibrous tissue is removed). Replicate whole plugs are fixed in neutral buffered 10% formaldehyde, embedded in paraffin and used to produce sections for histological examination after staining with Masson's Trichrome. Cross sections from 3 different regions of each plug
5 are processed. Selected sections are stained for the presence of vWF. The positive control for this assay is bovine basic FGF (150 ng/ml). Matrigel alone is used to determine basal levels of angiogenesis.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to
10 test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 44: Rescue of Ischemia in Rabbit Lower Limb Model

15 To study the in vivo effects of polynucleotides and polypeptides of the invention on ischemia, a rabbit hindlimb ischemia model is created by surgical removal of one femoral arteries as described previously (Takeshita *et al.*, *Am J. Pathol* 147:1649-1660 (1995)). The excision of the femoral artery results in retrograde propagation of thrombus and occlusion of the external iliac artery. Consequently, blood flow to the ischemic limb
20 is dependent upon collateral vessels originating from the internal iliac artery (Takeshita *et al.* *Am J. Pathol* 147:1649-1660 (1995)). An interval of 10 days is allowed for post-operative recovery of rabbits and development of endogenous collateral vessels. At 10 day post-operatively (day 0), after performing a baseline angiogram, the internal iliac artery of the ischemic limb is transfected with 500 mg naked expression plasmid
25 containing a polynucleotide of the invention by arterial gene transfer technology using a hydrogel-coated balloon catheter as described (Riessen *et al.* *Hum Gene Ther.* 4:749-758 (1993); Leclerc *et al.* *J. Clin. Invest.* 90: 936-944 (1992)). When a polypeptide of the invention is used in the treatment, a single bolus of 500 mg polypeptide of the invention or control is delivered into the internal iliac artery of the ischemic limb over a period of 1
30 min. through an infusion catheter. On day 30, various parameters are measured in these rabbits: (a) BP ratio - The blood pressure ratio of systolic pressure of the ischemic limb to that of normal limb; (b) Blood Flow and Flow Reserve - Resting FL: the blood flow

during undilated condition and Max FL: the blood flow during fully dilated condition (also an indirect measure of the blood vessel amount) and Flow Reserve is reflected by the ratio of max FL: resting FL; (c) Angiographic Score - This is measured by the angiogram of collateral vessels. A score is determined by the percentage of circles in an overlaying grid that with crossing opacified arteries divided by the total number in the rabbit thigh; (d) Capillary density - The number of collateral capillaries determined in light microscopic sections taken from hindlimbs.

The studies described in this example tested activity of polynucleotides and polypeptides of the invention. However, one skilled in the art could easily modify the exemplified studies to test the agonists, and/or antagonists of the invention.

Example 45: Effect of Polypeptides of the Invention on Vasodilation

Since dilation of vascular endothelium is important in reducing blood pressure, the ability of polypeptides of the invention to affect the blood pressure in spontaneously hypertensive rats (SHR) is examined. Increasing doses (0, 10, 30, 100, 300, and 900 mg/kg) of the polypeptides of the invention are administered to 13-14 week old spontaneously hypertensive rats (SHR). Data are expressed as the mean +/- SEM. Statistical analysis are performed with a paired t-test and statistical significance is defined as $p < 0.05$ vs. the response to buffer alone.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 46: Rat Ischemic Skin Flap Model

The evaluation parameters include skin blood flow, skin temperature, and factor VIII immunohistochemistry or endothelial alkaline phosphatase reaction. Expression of polypeptides of the invention, during the skin ischemia, is studied using in situ hybridization.

The study in this model is divided into three parts as follows:

- a) Ischemic skin
- b) Ischemic skin wounds
- c) Normal wounds

The experimental protocol includes:

- 5 a) Raising a 3x4 cm, single pedicle full-thickness random skin flap (myocutaneous flap over the lower back of the animal).
- b) An excisional wounding (4-6 mm in diameter) in the ischemic skin (skin-flap).
- c) Topical treatment with a polypeptide of the invention of the excisional wounds (day 0, 1, 2, 3, 4 post-wounding) at the following various dosage ranges: 1mg to 100 mg.
- 10 d) Harvesting the wound tissues at day 3, 5, 7, 10, 14 and 21 post-wounding for histological, immunohistochemical, and in situ studies.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the
15 invention.

Example 47: Peripheral Arterial Disease Model

Angiogenic therapy using a polypeptide of the invention is a novel therapeutic
20 strategy to obtain restoration of blood flow around the ischemia in case of peripheral arterial diseases. The experimental protocol includes:

- a) One side of the femoral artery is ligated to create ischemic muscle of the hindlimb, the other side of hindlimb serves as a control.
- b) a polypeptide of the invention, in a dosage range of 20 mg - 500 mg, is
25 delivered intravenously and/or intramuscularly 3 times (perhaps more) per week for 2-3 weeks.
- c) The ischemic muscle tissue is collected after ligation of the femoral artery at 1, 2, and 3 weeks for the analysis of expression of a polypeptide of the invention and histology. Biopsy is also performed on the other side of normal muscle of the
30 contralateral hindlimb.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to

test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 48: Ischemic Myocardial Disease Model

5

A polypeptide of the invention is evaluated as a potent mitogen capable of stimulating the development of collateral vessels, and restructuring new vessels after coronary artery occlusion. Alteration of expression of the polypeptide is investigated in situ. The experimental protocol includes:

10 a) The heart is exposed through a left-side thoracotomy in the rat. Immediately, the left coronary artery is occluded with a thin suture (6-0) and the thorax is closed.

 b) a polypeptide of the invention, in a dosage range of 20 mg - 500 mg, is delivered intravenously and/or intramuscularly 3 times (perhaps more) per week for 2-4 weeks.

15 c) Thirty days after the surgery, the heart is removed and cross-sectioned for morphometric and in situ analyzes.

 The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the
20 invention.

Example 49: Rat Corneal Wound Healing Model

 This animal model shows the effect of a polypeptide of the invention on
25 neovascularization. The experimental protocol includes:

 a) Making a 1-1.5 mm long incision from the center of cornea into the stromal layer.

 b) Inserting a spatula below the lip of the incision facing the outer corner of the eye.

30 c) Making a pocket (its base is 1-1.5 mm from the edge of the eye).

 d) Positioning a pellet, containing 50ng- 5ug of a polypeptide of the invention, within the pocket.

e) Treatment with a polypeptide of the invention can also be applied topically to the corneal wounds in a dosage range of 20mg - 500mg (daily treatment for five days).

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 50: Diabetic Mouse and Glucocorticoid-Impaired Wound Healing

Models

10

A. Diabetic db+/db+ Mouse Model.

To demonstrate that a polypeptide of the invention accelerates the healing process, the genetically diabetic mouse model of wound healing is used. The full thickness wound healing model in the db+/db+ mouse is a well characterized, clinically relevant and reproducible model of impaired wound healing. Healing of the diabetic wound is dependent on formation of granulation tissue and re-epithelialization rather than contraction (Gartner, M.H. *et al.*, *J. Surg. Res.* 52:389 (1992); Greenhalgh, D.G. *et al.*, *Am. J. Pathol.* 136:1235 (1990)).

The diabetic animals have many of the characteristic features observed in Type II diabetes mellitus. Homozygous (db+/db+) mice are obese in comparison to their normal heterozygous (db+/+m) littermates. Mutant diabetic (db+/db+) mice have a single autosomal recessive mutation on chromosome 4 (db+) (Coleman *et al.* *Proc. Natl. Acad. Sci. USA* 77:283-293 (1982)). Animals show polyphagia, polydipsia and polyuria. Mutant diabetic mice (db+/db+) have elevated blood glucose, increased or normal insulin levels, and suppressed cell-mediated immunity (Mandel *et al.*, *J. Immunol.* 120:1375 (1978); Debray-Sachs, M. *et al.*, *Clin. Exp. Immunol.* 51(1):1-7 (1983); Leiter *et al.*, *Am. J. of Pathol.* 114:46-55 (1985)). Peripheral neuropathy, myocardial complications, and microvascular lesions, basement membrane thickening and glomerular filtration abnormalities have been described in these animals (Norido, F. *et al.*, *Exp. Neurol.* 83(2):221-232 (1984); Robertson *et al.*, *Diabetes* 29(1):60-67 (1980); Giacomelli *et al.*, *Lab Invest.* 40(4):460-473 (1979); Coleman, D.L., *Diabetes* 31 (Suppl):1-6 (1982)). These

homozygous diabetic mice develop hyperglycemia that is resistant to insulin analogous to human type II diabetes (Mandel *et al.*, *J. Immunol.* 120:1375-1377 (1978)).

The characteristics observed in these animals suggests that healing in this model may be similar to the healing observed in human diabetes (Greenhalgh, *et al.*, *Am. J. of Pathol.* 136:1235-1246 (1990)).

Genetically diabetic female C57BL/KsJ (db+/db+) mice and their non-diabetic (db+/+m) heterozygous littermates are used in this study (Jackson Laboratories). The animals are purchased at 6 weeks of age and are 8 weeks old at the beginning of the study. Animals are individually housed and received food and water ad libitum. All manipulations are performed using aseptic techniques. The experiments are conducted according to the rules and guidelines of Human Genome Sciences, Inc. Institutional Animal Care and Use Committee and the Guidelines for the Care and Use of Laboratory Animals.

Wounding protocol is performed according to previously reported methods (Tsuboi, R. and Rifkin, D.B., *J. Exp. Med.* 172:245-251 (1990)). Briefly, on the day of wounding, animals are anesthetized with an intraperitoneal injection of Avertin (0.01 mg/mL), 2,2,2-tribromoethanol and 2-methyl-2-butanol dissolved in deionized water. The dorsal region of the animal is shaved and the skin washed with 70% ethanol solution and iodine. The surgical area is dried with sterile gauze prior to wounding. An 8 mm full-thickness wound is then created using a Keyes tissue punch. Immediately following wounding, the surrounding skin is gently stretched to eliminate wound expansion. The wounds are left open for the duration of the experiment. Application of the treatment is given topically for 5 consecutive days commencing on the day of wounding. Prior to treatment, wounds are gently cleansed with sterile saline and gauze sponges.

Wounds are visually examined and photographed at a fixed distance at the day of surgery and at two day intervals thereafter. Wound closure is determined by daily measurement on days 1-5 and on day 8. Wounds are measured horizontally and vertically using a calibrated Jameson caliper. Wounds are considered healed if granulation tissue is no longer visible and the wound is covered by a continuous epithelium.

A polypeptide of the invention is administered using at a range different doses, from 4mg to 500mg per wound per day for 8 days in vehicle. Vehicle control groups received 50mL of vehicle solution.

Animals are euthanized on day 8 with an intraperitoneal injection of sodium pentobarbital (300mg/kg). The wounds and surrounding skin are then harvested for histology and immunohistochemistry. Tissue specimens are placed in 10% neutral buffered formalin in tissue cassettes between biopsy sponges for further processing.

- 5 Three groups of 10 animals each (5 diabetic and 5 non-diabetic controls) are evaluated: 1) Vehicle placebo control, 2) untreated group, and 3) treated group.

Wound closure is analyzed by measuring the area in the vertical and horizontal axis and obtaining the total square area of the wound. Contraction is then estimated by establishing the differences between the initial wound area (day 0) and that of post
10 treatment (day 8). The wound area on day 1 is 64mm², the corresponding size of the dermal punch. Calculations are made using the following formula:

$$[\text{Open area on day 8}] - [\text{Open area on day 1}] / [\text{Open area on day 1}]$$

- 15 Specimens are fixed in 10% buffered formalin and paraffin embedded blocks are sectioned perpendicular to the wound surface (5mm) and cut using a Reichert-Jung microtome. Routine hematoxylin-eosin (H&E) staining is performed on cross-sections of bisected wounds. Histologic examination of the wounds are used to assess whether the healing process and the morphologic appearance of the repaired skin is altered by
20 treatment with a polypeptide of the invention. This assessment included verification of the presence of cell accumulation, inflammatory cells, capillaries, fibroblasts, re-epithelialization and epidermal maturity (Greenhalgh, D.G. *et al.*, *Am. J. Pathol.* 136:1235 (1990)). A calibrated lens micrometer is used by a blinded observer.

- 25 Tissue sections are also stained immunohistochemically with a polyclonal rabbit anti-human keratin antibody using ABC Elite detection system. Human skin is used as a positive tissue control while non-immune IgG is used as a negative control. Keratinocyte growth is determined by evaluating the extent of reepithelialization of the wound using a calibrated lens micrometer.

- 30 Proliferating cell nuclear antigen/cyclin (PCNA) in skin specimens is demonstrated by using anti-PCNA antibody (1:50) with an ABC Elite detection system. Human colon cancer can serve as a positive tissue control and human brain tissue can be used as a negative tissue control. Each specimen includes a section with omission of the primary

antibody and substitution with non-immune mouse IgG. Ranking of these sections is based on the extent of proliferation on a scale of 0-8, the lower side of the scale reflecting slight proliferation to the higher side reflecting intense proliferation.

Experimental data are analyzed using an unpaired t test. A p value of < 0.05 is
5 considered significant.

B. Steroid Impaired Rat Model

The inhibition of wound healing by steroids has been well documented in various *in vitro* and *in vivo* systems (Wahl, Glucocorticoids and Wound healing. In: Anti-
10 Inflammatory Steroid Action: Basic and Clinical Aspects. 280-302 (1989); Wahl *et al.*, *J. Immunol.* 115: 476-481 (1975); Werb *et al.*, *J. Exp. Med.* 147:1684-1694 (1978)). Glucocorticoids retard wound healing by inhibiting angiogenesis, decreasing vascular permeability (Ebert *et al.*, *Am. Intern. Med.* 37:701-705 (1952)), fibroblast proliferation, and collagen synthesis (Beck *et al.*, *Growth Factors.* 5: 295-304 (1991); Haynes *et al.*,
15 *J. Clin. Invest.* 61: 703-797 (1978)) and producing a transient reduction of circulating monocytes (Haynes *et al.*, *J. Clin. Invest.* 61: 703-797 (1978); Wahl, "Glucocorticoids and wound healing", In: Antiinflammatory Steroid Action: Basic and Clinical Aspects, Academic Press, New York, pp. 280-302 (1989)). The systemic administration of steroids to impaired wound healing is a well establish phenomenon in rats (Beck *et al.*, *Growth*
20 *Factors.* 5: 295-304 (1991); Haynes *et al.*, *J. Clin. Invest.* 61: 703-797 (1978); Wahl, "Glucocorticoids and wound healing", In: Antiinflammatory Steroid Action: Basic and Clinical Aspects, Academic Press, New York, pp. 280-302 (1989); Pierce *et al.*, *Proc. Natl. Acad. Sci. USA* 86: 2229-2233 (1989)).

To demonstrate that a polypeptide of the invention can accelerate the healing
25 process, the effects of multiple topical applications of the polypeptide on full thickness excisional skin wounds in rats in which healing has been impaired by the systemic administration of methylprednisolone is assessed.

Young adult male Sprague Dawley rats weighing 250-300 g (Charles River Laboratories) are used in this example. The animals are purchased at 8 weeks of age and
30 are 9 weeks old at the beginning of the study. The healing response of rats is impaired by the systemic administration of methylprednisolone (17mg/kg/rat intramuscularly) at the time of wounding. Animals are individually housed and received food and water *ad*

libitum. All manipulations are performed using aseptic techniques. This study is conducted according to the rules and guidelines of Human Genome Sciences, Inc. Institutional Animal Care and Use Committee and the Guidelines for the Care and Use of Laboratory Animals.

5 The wounding protocol is followed according to section A, above. On the day of wounding, animals are anesthetized with an intramuscular injection of ketamine (50 mg/kg) and xylazine (5 mg/kg). The dorsal region of the animal is shaved and the skin washed with 70% ethanol and iodine solutions. The surgical area is dried with sterile gauze prior to wounding. An 8 mm full-thickness wound is created using a Keyes tissue
10 punch. The wounds are left open for the duration of the experiment. Applications of the testing materials are given topically once a day for 7 consecutive days commencing on the day of wounding and subsequent to methylprednisolone administration. Prior to treatment, wounds are gently cleansed with sterile saline and gauze sponges.

 Wounds are visually examined and photographed at a fixed distance at the day of
15 wounding and at the end of treatment. Wound closure is determined by daily measurement on days 1-5 and on day 8. Wounds are measured horizontally and vertically using a calibrated Jameson caliper. Wounds are considered healed if granulation tissue is no longer visible and the wound is covered by a continuous epithelium.

 The polypeptide of the invention is administered using at a range different doses,
20 from 4mg to 500mg per wound per day for 8 days in vehicle. Vehicle control groups received 50mL of vehicle solution.

 Animals are euthanized on day 8 with an intraperitoneal injection of sodium pentobarbital (300mg/kg). The wounds and surrounding skin are then harvested for histology. Tissue specimens are placed in 10% neutral buffered formalin in tissue
25 cassettes between biopsy sponges for further processing.

 Four groups of 10 animals each (5 with methylprednisolone and 5 without glucocorticoid) are evaluated: 1) Untreated group 2) Vehicle placebo control 3) treated groups.

 Wound closure is analyzed by measuring the area in the vertical and horizontal
30 axis and obtaining the total area of the wound. Closure is then estimated by establishing the differences between the initial wound area (day 0) and that of post treatment (day 8).

The wound area on day 1 is 64mm^2 , the corresponding size of the dermal punch. Calculations are made using the following formula:

$$[\text{Open area on day 8}] - [\text{Open area on day 1}] / [\text{Open area on day 1}]$$

5

Specimens are fixed in 10% buffered formalin and paraffin embedded blocks are sectioned perpendicular to the wound surface (5mm) and cut using an Olympus microtome. Routine hematoxylin-eosin (H&E) staining is performed on cross-sections of bisected wounds. Histologic examination of the wounds allows assessment of whether the healing process and the morphologic appearance of the repaired skin is improved by treatment with a polypeptide of the invention. A calibrated lens micrometer is used by a blinded observer to determine the distance of the wound gap.

Experimental data are analyzed using an unpaired t test. A p value of < 0.05 is considered significant.

15 The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

20 **Example 51: Lymphadema Animal Model**

or The purpose of this experimental approach is to create an appropriate and consistent lymphedema model for testing the therapeutic effects of a polypeptide of the invention in lymphangiogenesis and re-establishment of the lymphatic circulatory system in the rat hind limb. Effectiveness is measured by swelling volume of the affected limb, quantification of the amount of lymphatic vasculature, total blood plasma protein, and histopathology. Acute lymphedema is observed for 7-10 days. Perhaps more importantly, the chronic progress of the edema is followed for up to 3-4 weeks.

25 Prior to beginning surgery, blood sample is drawn for protein concentration analysis. Male rats weighing approximately ~350g are dosed with Pentobarbital. Subsequently, the right legs are shaved from knee to hip. The shaved area is swabbed with gauze soaked in 70% EtOH. Blood is drawn for serum total protein testing.

Circumference and volumetric measurements are made prior to injecting dye into paws after marking 2 measurement levels (0.5 cm above heel, at mid-pt of dorsal paw). The intradermal dorsum of both right and left paws are injected with 0.05 ml of 1% Evan's Blue. Circumference and volumetric measurements are then made following injection of
5 dye into paws.

Using the knee joint as a landmark, a mid-leg inguinal incision is made circumferentially allowing the femoral vessels to be located. Forceps and hemostats are used to dissect and separate the skin flaps. After locating the femoral vessels, the lymphatic vessel that runs along side and underneath the vessel(s) is located. The main
10 lymphatic vessels in this area are then electrically coagulated suture ligated.

Using a microscope, muscles in back of the leg (near the semitendinosus and adductors) are bluntly dissected. The popliteal lymph node is then located. The 2 proximal and 2 distal lymphatic vessels and distal blood supply of the popliteal node are then and ligated by suturing. The popliteal lymph node, and any accompanying adipose
15 tissue, is then removed by cutting connective tissues.

Care is taken to control any mild bleeding resulting from this procedure. After lymphatics are occluded, the skin flaps are sealed by using liquid skin (Vetbond) (AJ Buck). The separated skin edges are sealed to the underlying muscle tissue while leaving a gap of ~0.5 cm around the leg. Skin also may be anchored by suturing to underlying
20 muscle when necessary.

To avoid infection, animals are housed individually with mesh (no bedding). Recovering animals are checked daily through the optimal edematous peak, which typically occurred by day 5-7. The plateau edematous peak are then observed. To evaluate the intensity of the lymphedema, the circumference and volumes of 2 designated
25 places on each paw before operation and daily for 7 days are measured. The effect plasma proteins on lymphedema is determined and whether protein analysis is a useful testing perimeter is also investigated. The weights of both control and edematous limbs are evaluated at 2 places. Analysis is performed in a blind manner.

Circumference Measurements: Under brief gas anesthetic to prevent limb
30 movement, a cloth tape is used to measure limb circumference. Measurements are done at the ankle bone and dorsal paw by 2 different people then those 2 readings are averaged. Readings are taken from both control and edematous limbs.

Volumetric Measurements: On the day of surgery, animals are anesthetized with Pentobarbital and are tested prior to surgery. For daily volumetrics animals are under brief halothane anesthetic (rapid immobilization and quick recovery), both legs are shaved and equally marked using waterproof marker on legs. Legs are first dipped in water, then
5 dipped into instrument to each marked level then measured by Buxco edema software(Chen/Victor). Data is recorded by one person, while the other is dipping the limb to marked area.

Blood-plasma protein measurements: Blood is drawn, spun, and serum separated prior to surgery and then at conclusion for total protein and Ca²⁺ comparison.

10 Limb Weight Comparison: After drawing blood, the animal is prepared for tissue collection. The limbs are amputated using a quillitine, then both experimental and control legs are cut at the ligature and weighed. A second weighing is done as the tibio-cacaneal joint is disarticulated and the foot is weighed.

Histological Preparations: The transverse muscle located behind the knee
15 (popliteal) area is dissected and arranged in a metal mold, filled with freezeGel, dipped into cold methylbutane, placed into labeled sample bags at - 80EC until sectioning. Upon sectioning, the muscle is observed under fluorescent microscopy for lymphatics..

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to
20 test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

**Example 52: Suppression of TNF alpha-induced adhesion molecule expression
by a Polypeptide of the Invention**

25 The recruitment of lymphocytes to areas of inflammation and angiogenesis involves specific receptor-ligand interactions between cell surface adhesion molecules (CAMs) on lymphocytes and the vascular endothelium. The adhesion process, in both normal and pathological settings, follows a multi-step cascade that involves intercellular adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule-1 (VCAM-1), and
30 endothelial leukocyte adhesion molecule-1 (E-selectin) expression on endothelial cells (EC). The expression of these molecules and others on the vascular endothelium determines the efficiency with which leukocytes may adhere to the local vasculature and

extravasate into the local tissue during the development of an inflammatory response. The local concentration of cytokines and growth factor participate in the modulation of the expression of these CAMs.

5 Tumor necrosis factor alpha (TNF- α), a potent proinflammatory cytokine, is a stimulator of all three CAMs on endothelial cells and may be involved in a wide variety of inflammatory responses, often resulting in a pathological outcome.

The potential of a polypeptide of the invention to mediate a suppression of TNF- α induced CAM expression can be examined. A modified ELISA assay which uses ECs as a solid phase absorbent is employed to measure the amount of CAM expression on TNF- α treated ECs when co-stimulated with a member of the FGF family of proteins.

To perform the experiment, human umbilical vein endothelial cell (HUVEC) cultures are obtained from pooled cord harvests and maintained in growth medium (EGM-2; Clonetics, San Diego, CA) supplemented with 10% FCS and 1% penicillin/streptomycin in a 37 degree C humidified incubator containing 5% CO₂.

15 HUVECs are seeded in 96-well plates at concentrations of 1×10^4 cells/well in EGM medium at 37 degree C for 18-24 hrs or until confluent. The monolayers are subsequently washed 3 times with a serum-free solution of RPMI-1640 supplemented with 100 U/ml penicillin and 100 mg/ml streptomycin, and treated with a given cytokine and/or growth factor(s) for 24 h at 37 degree C. Following incubation, the cells are then evaluated for

20 CAM expression.

Human Umbilical Vein Endothelial cells (HUVECs) are grown in a standard 96 well plate to confluence. Growth medium is removed from the cells and replaced with 90 μ l of 199 Medium (10% FBS). Samples for testing and positive or negative controls are added to the plate in triplicate (in 10 μ l volumes). Plates are incubated at 37 degree C for

25 either 5 h (selectin and integrin expression) or 24 h (integrin expression only). Plates are aspirated to remove medium and 100 μ l of 0.1% paraformaldehyde-PBS(with Ca⁺⁺ and Mg⁺⁺) is added to each well. Plates are held at 4°C for 30 min.

Fixative is then removed from the wells and wells are washed 1X with PBS(+Ca,Mg)+0.5% BSA and drained. Do not allow the wells to dry. Add 10 μ l of

30 diluted primary antibody to the test and control wells. Anti-ICAM-1-Biotin, Anti-VCAM-1-Biotin and Anti-E-selectin-Biotin are used at a concentration of 10 μ g/ml (1:10 dilution

of 0.1 mg/ml stock antibody). Cells are incubated at 37°C for 30 min. in a humidified environment. Wells are washed X3 with PBS(+Ca,Mg)+0.5% BSA.

Then add 20 µl of diluted ExtrAvidin-Alkaline Phosphatase (1:5,000 dilution) to each well and incubated at 37°C for 30 min. Wells are washed X3 with
5 PBS(+Ca,Mg)+0.5% BSA. 1 tablet of p-Nitrophenol Phosphate pNPP is dissolved in 5 ml of glycine buffer (pH 10.4). 100 µl of pNPP substrate in glycine buffer is added to each test well. Standard wells in triplicate are prepared from the working dilution of the ExtrAvidin-Alkaline Phosphatase in glycine buffer: $1:5,000 (10^0) > 10^{-0.5} > 10^{-1} > 10^{-1.5}$. 5 µl of each dilution is added to triplicate wells and the resulting AP content in each well is
10 5.50 ng, 1.74 ng, 0.55 ng, 0.18 ng. 100 µl of pNPP reagent must then be added to each of the standard wells. The plate must be incubated at 37°C for 4h. A volume of 50 µl of 3M NaOH is added to all wells. The results are quantified on a plate reader at 405 nm. The background subtraction option is used on blank wells filled with glycine buffer only. The template is set up to indicate the concentration of AP-conjugate in each standard well [
15 5.50 ng; 1.74 ng; 0.55 ng; 0.18 ng]. Results are indicated as amount of bound AP-conjugate in each sample.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the
20 invention.

Example 53: Assay for the Stimulation of Bone Marrow CD34+ Cell Proliferation

This assay is based on the ability of human CD34+ to proliferate in the
25 presence of hematopoietic growth factors and evaluates the ability of isolated polypeptides expressed in mammalian cells to stimulate proliferation of CD34+ cells.

It has been previously shown that most mature precursors will respond to only a single signal. More immature precursors require at least two signals to respond. Therefore, to test the effect of polypeptides on hematopoietic activity of a wide range
30 of progenitor cells, the assay contains a given polypeptide in the presence or absence of other hematopoietic growth factors. Isolated cells are cultured for 5 days in the presence of Stem Cell Factor (SCF) in combination with tested sample. SCF alone

has a very limited effect on the proliferation of bone marrow (BM) cells, acting in such conditions only as a "survival" factor. However, combined with any factor exhibiting stimulatory effect on these cells (e.g., IL-3), SCF will cause a synergistic effect. Therefore, if the tested polypeptide has a stimulatory effect on a hematopoietic progenitors, such activity can be easily detected. Since normal BM cells have a low level of cycling cells, it is likely that any inhibitory effect of a given polypeptide, or agonists or antagonists thereof, might not be detected. Accordingly, assays for an inhibitory effect on progenitors is preferably tested in cells that are first subjected to *in vitro* stimulation with SCF+IL+3, and then contacted with the compound that is being evaluated for inhibition of such induced proliferation.

Briefly, CD34+ cells are isolated using methods known in the art. The cells are thawed and resuspended in medium (QBSF 60 serum-free medium with 1% L-glutamine (500ml) Quality Biological, Inc., Gaithersburg, MD Cat# 160-204-101). After several gentle centrifugation steps at 200 x g, cells are allowed to rest for one hour. The cell count is adjusted to 2.5×10^5 cells/ml. During this time, 100 μ l of sterile water is added to the peripheral wells of a 96-well plate. The cytokines that can be tested with a given polypeptide in this assay is rhSCF (R&D Systems, Minneapolis, MN, Cat# 255-SC) at 50 ng/ml alone and in combination with rhSCF and rhIL-3 (R&D Systems, Minneapolis, MN, Cat# 203-ML) at 30 ng/ml. After one hour, 10 μ l of prepared cytokines, 50 μ l SID (supernatants at 1:2 dilution = 50 μ l) and 20 μ l of diluted cells are added to the media which is already present in the wells to allow for a final total volume of 100 μ l. The plates are then placed in a 37°C/5% CO₂ incubator for five days.

Eighteen hours before the assay is harvested, 0.5 μ Ci/well of [3H] Thymidine is added in a 10 μ l volume to each well to determine the proliferation rate. The experiment is terminated by harvesting the cells from each 96-well plate to a filtermat using the Tomtec Harvester 96. After harvesting, the filtermats are dried, trimmed and placed into OmniFilter assemblies consisting of one OmniFilter plate and one OmniFilter Tray. 60 μ l Microscint is added to each well and the plate sealed with TopSeal-A press-on sealing film. A bar code 15 sticker is affixed to the first plate for counting. The sealed plates is then loaded and the level of radioactivity determined

via the Packard Top Count and the printed data collected for analysis. The level of radioactivity reflects the amount of cell proliferation.

The studies described in this example test the activity of a given polypeptide to stimulate bone marrow CD34+ cell proliferation. One skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), antibodies, agonists, and/or antagonists and fragments and variants thereof. As a nonlimiting example, potential antagonists tested in this assay would be expected to inhibit cell proliferation in the presence of cytokines and/or to increase the inhibition of cell proliferation in the presence of cytokines and a given polypeptide. In contrast, potential agonists tested in this assay would be expected to enhance cell proliferation and/or to decrease the inhibition of cell proliferation in the presence of cytokines and a given polypeptide.

The ability of a gene to stimulate the proliferation of bone marrow CD34+ cells indicates that polynucleotides and polypeptides corresponding to the gene are useful for the diagnosis and treatment of disorders affecting the immune system and hematopoiesis. Representative uses are described in the "Immune Activity" and "Infectious Disease" sections above, and elsewhere herein.

Example 54: Assay for Extracellular Matrix Enhanced Cell Response (EMECCR)

The objective of the Extracellular Matrix Enhanced Cell Response (EMECCR) assay is to identify gene products (e.g., isolated polypeptides) that act on the hematopoietic stem cells in the context of the extracellular matrix (ECM) induced signal.

Cells respond to the regulatory factors in the context of signal(s) received from the surrounding microenvironment. For example, fibroblasts, and endothelial and epithelial stem cells fail to replicate in the absence of signals from the ECM. Hematopoietic stem cells can undergo self-renewal in the bone marrow, but not in *in vitro* suspension culture. The ability of stem cells to undergo self-renewal *in vitro* is dependent upon their interaction with the stromal cells and the ECM protein fibronectin (fn). Adhesion of cells to fn is mediated by the $\alpha_5\beta_1$ and $\alpha_4\beta_1$ integrin receptors, which are expressed by human and mouse hematopoietic stem cells. The factor(s) which integrate with the ECM environment and responsible for stimulating

stem cell self-renewal has not yet been identified. Discovery of such factors should be of great interest in gene therapy and bone marrow transplant applications

Briefly, polystyrene, non tissue culture treated, 96-well plates are coated with
fn fragment at a coating concentration of $0.2 \mu\text{g}/\text{cm}^2$. Mouse bone marrow cells are
5 plated (1,000 cells/well) in 0.2 ml of serum-free medium. Cells cultured in the
presence of IL-3 (5 ng/ml) + SCF (50 ng/ml) would serve as the positive control,
conditions under which little self-renewal but pronounced differentiation of the stem
cells is to be expected. Gene products are tested with appropriate negative controls in
the presence and absence of SCF(5.0 ng/ml), where test factor supernates represent
10 10% of the total assay volume. The plated cells are then allowed to grow by
incubating in a low oxygen environment (5% CO_2 , 7% O_2 , and 88% N_2) tissue
culture incubator for 7 days. The number of proliferating cells within the wells is
then quantitated by measuring thymidine incorporation into cellular DNA.
Verification of the positive hits in the assay will require phenotypic characterization
15 of the cells, which can be accomplished by scaling up of the culture system and using
appropriate antibody reagents against cell surface antigens and FACScan.

One skilled in the art could easily modify the exemplified studies to test the
activity of polynucleotides (e.g., gene therapy), antibodies, agonists, and/or
antagonists and fragments and variants thereof.

20 If a particular gene product is found to be a stimulator of hematopoietic
progenitors, polynucleotides and polypeptides corresponding to the gene may be
useful for the diagnosis and treatment of disorders affecting the immune system and
hematopoiesis. Representative uses are described in the "Immune Activity" and
"Infectious Disease" sections above, and elsewhere herein. The gene product may
25 also be useful in the expansion of stem cells and committed progenitors of various
blood lineages, and in the differentiation and/or proliferation of various cell types.

Additionally, the polynucleotides and/or polypeptides of the gene of interest
and/or agonists and/or antagonists thereof, may also be employed to inhibit the
proliferation and differentiation of hematopoietic cells and therefore may be
30 employed to protect bone marrow stem cells from chemotherapeutic agents during
chemotherapy. This antiproliferative effect may allow administration of higher doses

of chemotherapeutic agents and, therefore, more effective chemotherapeutic treatment.

Moreover, polynucleotides and polypeptides corresponding to the gene of interest may also be useful for the treatment and diagnosis of hematopoietic related disorders such as, for example, anemia, pancytopenia, leukopenia, thrombocytopenia or leukemia since stromal cells are important in the production of cells of hematopoietic lineages. The uses include bone marrow cell ex-vivo culture, bone marrow transplantation, bone marrow reconstitution, radiotherapy or chemotherapy of neoplasia.

Example 55: Human Dermal Fibroblast and Aortic Smooth Muscle Cell Proliferation

The polypeptide of interest is added to cultures of normal human dermal fibroblasts (NHDF) and human aortic smooth muscle cells (AoSMC) and two co-assays are performed with each sample. The first assay examines the effect of the polypeptide of interest on the proliferation of normal human dermal fibroblasts (NHDF) or aortic smooth muscle cells (AoSMC). Aberrant growth of fibroblasts or smooth muscle cells is a part of several pathological processes, including fibrosis, and restenosis. The second assay examines IL6 production by both NHDF and SMC. IL6 production is an indication of functional activation. Activated cells will have increased production of a number of cytokines and other factors, which can result in a proinflammatory or immunomodulatory outcome. Assays are run with and without co-TNF α stimulation, in order to check for costimulatory or inhibitory activity.

Briefly, on day 1, 96-well black plates are set up with 1000 cells/well (NHDF) or 2000 cells/well (AoSMC) in 100 μ l culture media. NHDF culture media contains: Clonetics FB basal media, 1mg/ml hFGF, 5mg/ml insulin, 50mg/ml gentamycin, 2%FBS, while AoSMC culture media contains Clonetics SM basal media, 0.5 μ g/ml hEGF, 5mg/ml insulin, 1 μ g/ml hFGF, 50mg/ml gentamycin, 50 μ g/ml Amphotericin B, 5%FBS. After incubation @ 37°C for at least 4-5 hours culture media is aspirated and replaced with growth arrest media. Growth arrest media for NHDF contains fibroblast basal media, 50mg/ml gentamycin, 2% FBS, while growth arrest media for

AoSMC contains SM basal media, 50mg/ml gentamycin, 50µg/ml Amphotericin B, 0.4% FBS. Incubate at 37C until day 2.

On day 2, serial dilutions and templates of the polypeptide of interest are designed which should always include media controls and known-protein controls.

- 5 For both stimulation and inhibition experiments, proteins are diluted in growth arrest media. For inhibition experiments, TNFa is added to a final concentration of 2ng/ml (NHDF) or 5ng/ml (AoSMC). Then add 1/3 vol media containing controls or supernatants and incubate at 37C/5% CO₂ until day 5.

- 10 Transfer 60µl from each well to another labeled 96-well plate, cover with a plate-sealer, and store at 4C until Day 6 (for IL6 ELISA). To the remaining 100 µl in the cell culture plate, aseptically add Alamar Blue in an amount equal to 10% of the culture volume (10µl). Return plates to incubator for 3 to 4 hours. Then measure fluorescence with excitation at 530nm and emission at 590nm using the CytoFluor. This yields the growth stimulation/inhibition data.

- 15 On day 5, the IL6 ELISA is performed by coating a 96 well plate with 50-100 ul/well of Anti-Human IL6 Monoclonal antibody diluted in PBS, pH 7.4, incubate ON at room temperature.

- On day 6, empty the plates into the sink and blot on paper towels. Prepare Assay Buffer containing PBS with 4% BSA. Block the plates with 200 µl/well of
20 Pierce Super Block blocking buffer in PBS for 1-2 hr and then wash plates with wash buffer (PBS, 0.05% Tween-20). Blot plates on paper towels. Then add 50 µl/well of diluted Anti-Human IL-6 Monoclonal, Biotin-labeled antibody at 0.50 mg/ml. Make dilutions of IL-6 stock in media (30, 10, 3, 1, 0.3, 0 ng/ml). Add duplicate samples to top row of plate. Cover the plates and incubate for 2 hours at RT on shaker.

- 25 Wash plates with wash buffer and blot on paper towels. Dilute EU-labeled Streptavidin 1:1000 in Assay buffer, and add 100 µl/well. Cover the plate and incubate 1 h at RT. Wash plates with wash buffer. Blot on paper towels.

- Add 100 µl/well of Enhancement Solution. Shake for 5 minutes. Read the plate on the Wallac DELFIA Fluorometer. Readings from triplicate samples in each
30 assay were tabulated and averaged.

A positive result in this assay suggests AoSMC cell proliferation and that the gene product of interest may be involved in dermal fibroblast proliferation and/or

smooth muscle cell proliferation. A positive result also suggests many potential uses of polypeptides, polynucleotides, agonists and/or antagonists of the gene/gene product of interest. For example, inflammation and immune responses, wound healing, and angiogenesis, as detailed throughout this specification. Particularly, polypeptides of the gene product and polynucleotides of the gene may be used in wound healing and dermal regeneration, as well as the promotion of vasculargenesis, both of the blood vessels and lymphatics. The growth of vessels can be used in the treatment of, for example, cardiovascular diseases. Additionally, antagonists of polypeptides of the gene product and polynucleotides of the gene may be useful in treating diseases, disorders, and/or conditions which involve angiogenesis by acting as an anti-vascular (e.g., anti-angiogenesis). These diseases, disorders, and/or conditions are known in the art and/or are described herein, such as, for example, malignancies, solid tumors, benign tumors, for example hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas; arteriosclerotic plaques; ocular angiogenic diseases, for example, diabetic retinopathy, retinopathy of prematurity, macular degeneration, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, rubeosis, retinoblastoma, uveitis and Pterygia (abnormal blood vessel growth) of the eye; rheumatoid arthritis; psoriasis; delayed wound healing; endometriosis; vasculogenesis; granulations; hypertrophic scars (keloids); nonunion fractures; scleroderma; trachoma; vascular adhesions; myocardial angiogenesis; coronary collaterals; cerebral collaterals; arteriovenous malformations; ischemic limb angiogenesis; Osler-Webber Syndrome; plaque neovascularization; telangiectasia; hemophilic joints; angiofibroma; fibromuscular dysplasia; wound granulation; Crohn's disease; and atherosclerosis. Moreover, antagonists of polypeptides of the gene product and polynucleotides of the gene may be useful in treating anti-hyperproliferative diseases and/or anti-inflammatory known in the art and/or described herein.

One skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), antibodies, agonists, and/or antagonists and fragments and variants thereof.

Example 56: Cellular Adhesion Molecule (CAM) Expression on Endothelial Cells

The recruitment of lymphocytes to areas of inflammation and angiogenesis involves specific receptor-ligand interactions between cell surface adhesion molecules (CAMs) on lymphocytes and the vascular endothelium. The adhesion process, in both normal and pathological settings, follows a multi-step cascade that involves intercellular adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule-1 (VCAM-1), and endothelial leukocyte adhesion molecule-1 (E-selectin) expression on endothelial cells (EC). The expression of these molecules and others on the vascular endothelium determines the efficiency with which leukocytes may adhere to the local vasculature and extravasate into the local tissue during the development of an inflammatory response. The local concentration of cytokines and growth factor participate in the modulation of the expression of these CAMs.

Briefly, endothelial cells (e.g., Human Umbilical Vein Endothelial cells (HUVECs)) are grown in a standard 96 well plate to confluence, growth medium is removed from the cells and replaced with 100 μ l of 199 Medium (10% fetal bovine serum (FBS)). Samples for testing and positive or negative controls are added to the plate in triplicate (in 10 μ l volumes). Plates are then incubated at 37°C for either 5 h (selectin and integrin expression) or 24 h (integrin expression only). Plates are aspirated to remove medium and 100 μ l of 0.1% paraformaldehyde-PBS(with Ca⁺⁺ and Mg⁺⁺) is added to each well. Plates are held at 4°C for 30 min. Fixative is removed from the wells and wells are washed 1X with PBS(+Ca,Mg) + 0.5% BSA and drained. 10 μ l of diluted primary antibody is added to the test and control wells. Anti-ICAM-1-Biotin, Anti-VCAM-1-Biotin and Anti-E-selectin-Biotin are used at a concentration of 10 μ g/ml (1:10 dilution of 0.1 mg/ml stock antibody). Cells are incubated at 37°C for 30 min. in a humidified environment. Wells are washed three times with PBS(+Ca,Mg) + 0.5% BSA. 20 μ l of diluted ExtrAvidin-Alkaline Phosphatase (1:5,000 dilution, referred to herein as the working dilution) are added to each well and incubated at 37°C for 30 min. Wells are washed three times with PBS(+Ca,Mg)+0.5% BSA. Dissolve 1 tablet of p-Nitrophenol Phosphate pNPP per 5 ml of glycine buffer (pH 10.4). 100 μ l of pNPP substrate in glycine buffer is added to each test well. Standard wells in triplicate are prepared from the working dilution of

the ExtrAvidin-Alkaline Phosphatase in glycine buffer: $1:5,000$ (10^0) $> 10^{-0.5} > 10^{-1} > 10^{-1.5}$. 5 μ l of each dilution is added to triplicate wells and the resulting AP content in each well is 5.50 ng, 1.74 ng, 0.55 ng, 0.18 ng. 100 μ l of pNPN reagent is then added to each of the standard wells. The plate is incubated at 37°C for 4h. A volume of 50 μ l of 3M NaOH is added to all wells. The plate is read on a plate reader at 405 nm using the background subtraction option on blank wells filled with glycine buffer only. Additionally, the template is set up to indicate the concentration of AP-conjugate in each standard well [5.50 ng; 1.74 ng; 0.55 ng; 0.18 ng]. Results are indicated as amount of bound AP-conjugate in each sample.

10

Example 57: Alamar Blue Endothelial Cells Proliferation Assay

This assay may be used to quantitatively determine protein mediated inhibition of bFGF-induced proliferation of Bovine Lymphatic Endothelial Cells (LECs), Bovine Aortic Endothelial Cells (BAECs) or Human Microvascular Uterine Myometrial Cells (UTMECs). This assay incorporates a fluorometric growth indicator based on detection of metabolic activity. A standard Alamar Blue Proliferation Assay is prepared in EGM-2MV with 10 ng /ml of bFGF added as a source of endothelial cell stimulation. This assay may be used with a variety of endothelial cells with slight changes in growth medium and cell concentration. Dilutions of the protein batches to be tested are diluted as appropriate. Serum-free medium (GIBCO SFM) without bFGF is used as a non-stimulated control and Angiostatin or TSP-1 are included as a known inhibitory controls.

Briefly, LEC, BAECs or UTMECs are seeded in growth media at a density of 5000 to 2000 cells/well in a 96 well plate and placed at 37-C overnight. After the overnight incubation of the cells, the growth media is removed and replaced with GIBCO EC-SFM. The cells are treated with the appropriate dilutions of the protein of interest or control protein sample(s) (prepared in SFM) in triplicate wells with additional bFGF to a concentration of 10 ng/ ml. Once the cells have been treated with the samples, the plate(s) is/are placed back in the 37° C incubator for three days. After three days 10 ml of stock alamar blue (Biosource Cat# DAL1100) is added to each well and the plate(s) is/are placed back in the 37°C incubator for four hours. The

plate(s) are then read at 530nm excitation and 590nm emission using the CytoFluor fluorescence reader. Direct output is recorded in relative fluorescence units.

Alamar blue is an oxidation-reduction indicator that both fluoresces and changes color in response to chemical reduction of growth medium resulting from cell growth. As cells grow in culture, innate metabolic activity results in a chemical reduction of the immediate surrounding environment. Reduction related to growth causes the indicator to change from oxidized (non-fluorescent blue) form to reduced (fluorescent red) form. i.e. stimulated proliferation will produce a stronger signal and inhibited proliferation will produce a weaker signal and the total signal is proportional to the total number of cells as well as their metabolic activity. The background level of activity is observed with the starvation medium alone. This is compared to the output observed from the positive control samples (bFGF in growth medium) and protein dilutions.

15 **Example 58: Detection of Inhibition of a Mixed Lymphocyte Reaction**

This assay can be used to detect and evaluate inhibition of a Mixed Lymphocyte Reaction (MLR) by gene products (e.g., isolated polypeptides). Inhibition of a MLR may be due to a direct effect on cell proliferation and viability, modulation of costimulatory molecules on interacting cells, modulation of adhesiveness between lymphocytes and accessory cells, or modulation of cytokine production by accessory cells. Multiple cells may be targeted by these polypeptides since the peripheral blood mononuclear fraction used in this assay includes T, B and natural killer lymphocytes, as well as monocytes and dendritic cells.

Polypeptides of interest found to inhibit the MLR may find application in diseases associated with lymphocyte and monocyte activation or proliferation. These include, but are not limited to, diseases such as asthma, arthritis, diabetes, inflammatory skin conditions, psoriasis, eczema, systemic lupus erythematosus, multiple sclerosis, glomerulonephritis, inflammatory bowel disease, crohn's disease, ulcerative colitis, arteriosclerosis, cirrhosis, graft vs. host disease, host vs. graft disease, hepatitis, leukemia and lymphoma.

Briefly, PBMCs from human donors are purified by density gradient centrifugation using Lymphocyte Separation Medium (LSM[®], density 1.0770 g/ml,

Organon Teknika Corporation, West Chester, PA). PBMCs from two donors are adjusted to 2×10^6 cells/ml in RPMI-1640 (Life Technologies, Grand Island, NY) supplemented with 10% FCS and 2 mM glutamine. PBMCs from a third donor is adjusted to 2×10^5 cells/ml. Fifty microliters of PBMCs from each donor is added to wells of a 96-well round bottom microtiter plate. Dilutions of test materials (50 μ l) is added in triplicate to microtiter wells. Test samples (of the protein of interest) are added for final dilution of 1:4; rhuIL-2 (R&D Systems, Minneapolis, MN, catalog number 202-IL) is added to a final concentration of 1 μ g/ml; anti-CD4 mAb (R&D Systems, clone 34930.11, catalog number MAB379) is added to a final concentration of 10 μ g/ml. Cells are cultured for 7-8 days at 37°C in 5% CO₂, and 1 μ C of [³H] thymidine is added to wells for the last 16 hrs of culture. Cells are harvested and thymidine incorporation determined using a Packard TopCount. Data is expressed as the mean and standard deviation of triplicate determinations.

Samples of the protein of interest are screened in separate experiments and compared to the negative control treatment, anti-CD4 mAb, which inhibits proliferation of lymphocytes and the positive control treatment, IL-2 (either as recombinant material or supernatant), which enhances proliferation of lymphocytes.

One skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), antibodies, agonists, and/or antagonists and fragments and variants thereof.

It will be clear that the invention may be practiced otherwise than as particularly described in the foregoing description and examples. Numerous modifications and variations of the present invention are possible in light of the above teachings and, therefore, are within the scope of the appended claims.

The entire disclosure of each document cited (including patents, patent applications, journal articles, abstracts, laboratory manuals, books, or other disclosures) in the Background of the Invention, Detailed Description, and Examples is hereby incorporated herein by reference. Further, the hard copy of the sequence listing submitted herewith and the corresponding computer readable form are both incorporated herein by reference in their entireties. Additionally, the contents of U.S. Provisional Applications Serial No. 60/164,744 and 60/215,140 are all hereby incorporated by reference in their entirety.

405

| | | | |
|---|------------|-------------------------------|------------|
| Applicant's or agent's file reference number | PS712PCT.. | International application No. | UNASSIGNED |
|---|------------|-------------------------------|------------|

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13 bis)

| | |
|--|-----------------------------|
| A. The indications made below relate to the microorganism referred to in the description on page <u>48</u> , line <u>N/A</u> | |
| B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/> | |
| Name of depositary institution American Type Culture Collection | |
| Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America | |
| Date of deposit 26 October 1999 | Accession Number PTA-868 |
| C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/> | |
| | |
| D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States) | |
| Europe In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC). Continued on the Attached Pages 2 & 3 | |
| E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable) | |
| The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit") | |
| | |

| | |
|---|--|
| For receiving Office use only | For International Bureau use only |
| <input type="checkbox"/> This sheet was received with the international application | <input type="checkbox"/> This sheet was received by the International Bureau on: |
| Authorized officer | Authorized officer |
| | |

ATCC Deposit No. PTA-868**Page No. 2****CANADA**

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

NORWAY

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

AUSTRALIA

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

FINLAND

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

UNITED KINGDOM

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

ATCC Deposit No.: PTA-868**Page No. 3****DENMARK**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

SWEDEN

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

NETHERLANDS

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

What Is Claimed Is:

1. An isolated nucleic acid molecule comprising a polynucleotide having a nucleotide sequence at least 95% identical to a sequence selected from the group consisting of:
- 5
- (a) a polynucleotide fragment of SEQ ID NO:X or a polynucleotide fragment of the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;
 - (b) a polynucleotide encoding a polypeptide fragment of SEQ ID NO:Y or a
10 polypeptide fragment encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;
 - (c) a polynucleotide encoding a polypeptide domain of SEQ ID NO:Y or a polypeptide domain encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;
 - 15 (d) a polynucleotide encoding a polypeptide epitope of SEQ ID NO:Y or a polypeptide epitope encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;
 - (e) a polynucleotide encoding a polypeptide of SEQ ID NO:Y or the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X,
20 having biological activity;
 - (f) a polynucleotide which is a variant of SEQ ID NO:X;
 - (g) a polynucleotide which is an allelic variant of SEQ ID NO:X;
 - (h) a polynucleotide which encodes a species homologue of the SEQ ID NO:Y;
 - 25 (i) a polynucleotide capable of hybridizing under stringent conditions to any one of the polynucleotides specified in (a)-(h), wherein said polynucleotide does not hybridize under stringent conditions to a nucleic acid molecule having a nucleotide sequence of only A residues or of only T residues.

2. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises a nucleotide sequence encoding a secreted protein.
- 5 3. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises a nucleotide sequence encoding the sequence identified as SEQ ID NO:Y or the polypeptide encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X.
- 10 4. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises the entire nucleotide sequence of SEQ ID NO:X or the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X.
- 15 5. The isolated nucleic acid molecule of claim 2, wherein the nucleotide sequence comprises sequential nucleotide deletions from either the C-terminus or the N-terminus.
- 20 6. The isolated nucleic acid molecule of claim 3, wherein the nucleotide sequence comprises sequential nucleotide deletions from either the C-terminus or the N-terminus.
- 25 7. A recombinant vector comprising the isolated nucleic acid molecule of claim 1.
8. A method of making a recombinant host cell comprising the isolated nucleic acid molecule of claim 1.
9. A recombinant host cell produced by the method of claim 8.
- 30 10. The recombinant host cell of claim 9 comprising vector sequences.

11. An isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence selected from the group consisting of:

(a) a polypeptide fragment of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;

5 (b) a polypeptide fragment of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z, having biological activity;

(c) a polypeptide domain of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;

10 (d) a polypeptide epitope of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;

(e) a secreted form of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;

(f) a full length protein of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;

15 (g) a variant of SEQ ID NO:Y;

(h) an allelic variant of SEQ ID NO:Y; or

(i) a species homologue of the SEQ ID NO:Y.

20 12. The isolated polypeptide of claim 11, wherein the secreted form or the full length protein comprises sequential amino acid deletions from either the C-terminus or the N-terminus.

13. An isolated antibody that binds specifically to the isolated polypeptide of claim 11.

25 14. A recombinant host cell that expresses the isolated polypeptide of claim 11.

15. A method of making an isolated polypeptide comprising:

30 (a) culturing the recombinant host cell of claim 14 under conditions such that said polypeptide is expressed; and

(b) recovering said polypeptide.

16. The polypeptide produced by claim 15.

17. A method for preventing, treating, or ameliorating a medical condition, comprising administering to a mammalian subject a therapeutically effective amount
5 of the polypeptide of claim 11 or the polynucleotide of claim 1.

18. A method of diagnosing a pathological condition or a susceptibility to a pathological condition in a subject comprising:

(a) determining the presence or absence of a mutation in the polynucleotide of
10 claim 1; and

(b) diagnosing a pathological condition or a susceptibility to a pathological condition based on the presence or absence of said mutation.

19. A method of diagnosing a pathological condition or a susceptibility to
15 a pathological condition in a subject comprising:

(a) determining the presence or amount of expression of the polypeptide of claim 11 in a biological sample; and

(b) diagnosing a pathological condition or a susceptibility to a pathological condition based on the presence or amount of expression of the polypeptide.
20

20. A method for identifying a binding partner to the polypeptide of claim 11 comprising:

(a) contacting the polypeptide of claim 11 with a binding partner; and

(b) determining whether the binding partner effects an activity of the
25 polypeptide.

21. The gene corresponding to the cDNA sequence of SEQ ID NO:Y.

22. A method of identifying an activity in a biological assay, wherein the
30 method comprises:

(a) expressing SEQ ID NO:X in a cell;

(b) isolating the supernatant;

- (c) detecting an activity in a biological assay; and
- (d) identifying the protein in the supernatant having the activity.

23. The product produced by the method of claim 20.

<110> Human Genome Sciences, Inc.

<120> 28 Human Secreted Proteins

<130> PS712PCT

<140> Unassigned

<141> 2000-11-08

<150> 60/164,744

<151> 1999-11-12

<150> 60/215,140

<151> 2000-06-30

<160> 190

<170> PatentIn Ver. 2.0

<210> 1

<211> 733

<212> DNA

<213> Homo sapiens

<400> 1

| | | | | | | |
|------------|-------------|------------|-------------|------------|------------|-----|
| gggatccgga | gcccaaactct | tctgacaaaa | ctcacacatg | cccaccgtgc | ccagcacctg | 60 |
| aattcgaggg | tgacaccgtca | gtcttcctct | tcccccaaaa | acccaaggac | accctcatga | 120 |
| tctcccggac | tcctgaggtc | acatgcgtgg | tggtggacgt | aagccacgaa | gaccctgagg | 180 |
| tcaagttcaa | ctggtacgtg | gacggcgtgg | aggtgcataa | tgccaagaca | aagccgcggg | 240 |
| aggagcagta | caacagcacg | taccgtgtgg | tcagcgtcct | caccgtcctg | caccaggact | 300 |
| ggctgaatgg | caaggagtac | aagtgcgaag | tctccaacaa | agccctccca | acccccatcg | 360 |
| agaaaaccat | ctccaaagcc | aaagggcagc | cccagagaacc | acaggtgtac | accctgcccc | 420 |
| catcccggga | tgagctgacc | aagaaccagg | tcagcctgac | ctgcctggtc | aaaggcttct | 480 |
| atccaagcga | catcgccgtg | gagtgggaga | gcaatgggca | gccggagaac | aactacaaga | 540 |
| ccacgcctcc | cgtgctggac | tccgacggct | ccttcttctc | ctacagcaag | ctcaccgtgg | 600 |
| acaagagcag | gtggcagcag | gggaacgtct | tctcatgctc | cgtgatgcat | gaggctctgc | 660 |
| acaaccacta | cacgcagaag | agcctctccc | tgtctccggg | taaatgagtg | cgacggccgc | 720 |
| gactctagag | gat | | | | | 733 |

<210> 2

<211> 5

<212> PRT

<213> Homo sapiens

<220>

<221> Site

<222> (3)

<223> Xaa equals any of the twenty naturally occurring L-amino acids

<400> 2

Trp Ser Xaa Trp Ser

1

5

<210> 3

<211> 86

<212> DNA

<213> Homo sapiens

<400> 3

gcgcctcgag atttccccga aatctagatt tccccgaaat gatttccccg aaatgatttc 60
cccgaaatat ctgccatctc aattag 86

<210> 4

<211> 27

<212> DNA

<213> Homo sapiens

<400> 4

gcggcaagct ttttgcaaag cctagggc 27

<210> 5

<211> 271

<212> DNA

<213> Homo sapiens

<400> 5

ctcgagattt ccccgaaatc tagatttccc cgaaatgatt tccccgaaat gatttccccg 60
aaatatctgc catctcaatt agtcagcaac catagtcccg cccctaactc cgcccatccc 120
gccctaact ccgcccagtt ccgcccattc tccgcccacat ggctgactaa ttttttttat 180
ttatgcagag gccgaggccg cctcggcctc tgagctattc cagaagtagt gaggaggctt 240
ttttggaggc ctaggctttt gcaaaaagct t 271

<210> 6

<211> 32

<212> DNA

<213> Homo sapiens

<400> 6

gcgctcgagg gatgacagcg atagaacccc gg 32

<210> 7

<211> 31

<212> DNA

<213> Homo sapiens

<400> 7

gcgaagcttc gcgactcccc ggatccgcct c 31

<210> 8

<211> 12

<212> DNA

<213> Homo sapiens

<400> 8

ggggactttc cc 12

<210> 9

<211> 73
 <212> DNA
 <213> Homo sapiens

<400> 9
 gcggcctcga ggggactttc ccgggggactt tccggggact ttccgggact ttccatcctg 60
 ccatctcaat tag 73

<210> 10
 <211> 256
 <212> DNA
 <213> Homo sapiens

<400> 10
 ctcgagggga ctttcccggg gactttccgg ggactttccg ggactttcca tctgccatct 60
 caattagtcg gcaaccatag tcccggccct aactccgccc atcccggccc taactccgcc 120
 cagtcccgcc cattctccgc cccatggctg actaatTTTT tttatttatg cagagggcga 180
 ggccgcctcg gcctctgagc tattccagaa gtagttagga ggcttttttg gaggcctagg 240
 cttttgcaaa aagctt 256

<210> 11
 <211> 2670
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (2664)
 <223> n equals a,t,g, or c

<400> 11
 aattccccggg tgcacccacg cgtccggcag catgctgtgt gccctgctcc tcctgcccag 60
 cctcctgggg gccaccaggg ccagcccccac ctccagggccc caggagtgtg caaagggctc 120
 caccggtgtg tgtcaggatc tgcagacagc tgccaggctg ggggctgtgg ggtactgcca 180
 aggggcccgtg tggaaacaaac ccaccgcgaa gtctctgccc tgcgacgtat gccaggacat 240
 agcagccgcc gctggcaatg ggctgaaccc tgacgccacg gagtctgaca tcctggcttt 300
 ggtgatgaag acctgtgagt ggctccccag ccaggagtct tcagccggat gcaagtggat 360
 ggtggatgcc cacagtccgg ccacccctgag catgctccgt ggggccccgg acagtgcccc 420
 ggcacagggt tgcacagcgc tcagcctctg tgagccgctg cagaggcacc tggccaccct 480
 gaggccactc tccaaagagg acacctttga ggctgtggct ccgttcattg ccaatgggcc 540
 ccttaccttc cacccccggc aggcgcctga aggagctctg tgccaagact gtgtacggca 600
 ggtctcccga ctccaggagg ctgtccgggtc caacttgacc ttggccgact tgaacatcca 660
 ggagcagtggt gagtccctgg ggctggcctt ggcgctctc tgcaagaact acctcttcca 720
 gttttttgtc cctgctgacc aagcactgag gcttctcccc ccgcaggagc tctgcaggaa 780
 ggggggattc tgtgaggagc taggggcacc tgcccgtttg actcaagtag tggccatgga 840
 cgggggtcccc tccctggagc tgggggtgcc aaggaaacag agcgagatgc agatgaaggc 900
 cgggtgtgacc tgtgagggtg gcatgaacgt ggtgcagaag ctggaccact ggctcatgtc 960
 caacagctct gagctcatga tcacccatgc cctggagcgc gtgtgctcgg taatgcctgc 1020
 ctctatcacg aaggagtgca tcactctggt ggacacctac agccccctct tgggtgcagct 1080
 tgtggccaaa atcaccaccg agaaggtgtg caagttcatc cgtctgtgtg gcaaccggag 1140
 gcggggccgg gcagtccatg atgcctatgc catcgtgccc tcccagagt gggatgcgga 1200
 gaaccagggc agcttctgca atgggtgcaa gaggctgctc acgggtgcct cccacaacct 1260
 ggagagcaag agcaccaagc gagacatcct ggtggccttc aagggtggct gcagcatcct 1320
 gccgctgccc tatatgatcc agtgcaagca cttcgtcacc cagtacgagc ccgtgctcat 1380
 tgagagtctc aaggacatga tggaccccggt ggctgtgtgc aagaagggtg gggcctgcca 1440
 cggcccccagg accccactgc tgggcaccga ccagtgtgcc ctgggcccga gcttctggtg 1500

| | | | | | | |
|-------------|------------|-------------|-------------|-------------|------------|------|
| caggagccag | gaggccgcaa | gctgtgcaac | gctgtgcaac | actgccagaa | gcatgtatgg | 1560 |
| aaagagatgc | acctccacgc | tggggaacac | gcgtgaccgt | ggctgccaga | gacccagagc | 1620 |
| ctgttagcga | ggcccatgag | gtgggtgctt | tccccatccc | catttcacaa | atgaaaaact | 1680 |
| gaagctctga | ggagggaggg | tgggaaggag | cagagctgaa | gttcaaaacc | aagtattcct | 1740 |
| gatcccga | gcctctctct | taacaacggt | gccgcacagc | tttgcccttg | aaagcatctc | 1800 |
| tactggaccg | gaacacactc | atgtgccccg | ctccctgacc | cagccaaggc | tgccctttca | 1860 |
| tctccaaggc | tgagatgttg | ccgggtgttc | catgagagcc | tgcccatggg | ctcaggtgcc | 1920 |
| cctttacctt | ctgctggatg | gacatctggc | tgtgagccag | gctgggggtca | tggccggggg | 1980 |
| gagcggaggg | aggggtggac | ggaggcttcg | agggcccatc | actagtaggg | tcattacctc | 2040 |
| ttgccaacag | ccgggggtgg | gagtcgtggg | ctcgctcagg | ccagagcttc | tcaacctgga | 2100 |
| gtccctgggg | gtggctgcca | aagggtgtgt | tgacaagcac | gtatccctgg | acatttcagg | 2160 |
| ggagaggtct | ggggcttttg | tcacattctc | caagggtctc | tgggtctcgg | agcagtcctc | 2220 |
| ccccatgtct | cagccactac | agggctccctc | tctctccttg | caccccagac | cctccgctgc | 2280 |
| cctggtaatg | agcagaagga | aagtcttggtg | gtgtgctcaa | agtcaggaga | gcaaaatatg | 2340 |
| ccaggcaaaa | gctcccggga | aaagcctgag | gagtcctggg | tggccaccgg | gatgtggagc | 2400 |
| agcaggggga | aagacggtga | acacagccct | ccagctgtct | gagcctcagt | tttctaattc | 2460 |
| gtagaatggg | gatgatcata | cctgcctcac | aagaatgttg | agacaattca | cagagacgtt | 2520 |
| ctggagcccc | tttccccgga | gacgggcatt | catgagctcg | ctgggaccag | aaaaccctac | 2580 |
| tcagggggccc | agcggggcac | ccaggagagt | ctggcgggtgc | aagcgctgta | taaaccacaa | 2640 |
| gcgttctcca | aaaaaaaaaa | aaanaaaaaa | | | | 2670 |

<210> 12

<211> 2187

<212> DNA

<213> Homo sapiens

<400> 12

| | | | | | | |
|-------------|------------|-------------|------------|-------------|-------------|------|
| gcgtccgaca | acatggctga | cgtgtcagcc | aaggacagca | gccaggagac | gctgggtgaac | 60 |
| ctggcggggc | tcttggtcag | cctcctgatg | ctccctctgg | tgctagggtg | ccctggcttc | 120 |
| agccttggat | gtttcttctt | cctcactgcc | ctccacatct | acgccaaacta | ccgcgcgggc | 180 |
| cgagccctgg | tcatggagac | cttgaacgaa | ggccggctcc | ggctggctct | gaagcactac | 240 |
| cttcagaggg | gagaggtact | cgacccaact | gcagccaatc | gcatggagcc | gctgtggaca | 300 |
| ggtttctggc | cagctccgtc | tctatccctg | ggggctccct | tacaccgctt | ggctctccagt | 360 |
| gtctttgagc | tgagcagct | ggttgagggg | caccaagaat | cctacctcct | ctgctgggac | 420 |
| cagtcacaaa | accaggtaca | ggtagttctg | aaccagaagg | caggccccaa | gaccatccta | 480 |
| agggccgcca | cacatgggct | gatgcttggg | gccctgcagg | gagatggacc | ccttcagca | 540 |
| gagctggagg | agctgaggaa | ccgggtgcgg | gcaggctcta | gaaagagagc | tgggtcgtcg | 600 |
| tcaaggagac | acacgaagtg | ttggacatgc | tgttcccaaa | gttcttgaaa | ggactgcagg | 660 |
| atgccggctg | gaagaccgag | aagcaccagc | tagaggtgga | tgagtggagg | gccacatggc | 720 |
| ttctgtctcc | cgaaaagaag | gtcttgtgag | cagcccagac | ggaggcccaa | gcccagggca | 780 |
| ggaacctgga | gcaaggacac | tttggccaca | gcaggacagg | ggaaaggcag | ctttattttt | 840 |
| ccttagggca | actgcagcgg | gtggggcagg | ccctcatggg | aagtgactgc | caatcagatg | 900 |
| cagtggggccc | caggcagagg | aaggccggga | gaaggggagc | caggaccttc | tcacccact | 960 |
| gccccttccc | cttttctggg | gagcaccgca | ggctcctcac | ccccacttcc | tgtgaggctg | 1020 |
| tggcttatgg | tgtccaacgc | agttgggtctt | aggcatagaa | gccccagagg | aacacggcca | 1080 |
| ctgccatcat | gagcagggca | ttgaggttga | ccacacgggc | ccagctcggg | tcctcgctga | 1140 |
| tgtcctccag | ccgcctggct | gctgccgctg | cctcctcctg | ggtaaggggc | ggaggactgc | 1200 |
| ccacccccacc | tctgctcatt | ccacaaaacc | agagcaggca | ctggcggaag | aggcttggtg | 1260 |
| ccggggcctg | gggctctgag | ggaaattgag | gccctgcagt | tagtttgctg | gaactcagct | 1320 |
| cctccagccc | cacctccag | catggtgccc | taccattcat | ctccatggca | ctctctgggc | 1380 |
| acccattctg | tacagggagt | gaggagcctt | gctgctcatc | agcatccagg | tcctcccgtt | 1440 |
| cctccttgct | atgccggaga | ctgaagacca | ggcgggtggg | ctggggaggg | tgggagcacg | 1500 |
| aacgaggtgg | gagttctgtc | cccccatgcc | tggccctaaa | gtctcttgca | caccagctcg | 1560 |
| tactgcctg | ccctaccac | ctctgtccag | tctacacacc | cagcccaggc | ttaactcatg | 1620 |
| ccaactccac | cctacatggc | tgccctgtgc | cctcgggata | aaccccaagc | ccctgagctt | 1680 |
| gtgtttaaag | ccgttgccct | tgtcccccca | gctttgtcag | ctcaggtctg | tctacacca | 1740 |
| gatggtagcg | cttgtgacac | tggcctggca | gtcctgctca | cagtgttctg | tgctgtgtg | 1800 |

| | | | | | | |
|------------|------------|-------------|------------|------------|------------|------|
| ctccccacct | ttcctcctgc | tgctgcagaa | acccggccat | ccttccacac | ccagatctct | 1860 |
| tgtcctgtcc | tcacccacc | ctgccaccat | cagccctgcc | tggagccacc | tgccctttgg | 1920 |
| caacaaaacc | aaaccttttt | gtgggcgttc | aagatgggat | tgtgcccacc | agtcagatcc | 1980 |
| tgtgttttga | gtcccaaagg | ccatgccaaag | gattggcctt | tgggaggcct | taatcaccaa | 2040 |
| cccatcaaca | tcaagcctcc | cccaggccgg | ttcaaataaa | tgtattttaa | taaaaaaaaa | 2100 |
| aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | 2160 |
| aaaaaaaaaa | aaaaaaaaaa | aaaaaaa | | | | 2187 |

<210> 13

<211> 2664

<212> DNA

<213> Homo sapiens

<400> 13

| | | | | | | |
|-------------|-------------|-------------|-------------|-------------|-------------|------|
| ccacgcgtcc | gctctgtcac | tggtgccttc | tggaagatgg | cttccccctg | gggagcattt | 60 |
| cctgcaaaga | ctcacaagtt | taaaataatt | taggatattt | ctctgtactt | gagagccagg | 120 |
| ataggatgag | agttggctct | gccctgatgc | tgtaaggcca | ctcctgtctc | agaccctgag | 180 |
| gtcctcagga | gtgggttccc | aagtactgtt | gtagttgtat | ggcaagatca | ctcttgatta | 240 |
| ttttaggagc | tgattttact | tttctacct | cttttaactg | cttccaaaag | atgaatctgg | 300 |
| ccaagaagag | caggggctca | ttcactcatt | tactcactca | ttcctgggtgc | ttgagtttat | 360 |
| tcctgaaggga | agctgaccaa | gggttgagag | aaaacaactt | tgatttttct | catgtctgcc | 420 |
| ccagcaaac | tcctttgtgg | actgattctc | cttctgtgcc | tggaagaaac | tgggataatc | 480 |
| ccaggacttt | tctgggtccct | tctaggtaat | tcagtgtctt | ccactaatgt | ggccccacac | 540 |
| aggctgggct | tgttgattag | tatagggtcca | gcaaaacagt | taaaactcta | catgtcttcc | 600 |
| gcatgtgaac | ctcttcccaa | cctcacctgg | tgctgtggca | gggtgggaggc | tgggtgtcag | 660 |
| gctggctgct | gccccctggg | ttctgcccag | ctggtagcta | tggcttccaa | ccctccagtg | 720 |
| ggaggaacaa | gagggataag | aggtgaaggac | aggagagggt | ctgtatggct | ggcatggcag | 780 |
| agagctggca | catgaggctg | gcaggaaatc | cagagggctc | tggggggcatt | ctcatggctc | 840 |
| cctaagtgc | cacactaccc | tgggggtgtc | ctgccacaac | tggccaactc | aggaggtgta | 900 |
| tcctctccag | ctgggtccctt | acactcactt | ccttagccct | agggtggccgt | ggtacctcta | 960 |
| gcctcttgcc | aagggtctgcc | ctcacatctc | caggaaacct | tctgggcagg | atttaagaca | 1020 |
| tcccaggcca | gtgttcaggc | actatgacct | cacctggctc | acaggaaatg | tgcacctttg | 1080 |
| atccactggg | ttaggggttat | ttccacatgc | agcctagtgt | cctactggga | gctccaggag | 1140 |
| tcaacttttg | cccttttagat | ttcttgagt | taagagggat | accaggctct | tgccaccac | 1200 |
| cagcctcaga | aggcacacag | taggcacttg | caggcctcgt | gaaacccctc | ttggaatgaa | 1260 |
| gtaaggagaa | agctgccccat | ccctctcatg | gtacagggtt | gggtggggaca | gactgagagc | 1320 |
| actgcccccc | tctcccaagg | aagccctcct | gtcaaaactg | agcttctctc | atcttcaaca | 1380 |
| cctcctctac | ctagatcacc | cagaatttca | catgggaaca | attggcactt | cttgtcttgg | 1440 |
| ggcctcagct | aaaattcaga | ctggaaaagc | tccatattaa | catcctgttc | taatgcaagt | 1500 |
| aaaaaaagag | ctgctatggg | ctttgcttct | tggtatttag | agtctagtgg | gagaatattg | 1560 |
| gctacttaat | gcagtctctt | ccattttcaa | tatgtaactg | aggcatcctt | ctactgataa | 1620 |
| cagccaaaac | ttcaagaaaa | aatattttta | gccttttaaa | atacatttgt | gggctggtag | 1680 |
| gaaattaagg | aataatcaga | gacccaaaac | taagtgaagc | tgagaaccca | gagataactt | 1740 |
| tattaaagga | gaatattaaa | catatacttc | acgcagaagg | aagagaattc | cagggtggaag | 1800 |
| atttgagatg | caaggggggaa | tgaaaaacaa | gaagggtgtg | actatatggg | taaatcaaata | 1860 |
| caaatcaaac | acctttgtgt | aaataataat | gggtgatata | gaatttttaa | aagagaaata | 1920 |
| agtatatagt | cataatatga | tacaaatttg | aagagggtga | tgagttaaaa | agttctaaga | 1980 |
| ttcttgtgtt | aactttggag | gatggttatc | taatttgtaa | taaaattttc | aggataatca | 2040 |
| tcaaccgtag | aaatgggggt | gaaatttcaa | actaatggaa | agaaaaaaga | atgagaaaaat | 2100 |
| ataatcaatc | aaaaggaaaa | gactagaaag | ggaaaaggaaa | aaagaaacac | aaaagagggtg | 2160 |
| gaacaaaagt | ccaaatataat | cagtaactac | aataaacata | aatggaataa | acataacaac | 2220 |
| taaaagataa | agattattaa | agtggattac | tatatgctgt | ttacagaaaa | cacatctaca | 2280 |
| ctctaagggt | acagaaaagg | tgaaaagtaa | aggatgggaa | aggggtatact | ggggaaatat | 2340 |
| aatgaaggga | aatatgatga | atctatatta | atgtaaaaaa | attgattttg | gggtaaaaag | 2400 |
| catcactgga | gatagagaga | atcactatgt | aatgattaaa | ctattcattt | catcaagcag | 2460 |
| ataatataat | tttgaatttt | attctttcca | taacataatt | tcaaaatctg | taaagcaaat | 2520 |
| attgacagaa | ctataaggaa | gaatatataa | attcatcatc | agagagaaat | atttttaata | 2580 |

ctgtaactca gtatTTTTata gaagaagtaa ataaaaagcca gtaaaggTTa tagaacattt 2640
 gaacaatgca aaaaaaaaaa aaaa 2664

<210> 14
 <211> 3393
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (3348)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (3371)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (3386)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (3390)
 <223> n equals a,t,g, or c

<400> 14
 aatcccttcc tatatttggT acgcctgccg gtaccggTcc ggaattcccg ggtcgaccca 60
 cgcgTccggg gaaaagggca TggtTgggag gttgagcagT aagagaccat taatactaag 120
 tatTTTgcct gaaatgcTtc tttataatta tttcagggaa ggaataagat aatctcagac 180
 tTactggggT ggggctTggg gaatgTttgg atgTtgctct tttcttctt cttctttttt 240
 tttttttttt tttttTgTt ggtgTtGttt ggcatTTttt tcttttcctt tttaaagaaa 300
 atgTttTcag gaaacatgaa caaacatact gccaatTata gtggggctgg gaaggcacag 360
 gagctggcgT cctctcagct gcatagTtgg gacggcaaac cctgctgtga gcttctcaga 420
 cttttcacct acttcaccta ttaagaagcc atgaggagTt gtaaactctt gttcagggaa 480
 gaaaagaggg caggaggaak taaccaatg cctttaaata caaaacagat gaaaaacaaa 540
 atactcattt tTccttttct attgtgggTt Tggggagcca aaccaagtgt gactgtgagc 600
 acgtTcactg agaggcatga caacattatt gaagaggtct tTccattatg tctggtTctc 660
 tTatttttctt aagtcaagcc ttaactatgw acatacttTa agatgatacg ggtctgtcac 720
 cataaacctg taaaagggca Tcgacagctt ttcagataat ccaggaatgt tgagacatgg 780
 ctaagtTtct agctgatact tagtcattcc ctcttgatat TgtTgggatg ggtagccaaa 840
 aggcaggctt gtgggagTta agagccaatg atacctggaa acatgggagg tagctggggg 900
 tacaagtggT Tgttgagact gaaggaagag gtggaaagaa gacacagaag gcaccagat 960
 gccaaaaggc ctggcatgaa TgagagTtgg catctgtcct gttgtggTgt tcccattctt 1020
 gccataccca tttgcccaga ttgttctgat tctcatgttc aattgactat cccacatgag 1080
 actacagTtt ataggcagTg aggaattgtt tcttctTggg aaatgtcctt ccagcaactt 1140
 cagaggaatg ggaagtgtt aatcttacgt gaagcaagg agcctggact aggggcagag 1200
 agaaaagaagc agcagcccac cttctttgtg tttgcacatg Tgcctatgtc gtgtacattg 1260
 gtacctgact gtggctcaga tctgcgtcgc agcagcgaga gaagaaatca ctccatatcc 1320
 gatgagagga agggTggcac araratggTg tctacaatta gagacatttc Tgactycacc 1380
 ttagcctaag caaactttat atactgagTa acattTgaag gttgtctttt aatggTgggg 1440
 ggtgtttttt cttttttaaa ctacagTgct Tgcacaagag agggaggggac tywkaawwkk 1500
 ttagggcagg Tgaggggagac agkrgatggs ctgrgatgac ttgagtccat catactattg 1560
 cttggcaggT gTcctcccc atgTttgatt caaattccat gagtgcacta cttttcccca 1620
 ggaatgggac TgagagggtT gtctccagca actcagTctg cacagggctc cccgttcagg 1680

| | | | | | | |
|------------|------------|-------------|------------|-------------|-------------|------|
| ctgccttttg | tggttgtgct | tttgtaagtt | tctttctctg | cacttcgact | tacctctgaa | 1740 |
| tcagaaagca | agcccagcag | gtgaatgagg | gatgtctgct | tggcattgcc | caatctaacc | 1800 |
| agggaggctg | gctggccacc | cactgtccgc | tagaggggag | agctagcagg | tggttggtatg | 1860 |
| aactcaggaa | tagaaacacg | aggccttttt | aaatacgagg | gagaagaatc | catgatgcat | 1920 |
| acctgtaacc | ccctagaacc | caagtgccag | aattcctaga | tgctgcttct | gtttgaacaa | 1980 |
| aatgtcactg | cttttacact | tgaaaaaac | acactcgaaa | aatgttcaac | tccatgaaaa | 2040 |
| atattttttg | gctttaagaa | attgtttggg | gtttaactgt | ttcctttgat | tgccattcca | 2100 |
| ccagtaaat | gttggttgat | ttgcactgca | ctctgggggt | gggggtggga | ggggagggtc | 2160 |
| cttatacaga | gccgaacctg | gggttgctca | ggaagtgggc | caggggaatgt | ggaagtcgtt | 2220 |
| gacattgcct | gggccaaaag | agtgggagat | agtttttctc | ccctcagccc | actcctggta | 2280 |
| gcactgtcg | ccagcctggg | acaaagccag | gcctttttct | tctgtgagca | tctcatcact | 2340 |
| gtccagcagc | agggtgaaaa | agggggacaa | caaccagacc | tattttttct | cccccathtt | 2400 |
| ttccaaat | tgctgtgcca | aatgtttaaa | attttataaa | tatgaatcta | ttgaaatttc | 2460 |
| cttaatcaag | agcttcttcg | tgtaaagttt | gcttttttag | ctatagaaaa | agaaaacagt | 2520 |
| aaatatctct | taatggcatc | cagccttgct | gagctcacct | tttttcctga | agaatgggta | 2580 |
| ggagtgaat | atttaatgta | aacatttcac | caagtccctt | taccctaatt | ttgaagctgc | 2640 |
| attaaacca | actcactaac | acaggggaatg | attgcaccct | agtcctctgt | gggccaagaa | 2700 |
| actttcagaa | gcattaaaaa | aatagttgaa | gtatatcact | tctcaccaag | tggttagagt | 2760 |
| cagttggctg | tttgtccctt | gttttttatt | tattccataa | ttatgtttgt | gctttttgtt | 2820 |
| ttgtaaacag | taatggaacg | tacattttta | ttttgtttag | aagacaactt | tgattcaatc | 2880 |
| tttcaagaac | tgttccattc | ttggtttctt | cttaggggga | taaaaagtta | ccagttaatt | 2940 |
| tgttttgaga | tatttaagca | ttctttgagt | tatagaattg | tgatgcaggg | atttgtgaat | 3000 |
| gagacattta | catgtgaaag | gtgacttcac | tagttacctg | cttgagcaga | gtaaagtgtg | 3060 |
| tatatgtaca | taaaatgtaa | gtaatcttaa | ctccattgtg | cagtgccttt | tagatgttcc | 3120 |
| gctttctata | aagtcttcaa | atttttgcac | atatattata | tatatatgat | gtaatgttat | 3180 |
| agaaatatat | gtataatata | catatttttt | ccaggggtat | ctgatatgctc | tgtattttgt | 3240 |
| tatggaagtt | gaaagaaaaa | agtattttac | ctcagaaatt | aaataaaaaa | atacttttaa | 3300 |
| gtaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | 3360 |
| aaaaaaaaaa | naaaaaaaaa | aaaaanaaan | caa | | | 3393 |

<210> 15

<211> 1604

<212> DNA

<213> Homo sapiens

<400> 15

| | | | | | | |
|-------------|-------------|-------------|------------|-------------|-------------|------|
| ccacgcgtcc | ggcagacaca | ggcacttatt | cattcatctc | attgaaaagc | tacgagttgg | 60 |
| ttccttattg | cctctccata | atagaaaaac | tctttaatga | gctctctttt | tgtttttcaa | 120 |
| atcagatatg | caaagaagct | cataacaatt | ttttttaaaa | atgcaaaaca | agaatctcca | 180 |
| attatgggag | caaaatcttc | agcttctggg | ttcctgtctc | actgaggaaa | tggtattgaa | 240 |
| atggcaagga | ggaaatgagg | aggcaaaactt | tcatgtctat | tttagttttc | caatgcagtc | 300 |
| ctatttccct | tggaactttgt | ataaacaagg | aaaggacagt | tgtagtttca | gttattacag | 360 |
| ataacctgtg | tctttaaagt | aatgtatct | taaataagta | ggactcccat | aaatgactac | 420 |
| actttttcaa | aatatgactc | cccagcttat | aacaagaata | atagcaaaaca | tcacttttatt | 480 |
| aagcaattac | tatgtaaaaag | acacttagtg | cttagcacac | actggaaaata | ttgttgactg | 540 |
| gctatatattt | ccccagaaaat | cccatttctg | aaagcctatt | acaaagaaat | aaaatcatca | 600 |
| gtataacaaa | ggagtgtgtg | tgtgtgtggg | tgtgagtgtg | tgtgggtgtg | agtgtgtggg | 660 |
| gggtgtgagt | gtgtgtgggt | gtgagagtgt | gtgtgtgagt | gtgagtgtgt | gtgtgtgtaa | 720 |
| gtgcacacac | atgttgtagt | attgttcata | gtggcaaaaa | ctagaaacaa | agtgaatatt | 780 |
| gataacatgg | gcacagatga | caaattatat | ctccaaatta | tgaaacagaa | tccagccatt | 840 |
| aaaatcagag | ctttgccacg | tgactaggat | gaagttacaa | aaagtattgt | tgagtgaagaa | 900 |
| aagcaggata | cataggatac | atggaattga | gtataatatg | atttcttttt | tttttttttt | 960 |
| gagatggagt | ctcgtctctg | cactcaggct | ggagtgcagt | ggcacaaatc | cagctcactg | 1020 |
| caacctccgc | ctcccggtat | caagcaattc | tcctccctca | gcctcctgag | tagctggggac | 1080 |
| taccgtcacc | tgccaccacg | ccccagctaa | tttttgtatt | tttagtagag | atgggctttc | 1140 |
| accatattgg | tcacgtgat | ctcaaaatcc | tgacctcagg | tgatccacct | gcctcagcct | 1200 |
| cccaaagtgc | tgggattaca | gtcgtgagcc | actgcacctg | gccgatttct | tttttaaaat | 1260 |

| | | | | | | |
|------------|-------------|------------|------------|------------|------------|------|
| gatcaaaaaa | ccatttatat | gtgggaatat | agctatatac | ttttattatt | gaattaccat | 1320 |
| ggaaaaaac | atggaagagg | gaggccaagg | caggagaatc | acttgaggcc | caggagtttg | 1380 |
| agaccagcct | gggcaacaaa | gcgagaccct | catctctact | aaaaatacaa | aaattacctg | 1440 |
| ggcctggtga | cacatgcctg | taatcccagc | tactcagaag | actaaggcaa | gtgaatcgct | 1500 |
| tgaacccgag | acgtggaggt | tgcagtgaag | tgagccaaga | tcgcgccgtt | gcactccagc | 1560 |
| ctggtgacag | agtgaagagtc | tttctcaaaa | aaaaaaaaaa | aaaa | | 1604 |

<210> 16

<211> 3541

<212> DNA

<213> Homo sapiens

<400> 16

| | | | | | | |
|-------------|-------------|-------------|-------------|-------------|-------------|------|
| gcgtccgcta | acctgtagat | aactatagat | gaaactcaca | ttttccttcc | cctgggttcac | 60 |
| cttgacagca | ttgcagctct | ggtcagccac | agaatgccag | gcagtgggtg | acaccatgat | 120 |
| agccgtgtgg | tcagaaggga | agggcacggg | agtgagctgg | gaaccttggc | tgttggggaa | 180 |
| gttgcaaagc | tccagtttcc | tctaagtgtg | cctttatgat | ccgcactggc | gacagcaggc | 240 |
| tttgtctttc | cttagaaaaga | ctttgtgcct | gttctcactt | gaccagatat | cgtgtacgga | 300 |
| gcacttgctg | gcagtaggcc | aagctctgca | cattcatgaa | ttgctttgtc | tcacaacacc | 360 |
| ctgagccgta | ggtatcttta | ttgaccatat | tttccagatg | aagagacgaa | agcccaagag | 420 |
| aaccctgtta | atgctcggcc | agatcagagg | agacaaaaca | agcacccttg | ccttcatggg | 480 |
| ttgtgtggat | gaatgagggt | cggcctaggg | ctgtcccttc | gcggtctgtc | cggctctgtcg | 540 |
| gtacaggcag | cagatgggcc | acttgttcct | ggaattgcag | aagcagataa | ccaagtgtag | 600 |
| aaaaatgtatg | tatccctctt | ctaccccaac | agaagcacccg | tctttttgct | ttttcttttt | 660 |
| tttttttagtt | acagatatga | taggtaggtt | tttttctatg | gtgtaagcat | ttctgctggt | 720 |
| ggattttgtt | tgctgtttga | agacctgcct | aactcgtctc | agtaggctgt | ataaatcggt | 780 |
| taatgaagat | tggccctttt | ctgaagttag | aaaaaggatg | gttgctgaag | aacctatctc | 840 |
| gagaatacac | tattgacaga | ctcacggtgt | ctgtgtgtct | cacatcctgg | aattacttgc | 900 |
| acaccattgt | ggctatgtgt | tatatcccag | aatatggaat | tgattttgat | gtttagaagg | 960 |
| ccaaagctta | cttagtaaat | gccacactga | ttttagctta | ccattttgta | gtttttttct | 1020 |
| gtgggaaata | acaaaaaaat | tattgattga | ggatctatat | tgattttaca | gagctatttt | 1080 |
| agaggaagta | tttctcattt | aatgaacact | ttacattgga | ctttccatct | taacaagaca | 1140 |
| ctgccccagg | tggaaactctg | tcttcacaac | acctgagctg | cccagtgggtg | tcagattgac | 1200 |
| tgggtgtcagt | ttaaccccg | taagtttaaa | ggatctgtgc | accctcatct | ggaagcaaag | 1260 |
| ctccgccttc | ggctgctgtg | gcccgcctgc | ccgcagagag | gcttgctggg | aggtccaggc | 1320 |
| cgcagttccc | gtccagctcc | tggagtcaga | gctgcaggat | ggctgttggg | ggtccttgaa | 1380 |
| gcctgtccag | atgcacttgg | gatgggtttt | tttgttttgt | ccagactgtt | tctgtagatt | 1440 |
| cctaataataa | tgagccatta | taattttgaa | caattttgtaa | agagtacttc | cagaggaacc | 1500 |
| tgtaaacat | tcaaccatta | ggtctttgct | gtaagtgtct | gttttttaatt | ttctggcagt | 1560 |
| gcttctgaga | attctgcaaa | gcctcaagaa | aaaatagagt | aagctgtctt | agccaggccc | 1620 |
| ctgttagcttt | ggtcacagca | cttccctggtg | tgacaaggac | tcagaccccc | cgggagcccc | 1680 |
| ttggctatcc | tcagcactgt | ttattcctcc | tcaggctctg | cctggcaccc | taccctccct | 1740 |
| ggcactcttt | ctttgtcagg | ccactcgttg | ccaccctgag | actccagctc | atgtctgatt | 1800 |
| tgtgcatcct | acttaactga | tgtgtttttc | ctcctctctg | tgtgtgagga | tgtgtgcacc | 1860 |
| acctcacatt | ggctcttgag | tgtgaagtgt | cttgctgcac | ggcagaaaca | gacctagaga | 1920 |
| agtcccttgg | tttggccagt | attgtgttgc | tgacaggcgg | tggatcaaga | gttaaaatgt | 1980 |
| aggaagtcca | tgttctgtcc | cctttgacaa | gccattttct | taatctgtac | aaagggaaca | 2040 |
| atgtctgcac | cacgaggttg | gggagatttt | cagtgaagat | ggatgtgaag | agtgtttcgt | 2100 |
| aaaccataaaa | gcagtttatt | gtcattgatg | attctcagct | ttgcatgaac | ctgagggaca | 2160 |
| ccagaccact | tgctctagga | gcccataattc | tgggttcttt | gtcctaaact | actttttggg | 2220 |
| ccttttctcta | aaaatacaca | ttctaagagt | caaggtaaga | gtgttcctaa | tttgggtgcat | 2280 |
| ggatgagagt | tcattgttagt | tttcatttag | gctggaagtt | aaacctgagc | cattttcctg | 2340 |
| tccagctgca | ctttgccaca | ggctggggcac | actggctcat | acctgtaatc | ctagcacttt | 2400 |
| gggagactga | ggaggattac | ttaagaccac | gtgttcaaga | ccagcctagg | agtatagtga | 2460 |
| ggtgccatct | cttttaattt | ttaaaaaata | gtcctagtga | ctcttaccag | gccagtcacc | 2520 |
| atctccagga | gccccctgt | gggagtatgt | tcagtgtctg | ccctttgtga | agtggggcct | 2580 |
| tttgtccgc | cacctacccc | cagggcactt | aaggcctggg | tgtgtttctg | ccggcagggg | 2640 |

| | | | | | | |
|-------------|-------------|------------|------------|-------------|------------|------|
| cttgaaggtg | tctggcagtg | aaggaaggag | aggatgaagg | gaggctggag | agggagtggc | 2700 |
| gagcggccag | aggcaggcgg | ggccgtggga | agggaagca | gcacaggcaa | gctgagtga | 2760 |
| cattggtgat | gagatacaga | gatccatgat | tgaacttcta | gtgtatctta | gcccactggg | 2820 |
| gctgccatga | caaaacacca | tagagtgtgg | ctcatcaaca | gcaggagtct | gttgctcaca | 2880 |
| gctctggagg | ctggagggtcc | agggtcaagg | cactggcaga | ttgcatgtct | ggtagggccc | 2940 |
| cgcttcctgg | tccatagcca | gcaccttctc | actgcaccc | tatgtggtgg | aagaggcagg | 3000 |
| ggagctctct | ggggctctct | ttgtaacccc | attcttgagg | cttggtccact | ctcatgacct | 3060 |
| aatcacctcc | cagaggccct | gcctcctaac | accattaccg | tagagtgaga | gcttcagcct | 3120 |
| atgaattttg | gggggacatg | aacattcaag | actaatattc | tttcagtatt | ccagttactg | 3180 |
| aatggtgtgt | agtattcagt | ggagaaggta | gctgaaatta | aggggttaaa | attgtaataa | 3240 |
| tggccggttg | cggtggctca | cgctgtaat | cccggcactt | tgggaggctg | aggccagtgg | 3300 |
| atcacaaggt | caggagtctg | agatcagcct | gaccaacatg | gtgaaacccc | gtctctacta | 3360 |
| aaaatacaaaa | aaattagctg | ggcgtggtgg | cacacgcctg | taatcccagc | tactcaagag | 3420 |
| gctgaggcag | gagaatcgct | tgaacctggg | aggcagaggc | tgcactgctg | agctaagatc | 3480 |
| gtgccactgc | actccagcct | gggcgacaga | gcgagactcc | atctcaaaaa | aaaaaaaaaa | 3540 |
| a | | | | | | 3541 |

<210> 17

<211> 1687

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (1568)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (1652)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (1654)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (1660)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (1661)

<223> n equals a,t,g, or c

<400> 17

| | | | | | | |
|------------|------------|------------|------------|-------------|------------|-----|
| ccacgcgtcc | gggcagtggg | gtgagggcac | acaagcagtt | caggggtccca | gcaggaagtg | 60 |
| gggctgcagg | gccgggggtg | gtcctgggcc | tggccatcag | gcagcctagc | agggtgttct | 120 |
| gggcatggag | ggggcctggt | gtggctgagg | gcatgccag | ggctccctgg | aggatcccgc | 180 |
| tctgtgccct | gccaccctg | tgcctgggga | gccctctgcc | ctcacagccc | acccacccca | 240 |
| ttttctatga | ccacagagct | ccgacctgga | agatggctca | cccaggaggt | cccaggagct | 300 |
| ctcactcccc | caggggacct | ggaggacacc | cagctctcag | acaaaggctg | ccttgccggc | 360 |
| ggggggagcc | cgaaacagcc | ctttgcagct | ctgcaccagg | agcagggtttt | gcggaacccc | 420 |
| catgtaaggc | ttccccgggg | tggggtcctc | ccagccgtgg | gcctcagggt | gaccgatcac | 480 |
| agggagagtg | gtccctgcc | ctgggcaccc | cctgcggtgg | ccccgacgac | agctgaggag | 540 |

| | | | | | | |
|-------------|------------|------------|-------------|------------|------------|------|
| tgaccacaag | gtctctgccc | acagtgtctg | gggtgcggtg | tctgggctgc | gaagtggatc | 600 |
| ccccctcttt | cttgggcact | gcagcagctt | ggggggcttt | ttggacgtgg | atgtgcctgg | 660 |
| tccttggtttc | ccgagggcct | ttacagtggg | tgaggagggtg | aacacagggg | gtcctgagag | 720 |
| caagcaccac | ctcgggcttt | ggtgtagaaa | caatggcccc | gaccccaggc | cggagccgtg | 780 |
| gcttggcctc | ctgggtgtgt | cttggcatct | gaaatgcagg | ctacccacac | cggctcacct | 840 |
| ccaggggtac | aggcagggtc | cacagggaga | gcttggcgct | gagctgaggc | tgtctgggct | 900 |
| cctcgcctcc | caaccagtct | gcagttacag | gggccagtgg | atgggcgggt | gagaaggacg | 960 |
| ggttccctca | ggggagcccg | ccggagcccg | agccttcccc | cttctccagg | acgcaggcct | 1020 |
| gagcagcggg | gagccgcccg | agaaggagcg | gcggcgccctc | aaagagagtt | ttgagaacta | 1080 |
| ccgcaggaag | cgcgcctca | ggaagatgca | gaaaggatgg | cggcagggtg | gaggaggacc | 1140 |
| gggagaacac | cacgggcagc | gacaacaccg | acactgaggg | ctcctagccg | cagcagccgc | 1200 |
| aggccccgac | cagggcacac | ccaccggccc | ggcctcctgc | cacccggggg | tgccgacgcc | 1260 |
| ctggggcgca | gacttccccg | agccgtcgct | gacttggcct | ggaacgagga | atctggtgcc | 1320 |
| ctgaaaggcc | cagccggact | gccgggcatt | ggggccgttt | gttaagcggc | actcattttg | 1380 |
| cggaggccat | gcgggtgtc | accaccccc | tgcacacgcc | atctgtgtaa | cttcaggatc | 1440 |
| tgttctgttt | caccatgtaa | cacacaatac | atgcatgcat | tgtattagt | ttagaaaaa | 1500 |
| cagctgcgta | aataaacagc | acgggtgacc | cgcaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | 1560 |
| aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | 1620 |
| aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | ananaaaaa | naaaaaaaa | aaaaaaaaaa | 1680 |
| aaaaaaa | | | | | | 1687 |

<210> 18

<211> 2075

<212> DNA

<213> Homo sapiens

<400> 18

| | | | | | | |
|-------------|-------------|------------|------------|-------------|-------------|------|
| ccacgcgtcc | gtactgcctt | ttttcaacct | cagatgtgac | tttcattata | ggaagtcttc | 60 |
| aggcattttc | tcttgggaata | atacctcttc | tctcttctct | ttatgtcctt | gtgccgcatt | 120 |
| ctgggttatt | ccttttagctc | taggttaagt | tcactaatct | ttccttttagc | tgtatttcat | 180 |
| tattgtttta | gctgtccatt | gcatttttaa | ctttctttca | aatatcttcc | cttcccttcc | 240 |
| tttcccttct | cttccctgcc | ctgccctgcc | ctgccctgcc | ctgccctccc | gtccctccc | 300 |
| ctccctgcc | ctccctccc | ctccctccc | ctcccttgc | cttccctctc | ttttttttaga | 360 |
| taaagcctgg | gcaacatag | gagaccccat | ctctacaaaa | aaatttttaa | ttagttgggc | 420 |
| atggtgttaa | gtgcctgtag | tcccagttat | tcaggagtct | gatgtgggag | gttggcttga | 480 |
| gcttcagagg | caatggttgc | aatgagccaa | ggtcacacca | ctgcactcca | tcctgggcaa | 540 |
| cagagccaga | ccctgtctca | aaaacaaaac | aaaacaaaag | aacataccaa | aagcactacc | 600 |
| cataaagaac | caattgataa | attggacttt | atcaaaaatt | aaaacttctg | ctgttcaaaa | 660 |
| acattattaa | aagaatgaaa | agacaagcca | ttaactagga | gaatataact | tcaaatcaca | 720 |
| catttaataa | aagacttata | tccagaacat | atgaagaact | ttctaaagt | aataagaaaa | 780 |
| aaaaaaccca | ataaaaatag | gggaaatatt | ttactagaaa | cttcaaaaaa | gacatacagt | 840 |
| aggcaaatga | aaagatcatt | agtcattagg | gaaatgcaaa | ttaaaaccac | aatgagatag | 900 |
| caatacatat | ctactagaaa | gcctaaaatt | aaaagaactt | tttatattat | gtgttgacga | 960 |
| gaatgtagag | caactagaac | tgcaaatggg | agtataaatg | atacaacaac | tttgggaacat | 1020 |
| atttcgatag | tttcttttaga | aaacatacat | taacctgcc | catgattcag | ttatcccatg | 1080 |
| cctagggtatt | tactcaaaaag | aaatgagggg | atattctatg | caaagggtgt | aaaataatgt | 1140 |
| tcataacagt | tttacatgtc | aaagccaaaa | acttgtaatt | acccaaatag | ccatcaacaa | 1200 |
| gtgaatagat | aagtaaactg | tgatatatcc | atacaatgga | atactattca | acaataaaaa | 1260 |
| gaatgcagct | acatgaatga | atctcaatta | tgctagttaa | agaagacatt | taaaatcaat | 1320 |
| acacactata | tgattctgtt | tattttaa | tctataaatg | caagataaga | tacagtaaat | 1380 |
| aaaagattcg | tggttgctcg | gggattgggg | gagggagcaa | agagaataga | aatgtgatta | 1440 |
| caaagggtgtg | atataataag | aaatatatac | agatgcttct | caatttatga | tagcagggtt | 1500 |
| acctccta | aaacctatca | taaattttaa | ataccatgtc | aaaaatgtat | ttaagacacc | 1560 |
| taatctgctg | aacatctcat | tgcttagtgt | agcctacctt | aatgtgtctc | agaacacatt | 1620 |
| agcctacagt | tggaacaaa | catctaacac | aatccctatt | ttataataaa | gtgttgaata | 1680 |
| tctcatgtaa | tttattgaat | actaaagtga | aaaacagtat | aagtattgaa | gtagtttgtg | 1740 |
| gaaataagta | attcaaaatc | taagctgggc | tggtgtgtgt | agctcacgcc | tgtaatccca | 1800 |

| | | | | | | |
|------------|------------|------------|-------------|------------|------------|------|
| acactttggg | aggctgaagc | gggtggattg | cttgagggtca | ggagttcaag | accagcctgg | 1860 |
| ccaacatggt | gaaaccctgt | ctctactaaa | attacaaaaa | ttagccaggc | atggtggcac | 1920 |
| atgcctgtaa | acccagctat | gtcaggaggt | tgaggcacaa | gaattgcttg | aaccaaggag | 1980 |
| acagagggtg | caatgagccg | agattgcgcc | actgcactcc | agcctgggtg | acacagagag | 2040 |
| gctctgactc | aaaaaaaaaa | aaaaaaaaaa | aaaaa | | | 2075 |

<210> 19

<211> 4699

<212> DNA

<213> Homo sapiens

<400> 19

| | | | | | | |
|-------------|------------|-------------|------------|------------|-------------|------|
| ccacgcgtcc | gttctgtcca | tgtcagcatc | cctgccttaa | gagaactcct | cctggccaat | 60 |
| ggcattgcc | cccagcagt | ggaccaagac | tctccaagac | ctccaggact | ggatcccatt | 120 |
| gcctggagaa | actccagcaa | gggtctctca | tggcttgagc | atggcacagt | aaggggcagc | 180 |
| caaccagtc | catgatgact | tttgctccaa | cttcttcatt | tttctaaaag | cccagtggct | 240 |
| ttattcactc | ctcctaaatt | gcctgctacc | agaaggaact | tcacctgaa | gaaatgcatt | 300 |
| ccattaccag | ttccagggaa | agtgtccctt | tcccaaagc | ctcaggcccg | tgggcctctg | 360 |
| aggttccact | ggatgcggct | ccccaggag | tgggcctgga | gatccgctca | gccgcctgc | 420 |
| ctcccactct | ctatcttggg | accgtgggtc | gccctgaagg | gtggttccag | ccccgcgtat | 480 |
| gctgcacttt | gctgctgcag | gtccgggtcc | ctccagggtt | tttcaatgag | agttcccacc | 540 |
| ccaacttgg | gcatttcatt | tttgcttacc | taaagcaaga | atctcaaagt | ttgtttgaat | 600 |
| aagaggccca | ttcacaagtc | gtgcccttgt | gagcaccctt | ttgctgagtg | cctcgtgggt | 660 |
| gccaaacact | gtacttggcg | ctttgcattc | atgctttgac | cctcacaacc | tccctttata | 720 |
| gatgagga | ttcacgctcc | atgtgcccaa | ggacacacca | gccaggagga | agagggaaga | 780 |
| ggatttgagt | ccaggcctgc | ctgacttcag | tgctgtgtcc | ctcccctctg | taaaagacaa | 840 |
| accaggcagg | gaagagatag | aagctggcta | agtgggtggc | aagtggcttc | attgtgaggg | 900 |
| gggtgagggg | gcattcgtgg | ctgctccac | accaccatc | cctccagccc | acagcagcct | 960 |
| ataccaaagg | ctgccccgga | ctcaacagct | gcgtctccat | gacaggagaa | tgccagggtc | 1020 |
| ctgggtgggtg | gcacagagcc | ctgttccctg | cttctaagct | gttgccccct | ggggaataga | 1080 |
| gcttatggag | caacgggagc | tgaggtggcc | tcgagggcca | gcctggaggg | ctctgagcat | 1140 |
| agcatgcagc | ggtcccatag | ggaggcaggt | aaccaagggc | caggcagagc | tgcttccctgt | 1200 |
| gcctctccgg | catttgtaat | ggttttttct | tttttcacgc | actgccaaat | ctgttttctc | 1260 |
| ccttgaagaa | acccagggca | gatgtgggtc | tctgtcaaac | tgaggtctgg | agaggcatga | 1320 |
| ctaaatcagc | gcagaagcaa | aatcaggacc | cagcatttct | agctcccagc | ccagagtaag | 1380 |
| agagccagct | agaacccctt | cccattctac | ccgaccgagg | cagccctgcc | ctgccctgcc | 1440 |
| ccgccccctc | ccaccctgcc | ctgccccaga | gatctccagt | tcacagtgtt | catcaggcga | 1500 |
| gatacttttg | gacccacaaa | gccagttttt | tggggggatg | ctcagtgtct | acctgcagtg | 1560 |
| ccaaagtgtt | gcttgaacag | ccctcttctg | ccgcgcctcc | cagctccacc | ccctatcaac | 1620 |
| gtgggtgctg | gctgcttctg | cgtgaggagc | ttcctctgat | cacctgtagc | atcttttagtg | 1680 |
| ccttcgtcca | gtcggcctgc | aaatcccatt | tcccagagag | gcttgggcag | ggcccacttg | 1740 |
| ccacttggcc | ctgtgcccag | agcgttgggg | gcagatggca | gccaggggaa | gaaggaggtt | 1800 |
| ggggaagggg | cgccaccgct | tgagggtccc | ctcccttgag | aggaggcccc | atggcagttt | 1860 |
| tcttgagttc | ttgatccact | ttggttttaa | caacttttgt | gaagctatgt | gagatttgac | 1920 |
| tgcatttcaa | aagacaaaac | acatgttttt | ttctgatttg | ctctttgcct | ttcagcgacg | 1980 |
| ctttcagata | ctttgtgagt | gtctactctg | tgctaggtgc | tgagtagact | gggatgtgtg | 2040 |
| tggctccatc | cctggctgga | agagctttga | gtcctgtctg | ctgacttcac | tgaaggagtt | 2100 |
| tctctcttgc | ttcatctctt | cctcccgggg | gtcagctggg | taggagcaga | cttgccctgc | 2160 |
| cttctcttag | gcagaaagtt | ttcctgcaaa | tgaaagaatc | tgccattctc | ttttgctatg | 2220 |
| atgatgatga | ttattctgac | atgaccgcg | ggtactgaag | ccacagtctt | ctctgttcac | 2280 |
| tggtctctcc | ttgagtcata | tctatggcct | gaggacgtct | gtatccagct | ccttttccta | 2340 |
| caatggtaga | cacttcccca | ggcagtggtt | gggtcggggt | ggccagtcac | tctcctagct | 2400 |
| gtccttgctt | ggggacttag | atatacccca | tccaggactg | tggtcgctat | tctgtgagcg | 2460 |
| attgatggcg | catatgcaga | actcccctgg | agtcattggt | ctagaaagtg | tccaggccca | 2520 |
| tcgtgacaat | ggtgccaaag | accgggctgg | tgaccagtgc | ctaaacacac | acgctccctt | 2580 |
| aatccttacg | gcaagcctgt | gaggcgagata | ggaccagccc | tgatctatag | tgagaaaatt | 2640 |
| aaggttcaga | gggagaaggt | ccaaatagtg | gccaagctgg | gactgagatc | tcgagtttgg | 2700 |

| | | | | | | |
|-------------|-------------|-------------|-------------|-------------|-------------|------|
| gtttgtttgg | gtttgtttgt | tgttgtttctt | tccctcgtgt | ccatgacttt | tctctgccat | 2760 |
| tagcagggaa | gtgtgcggag | ggcttccagg | gaatcaagt | accctccctc | tgaatttcgg | 2820 |
| tgcctcttcc | ccccgcccct | tgctcacaca | tttcagtagg | tcacaggcca | aatggcagcc | 2880 |
| caaaaggtgg | gcagagattt | tttttttctt | cccctcacc | ctgcttatta | acacaattgt | 2940 |
| gacaactact | ttaccttaca | tcccagggca | aacggacagc | ttgcgggtacc | tccaaaaatt | 3000 |
| tggaaaacct | tgagtctaag | accacctctt | gataaaagag | gcctttgctc | acattttctg | 3060 |
| ctcggaatct | ttctttgcc | ttgggtgggtg | tatgtcgtcc | cagctgagac | tgcgggagag | 3120 |
| ctgagctcca | gcttaaaccg | cttttaaatgg | cccttcccta | ctgggggatc | gctggagccc | 3180 |
| atgccagggt | acggtagtca | ttttaaacct | ctgccacaag | gccacgccat | agaatcaatt | 3240 |
| tccaagtcaa | tgttcttggc | cctaaaatgt | ttgcgcccac | tgaggatttc | attaccaggt | 3300 |
| ggaagatggg | ggaacatatg | ccacgttaaa | gagcaagtgc | tgagtgtgcc | aggacttgag | 3360 |
| aaggacctta | ggctgaggag | tggacagtct | ggtattttatt | tcgcctctcc | caagtaccca | 3420 |
| gcacagagtg | ggaaaaagaa | gaaactgctc | ctgccttcca | gtagctggca | acttgacatc | 3480 |
| aagaacaaaa | acattcgtag | ctggaccccc | gtcctttcat | cgtgcccag | tcttacatgt | 3540 |
| tggcttaagc | tgcccccttg | gaggccgtac | ctttctcttt | ggactgtgac | agtggacacc | 3600 |
| taacagcctg | agccatgtgt | ggctgtctta | tcccccaagg | gttgtccagg | cccgttctgc | 3660 |
| agggctggtg | tgtagactgc | agacatccat | acctcaccac | agaccaaaga | tgacctcgtg | 3720 |
| tcagactgtg | ggctgatgag | aggtagagca | gcattgcacg | aggcctgagg | gtgcagggcg | 3780 |
| ccctctcttg | gcctggagga | attgctccta | actagagtaa | gtttccacga | gggtcccagg | 3840 |
| cagagctgca | gagctggaac | cggaggctcc | acagtccctg | cctgctcatg | gacctccttc | 3900 |
| agagcacctt | tctacagact | ggactgcccc | gtcccggtgg | gtggcatctg | gtttctggtg | 3960 |
| ctattctgcc | aagtattcga | gtctctcctc | atgtttcaac | attccatctt | cccgtttcta | 4020 |
| tcctcgactc | caaagtaagc | cttcttagct | ccaatcaggg | atgaggggct | caacctcttc | 4080 |
| tgtcctcaaa | gaggccaaac | gcagtgccac | agtcggtagc | cttcactttt | agatgtccta | 4140 |
| ttcatgtaaa | aaagaagggtg | ccccaccag | gcttacatca | gcaataagca | attctaattgc | 4200 |
| aacgatgggtg | tccacatttt | accccagtg | gtgcccatgt | atgcctttgt | gcccgtgtaa | 4260 |
| ttattgttag | cgcccccttc | acttagaggg | gtgatgataa | actgtggcca | ccttgattac | 4320 |
| aaccacatt | tcctgctttg | gggagcttcc | aagtaacagg | ccatttctta | cctccctcca | 4380 |
| ggaacagtgg | gcaactgcca | ccacctcgtg | tctgctcata | ggatgacgct | ggagatcccc | 4440 |
| acacttactc | tacctctttg | gcaaattggc | attccgggtg | tggtttttgt | ttcctttaac | 4500 |
| acattaaata | aatgagtata | taggatgtga | ggggaggggt | gagaacaact | agctgtagca | 4560 |
| tgtgtaggct | atatacttta | ccattcgact | tctttccttt | ttttttttta | aataaaaaaa | 4620 |
| gtgcttgact | ggtttcaagc | ttcatcatga | aaaaaaaaa | aaaaaaaaa | aaaaaaaaa | 4680 |
| aaaaaaaaa | aaaaaaaaa | | | | | 4699 |

<210> 20

<211> 2762

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (2711)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (2730)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (2752)

<223> n equals a,t,g, or c

<400> 20

ccacgcgtcc gcggagtcgg tgggtgcaga tggcggcggc agttgtggtg gcggaggggg

60

| | | | | | | |
|-------------|-------------|-------------|-------------|-------------|-------------|------|
| acagcgactc | ccggcccggg | caggagttgt | tagtggcctg | gaacaccgtg | agcaccggcc | 120 |
| tgggtgccgc | ggctgcgctg | gggctgggtg | cttcccggac | cagcgggtgca | gtcccgccaa | 180 |
| aggaagagga | gctccgggcg | gcgggtggag | ttctgagggg | ccacgggcta | cactcgggtcc | 240 |
| tggagagtg | gttcgtggag | gtgctgcaga | acgatctgca | ggccaacatc | tccccgtagt | 300 |
| tctggaatgc | catctcccaa | tgcgagaact | ctgcggatga | gccccagtgc | cttttgctac | 360 |
| tccttgacgc | ttttggcctg | ctggagagcc | gcctggatcc | ctacctgcgt | agcctagagc | 420 |
| tgctggagaa | atggactcgc | ctgggcttgc | tgatgggcac | tggtgctcag | gggctgcgag | 480 |
| aagaagtcca | cactatgttg | cgcggagtct | tgttcttttag | cacccccaga | accttccaag | 540 |
| agatgatcca | gcgtctgtat | gggtgcttct | tgagagtcta | tatgcagagt | aagaggaagg | 600 |
| gggaaggggg | cacagaccgg | gaactggaag | gggagctgga | cagccgggtat | gcccgtcgcc | 660 |
| ggtactaccg | gctcctgcag | agcccgtgtg | gtgcaggggtg | cagcagtgac | aagcaacagt | 720 |
| gctggtgtcg | ccaggctctg | gagcagttcc | atcagctcag | ccaggctcta | cacaggctca | 780 |
| gtctgctgga | gcgggtcagt | gccgaggctg | tgaccaccac | cctgcaccag | gtgacccggg | 840 |
| agaggatgga | ggaccgttgc | cggggcgagt | acgagcgctc | cttcctgcgt | gagttccaca | 900 |
| agtggatcga | gcgggtgggt | ggctggctcg | gcaagggtgtt | cctgcaggac | ggccccgcca | 960 |
| ggcccgcatc | tcccagggcc | ggcaacaccc | tgcgcgctg | gcgctgccac | gtgcaaagggt | 1020 |
| tcttctaccg | catctacgcc | agcctgcgca | tcgaggagct | cttcagcatc | gtccgagact | 1080 |
| tcccagactc | ccggccagcc | atcgaggacc | tcaagtactg | cctggagagg | acggaccaga | 1140 |
| ggcagcagct | gctcgtgtcc | ctcaaggctg | ccctggagac | tcggctcctg | catccaggcg | 1200 |
| tcaacacgtg | tgacatcatc | accctctata | tctctgccat | caaggcgctg | cgcgtgctgg | 1260 |
| acccttccat | ggatcatcctg | gaggtggcct | gtgagcctat | ccgccgctac | ctgaggacgc | 1320 |
| gggaggacac | agtgcggcag | attgtggctg | ggctgacggg | ggactcggac | gggacagggg | 1380 |
| acctggctgt | tgagctgtcc | aagaccgacc | cggcgagcct | ggagacaggc | caggacagtg | 1440 |
| aggatgactc | aggcgagcca | gaggactggg | tcccggaccc | tgtggatgcc | gatccaggga | 1500 |
| agtcgagctc | caagcggcgt | tcatacgaca | tcatacgcc | gctggtcagc | atctacggca | 1560 |
| gcaaggacct | cttcatcaat | gagtaccgct | cgtgctggc | cgaccgcctg | ctgcaccagt | 1620 |
| tcagcttcag | ccccgagcgg | gagatccgca | acgtggagct | gctgaagctg | cgctttggcg | 1680 |
| aggccccaat | gcacttctgt | gaagtcatgc | tgaaggacat | ggcggactcc | cgccgcatca | 1740 |
| atgccaaacat | ccgggaggag | gatgagaagc | ggccagcaga | ggagcagcca | ccgttcgggg | 1800 |
| tctacgctgt | catcctgtcc | agtgaattct | ggccgccctt | caaggacgag | aagctggagg | 1860 |
| tccccgagga | tatcagggca | gccctggagg | cttactgcaa | gaagtatgag | cagctcaagg | 1920 |
| ccatgcggac | cctcagttgg | aagcacaccc | tgggcctggg | gaccatggac | gtggagctgg | 1980 |
| ccgaccgcac | gctgtctgtg | gcggtcaccc | cagtacaggc | ggtgatcttg | ctgtattttc | 2040 |
| aggaccaagc | cagctggacc | ctggaggaac | tgagcaaggc | ggtgaagatg | cccgtggcgc | 2100 |
| tgctgcggcg | gcggatgtcc | gtgtggctgc | agcagggtgt | gctgcgtgag | gagccccccg | 2160 |
| gcaccttctc | tgctattgag | gaggagcggc | ctcaggaccg | ggacaacatg | gtgctcattg | 2220 |
| acagtgcaga | cgagagcgac | tccggcatgg | cctcccaggc | cgaccagaag | gaggaggagc | 2280 |
| tgctgtctct | ctggacgtac | atccaggcca | tgctgaccaa | cctggagagc | ctctcactgg | 2340 |
| atcgtatcta | caacatgctc | cgcattgttg | tgggtgactg | gcctgcactg | gccgagattg | 2400 |
| acctgcagga | gctgcagggc | tacctgcaga | agaagggtcg | ggaccagcag | ctcgtctact | 2460 |
| cggccggcgt | ctaccgcctg | cccaagaact | gcagctgaca | catcgcccgc | ccgcccgcgc | 2520 |
| gcccgcagag | cgctgccctg | cagggtgctct | cgtcctcccg | tgccagcccc | cgcccgcgcc | 2580 |
| tgctccagaa | tgcactgctg | aggagcatgc | ccacccccac | ccccgcagtg | tgcaatttaa | 2640 |
| agcaagtcag | atcatcaaaa | aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | 2700 |
| aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | 2760 |
| at | | | | | | 2762 |

<210> 21

<211> 1133

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (1061)

<223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1078)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1102)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1107)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1111)
 <223> n equals a,t,g, or c

<400> 21

| | | | | | | |
|-------------|------------|-------------|-------------|------------|-------------|------|
| ggggcgggcg | gctccctgcc | tgctgctggg | cggaggggaag | gcggaagag | ctgaggagcc | 60 |
| cctggaagag | ctccaggaa | ccctgcgctg | tgggataaag | gaatgagggt | cagaaagggg | 120 |
| cagggagttg | cccgcagccg | caccgcacgt | cttcagcccc | accgttgtcc | tgacctctct | 180 |
| gtcccggtccc | ctgcccagtc | tcaccatggc | cttctggaca | cagctgatgc | tgctgctctg | 240 |
| gaagaatttc | atgtatcgcc | ggagacagcc | ggtccagctc | ctggtcgaat | tgctgtggcc | 300 |
| tctcttcttc | ttcttcatcc | tggtggctgt | tcgccactcc | cacccgcccc | tgagcacca | 360 |
| tgaatgccac | ttcccaaaca | agccactgcc | atcggcgggc | accgtgccct | ggctccaggg | 420 |
| tctcatctgt | aatgtgaaca | acacctgctt | tccgcagctg | acaccgggcg | aggagcccgg | 480 |
| gcgcctgagc | aacttcaacg | actccctggt | ctcccggctg | ctagccgatg | cccgcactgt | 540 |
| gctgggaggg | gccagtggcc | acaggacgct | ggctggccta | gggaagctga | tcgccacgct | 600 |
| gagggctgca | cgagcacgg | cccagcctca | accaaccaag | cagtctccac | tggaaaccacc | 660 |
| catgctggat | gtcgcgagc | tgctgacgtc | actgctgcgc | acggaatccc | tgggggttggc | 720 |
| actggggccaa | gcccaggagc | ccttgcacag | cttggtggag | gccgctgagg | acctggccca | 780 |
| ggagctcctg | gcgctgcgca | gcctgggtgga | gcttcgggca | ctgctgcaga | gaccccgagg | 840 |
| gaccagcggc | cccctggagt | tgctgtcaga | ggccctctgc | agtgtcaggg | gacctagcag | 900 |
| cacagtgggc | ccctccctca | actggtacga | ggctagtgc | ctgatggagc | tggtggggca | 960 |
| rgarccagaa | tccgcctgca | gacagcagct | gagcccgtg | ctcggagctk | attggagcct | 1020 |
| ggacagcacc | cgggtgtccg | tggtctggaa | cgctgaagct | ntatcctcgg | aagtactntt | 1080 |
| gacagatcac | tttaccgaag | tnatggncag | ngaacgactt | cagagttacc | tgt | 1133 |

<210> 22
 <211> 1714
 <212> DNA
 <213> Homo sapiens

<400> 22

| | | | | | | |
|-------------|------------|------------|------------|-------------|-------------|-----|
| agccgctcct | gccgcctg | tcgctggagc | tttgctcttc | taggcccggca | gcgctctctc | 60 |
| tccatgggtcc | tgtctgtcag | cgctgttttg | ggagcccgc | ggtgaggccg | ggccacgctc | 120 |
| agacacttcg | atcgtcgagt | ctgtcactgg | gcatggcggg | tcagttccgc | agttacgtgt | 180 |
| gggacccgct | gctgatcctg | tcgcaatcgt | cctcatgcag | accgtgtatt | acggctcgct | 240 |
| gggcctgtgg | ctggcgctgg | tgagcgggct | agtgcgaaca | gcccctcgct | ggaccagatg | 300 |
| ttcgacgccg | agatcctggg | cttttccacc | cctccaggcc | ggctctccat | gatgtccttc | 360 |
| atcctcaacg | ccctcacctg | tgccctgggc | ttgctgtact | tcatcacggc | gaggaaagca | 420 |
| gtgtctggat | ttactgtca | ctgtccattt | ctttcacctc | ctgggctgct | ggttctacag | 480 |
| ctcccgtttc | ccctcggcgc | tgacctgggt | gctgggtcaa | gccgtgtgca | ttgcaactcat | 540 |
| ggctgtcatc | ggggagtacc | tgtgcacg | gacggagctc | aaggagatac | ccctcaactc | 600 |

| | | | | | | |
|-------------|-------------|-------------|-------------|-------------|-------------|------|
| agccccctaaa | tccaatgtct | agaatcaggc | ccttttgaca | tcctgctgac | acttggggccc | 660 |
| ctttaacacc | ttgggctgct | cagaccctcc | agatgaggtc | cagcccagat | ctgagaggaa | 720 |
| ccgggaaatg | tgaagtctct | gttgggtgtg | gagagatagt | gagggcctgt | caaagaaggc | 780 |
| cggtagcagt | cagcatgaca | gctgccagaa | tgacctctgt | ctgttgaagc | cttgggtatct | 840 |
| gagaggtcag | gaaggggacc | tctttgaggg | taataacaga | attggaacca | tgccactctt | 900 |
| gagccacaat | acctgtcacc | agcctgttgt | tttaagagag | aaaaaaaaatc | aaggatatct | 960 |
| gattggagca | aaccacttct | ttagtcattct | gtcttaccct | cctgggacag | ctgttacctt | 1020 |
| tgcaagtgtg | ccgaatcaca | gcagttacct | ttgcagtgtt | gccgaatcac | agcagttctg | 1080 |
| ttggagaaac | gcttggtttc | cggatccaga | gccacagaaa | gaaatgtagg | tgtgaagtat | 1140 |
| taggctgctg | tcaggagag | gatggcagat | ggaggcatca | agcacaagga | aaatgcacaa | 1200 |
| cctgtgccct | gtttatacaca | cgttcattgtg | caccacaagaa | cctatgactt | tcttccagtt | 1260 |
| ccttctacca | gggtcccatc | ctgctgccag | ctctcaacat | agcaaggcca | taggaccag | 1320 |
| ataagaatcc | cagcgttgct | caaagtctaa | ccattcataa | aagacactgg | cctgtcttct | 1380 |
| aggaatgacc | aggcacccca | gcttccactg | gacttccaat | tttttttttc | ctggccttaa | 1440 |
| tttaagaatt | ctttggcggg | aagggatga | tgggttccca | gagacaagaa | gccccacctt | 1500 |
| ctggcctggg | ctgtgctgat | agtgtgagg | gagataggaa | tttgctgcta | agatttttct | 1560 |
| ttggggtgga | gtttcctctg | tgaggggctt | gcagctatcc | ttcctgtgta | tacaaatata | 1620 |
| gtattttcca | tggttcaaaa | aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | 1680 |
| aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | aaaa | | | 1714 |

<210> 23

<211> 2679

<212> DNA

<213> Homo sapiens

<400> 23

| | | | | | | |
|-------------|------------|------------|------------|------------|------------|------|
| ccacgcgtcc | gcctcagcgg | ccggggccac | ggccccgagc | agccatgctg | ggcgcgcggg | 60 |
| cctgggttggg | ccgcgtcctt | ctgctgcccc | gcgcgggtgc | aggcctcgcc | gcaagccgca | 120 |
| gggtgtcctgg | agtctggccc | aggacctggc | cccacaggag | tcccagcagg | ggtagctcct | 180 |
| ccggggacaa | ggaccgaagt | gcgacggtea | gtagtccagt | gcccctgcct | gctggaggga | 240 |
| aaggaagcca | tccttcatct | acaccccaga | gggtcccca | ccgcctgac | cacgagaagt | 300 |
| caccatacct | cctacaacat | gcctacaatc | ctgtggactg | gtacccctgg | ggacaggaag | 360 |
| ccttcgacaa | ggccaggaag | gaaaacaagc | cgattttcct | ctcagtcggg | tactccacct | 420 |
| gccactgggt | ccacatgatg | gaagaggagt | ccttccagaa | tgaggagatt | ggccgcctgc | 480 |
| tcagttagga | ctttgtgagt | gtgaaggtag | accgtgagga | gcggcctgac | gtggacaagg | 540 |
| tgtacctgac | gttcgtgcag | gccaccagca | gcggcggggg | ctggcccatg | aatgtgtggc | 600 |
| tgactcccaa | cctccagccc | tttgtcgggg | gcacctatct | ccctcctgag | gatggcttga | 660 |
| cccgagtctg | cttccgcaca | gtgttgctga | gaatacgaga | acagtggaaa | cagaacaaga | 720 |
| acaccctgct | agaaaatagc | cagcgtgtca | ccactgccct | gctggcccga | tcagagatca | 780 |
| gcgtgggtga | ccgccagctg | ccgccctctg | ccgccaccgt | gaacaatcgc | tgcttccagc | 840 |
| agctggatga | gggctatgat | gaggaatacg | gtggcttcgc | tgaggccccc | aagtttccca | 900 |
| cgccgggtgat | cctgagcttc | ctgttctcct | actggctcag | ccatcgactg | actcaggatg | 960 |
| gctctcgggc | ccagcagatg | gccttgcata | ccctgaaaat | gatggctaac | gggggcatcc | 1020 |
| gggacctatg | ggggcagggc | tttcaccgct | actccacaga | ccgccagtgg | cacgtccctc | 1080 |
| actttgagaa | gatgtcttat | gaccaggcac | agctcgctgt | ggcctattcg | caggccttcc | 1140 |
| agctctctgg | tgatgaattc | tactctgacg | tggccaaagg | catcctgcag | tacgtggctc | 1200 |
| ggagcctgag | ccaccgggtc | ggaggcttct | atagcgcaga | agatgcagac | tcgccccag | 1260 |
| agcggggcca | gcggcccaaa | gagggcgctt | actatgtgtg | gacgggtcaa | gaggttcagc | 1320 |
| agctcctccc | ggagcctgtg | ttgggtgcca | ccgagccgct | gacctcaggc | cagctcctca | 1380 |
| tgaagcacta | cggcctcaca | gaggctggta | acatcagccc | cagtcaggac | ccaagggggg | 1440 |
| agctgcaggg | ccagaatgtg | ctgaccgtcc | ggctactcgt | ggagctgact | gctgcccgct | 1500 |
| ttggcttgga | tgtggaggcc | gtgcggacct | tgctcaattc | agggctggag | aagctcttcc | 1560 |
| aggcccgga | gcctcggccc | aagccgcacc | tggacagcaa | gatgctggct | gcctggaatg | 1620 |
| gcttgatgg | gtcaggctat | gctgtgactg | gggctgtcct | gggccaagac | aggctgatca | 1680 |
| actatgccac | caatgggtgc | aagttcctga | agcggcacat | gtttgatgtg | gccagtgggc | 1740 |
| gcctgatgcg | gacctgtac | accggccctg | gggggactgt | ggagcacagc | aaccacacct | 1800 |
| gctggggctt | cctggaggac | tacgccttcg | tggtgcgggg | cctgctggac | ctgtatgagg | 1860 |

| | | | | | | |
|------------|------------|------------|------------|------------|------------|------|
| cctcacagga | gagtgcgtgg | ctcgagtggg | ctctgcggct | gcaggacaca | caggacaggg | 1920 |
| tcttttgga | ctcccagggt | ggcggctact | tctgcagtga | ggctgagctg | ggggctggcc | 1980 |
| tgcccctgcg | tctgaaggac | gaccaggatg | gagcagagcc | cagcgccaat | tccgtgtcag | 2040 |
| cccacaacct | gctccggctg | catggcttca | cgggccacaa | ggactggatg | gacaagtgtg | 2100 |
| tgtgcctatt | gaccgccttt | tccgagcgca | tgcgtcgtgt | cccgggtggc | ttgcccgaga | 2160 |
| tggtcgcgc | cctctcagcc | cagcagcaga | ccctcaagca | gatcgtgatc | tgtggagacc | 2220 |
| gtcaggccaa | ggacaccaag | gccctgggtg | agtgcgtcca | ctctgtctac | attcctaaca | 2280 |
| aggtgctgat | tctggctgat | ggggaccctt | cgagcttcc | gtcccgcag | ctgcctttcc | 2340 |
| tgagtaccct | ccgacggttg | gaagaccagg | ccactgcata | tgtgtgtgag | aatcaagcct | 2400 |
| gctcagtgcc | catcactgat | ccctgcgaat | tacgaaaact | actacatcca | tgactgcccc | 2460 |
| aacccccctt | gggtggggca | gaaggtgaag | catcccaact | gactagagac | tcaggccctg | 2520 |
| cagggcccta | tagaacctgt | ggccatccct | gagcaccctg | ccaccaggtg | acctcgcca | 2580 |
| tactcactgc | cccccttggg | caccactca | ccctagaata | aacttaacag | tgtcccgtgg | 2640 |
| taaaaaaaaa | aaaaaaaaaa | aaaaaaaaag | ggcggccgc | | | 2679 |

<210> 24

<211> 1786

<212> DNA

<213> Homo sapiens

<400> 24

| | | | | | | |
|-------------|------------|-------------|------------|-------------|------------|------|
| tgcaggttca | gatgtgtata | agagacaggg | tctatataag | cagagctcgt | ttagtgaacc | 60 |
| gtcagatcgc | ctggagacgc | catccacgct | gttttgacct | ccatagaaga | caccgggacc | 120 |
| gatccagcct | ccggactcta | gcmtaggcgc | cgggacggat | aacaatttca | cacaggaaac | 180 |
| agctatgacc | actaggcttt | tgcaaaaagc | tatttaggtg | acactataga | aggtacgcct | 240 |
| gcaggtagcg | gtccggaatt | cccgggtcga | cccacgcgtc | cgatgaagcc | ggccactgcc | 300 |
| tctgctctgc | tctgctcct | gctgggcctg | gcctggaccc | aggggagcca | cggctggggt | 360 |
| gcggacgcgt | catcactgca | gaaacgtgca | ggcagagccg | atcaggtgag | tctctgtcct | 420 |
| caagttaccc | tgcaggggcc | ttgggtcccc | ttggccttgc | tcccgggtct | gggcaatctg | 480 |
| aaattctctt | tactcctcc | attcaatggc | tttctcagcc | gggtgcagga | tggcaggagg | 540 |
| tggcagctgt | aacttccaag | gtgagacatc | taccatggat | ccgctgggtt | acgtcctggc | 600 |
| acttcttttc | ttggtctcct | accctcacct | cccaagtaac | acttgaagag | ttggcagagg | 660 |
| tcttgagggc | aggaggagga | gtaggaggag | gaggcgccgc | ccatttgggc | taggacagta | 720 |
| aaggcagggtg | aggggtgggg | ccctggctgg | ctggaggaga | ttgtataaaa | gaaaacacgt | 780 |
| gacacaagga | ggggagcagg | cagggtgtttt | aggaaaggct | tgggtccaggc | aggaagggtg | 840 |
| gatgttttatt | ccagagcaaa | atggccttcg | tccgtctgtt | ctaaccgcca | tgcccacttt | 900 |
| cttccacaga | actacaatta | caaccagcat | gcgtatccca | ctgcctatgg | tgggaagtac | 960 |
| tcagtcaaga | cccctgcaaa | ggtgagaccg | gcaaccaccc | tgtcttcccg | ctgggggctc | 1020 |
| tgcagtgggg | atggggagag | aggctttcac | tgccgggggc | ctctcggttg | gttgccggag | 1080 |
| gccggatccc | gctctctttg | agtccaggcc | tccccacccc | tttgcaaggg | agactcgggt | 1140 |
| tccctgagac | agtttggaag | tggagggcag | gagggttcaa | tgcgcctcac | acccaccag | 1200 |
| tcagaaaagg | ttaaaaagga | gcagcagcag | caacagagaa | aactgagaac | caacaccctc | 1260 |
| tccaagaaga | atataggctt | taaaaaatca | tgacacctgc | aaaacacgca | agcagctctt | 1320 |
| cctcacaggt | tccctctctg | tggacctttt | tttgcggggg | aagatggaat | ggagggcgag | 1380 |
| gaaagcagac | tagcacttag | gggaggccgg | cctgtggatg | cctggttggc | ggtgaaatga | 1440 |
| atgcttttatt | ccacaaagga | aatgaaaaga | aaaaatacgt | gagaatttgg | ttttattccg | 1500 |
| ttgcaaagaa | ggcaatctca | taacatcagt | aatgaatat | gctaagtaca | aatttggtcg | 1560 |
| acaaagcaac | agcaggctgt | gtatttccat | ctctcatgac | aataaagtcc | ccgcccttgg | 1620 |
| ggagaggacc | ttccactgag | ggacgagaaa | ctacaggctg | gtgagaggag | ggacgtgcag | 1680 |
| cttctactgc | ctgtggcttc | cactctgctg | agttatctgt | ggcttccact | ctgccggggt | 1740 |
| atctgtggtc | ctgtctctta | tacacatctc | aacctgcagg | gggggc | | 1786 |

<210> 25

<211> 3431

<212> DNA

<213> Homo sapiens

<400> 25

| | | | | | | |
|-------------|-------------|-------------|-------------|-------------|-------------|------|
| gcgtccgaaa | ttaataagct | cctaacataa | atttgatgtc | atttgcagat | gagaaaaatgg | 60 |
| ggcttgatga | agttgattgc | ttcaatgatg | caaccagtc | tgcttgaact | cctctctgtc | 120 |
| tggaggaagg | aaggcagaga | ctctagaaac | atacatgatt | cccacagcat | gtatgtgctg | 180 |
| aggaagagac | tgagtgggtc | ttggcttcag | gcaggctctgt | actctactgt | gatatcagca | 240 |
| gcactcatcc | ttgaatctcc | aagagcttgt | cttccttcaa | aaggctgatt | cagacacaga | 300 |
| gtctagactt | gtttctcttt | tcttaaaagt | acacatcagg | aagaaaacct | aaggcctctt | 360 |
| gtaaaaacat | cactgacatg | cttcctgggt | atttctcagc | agctatattgt | ccccagata | 420 |
| tattttttca | ctgccatttc | acccaatggt | aattgaaatg | ctaaactaaa | tgtttagcca | 480 |
| tccttgcatg | catggataat | taataaactg | attataaaat | tatagtgatg | agggattact | 540 |
| aagccatgat | aaattacaat | cattatttca | gtcttgacta | taaagattat | tataaggatg | 600 |
| atagaggttt | cttatggatt | tttagtcatt | tacacataat | gaagtaaaaa | tgtctttttc | 660 |
| aattttcaaa | tatgtcaccg | aagccaaaat | tggtatctgt | catttttaag | ttttgtatat | 720 |
| gcttgatttc | tgttgtattt | gtttcagatt | gtgtttgctt | catgaaaagg | atgttttctt | 780 |
| ttttcaggta | gtgaggggtt | taactacatt | ttatttctcc | caaaaagctta | ttaatcttgg | 840 |
| caataaaaaat | tctatatattc | tccagcaaaa | tctaagatgc | ttacctcaaa | gcagcttttg | 900 |
| aagtattttc | tatcagtatt | gttctgatta | tgctgtgagg | agcctgagga | attcctcaga | 960 |
| gacacctgag | gacaggggca | aggaaaaata | tcagagttaga | ctgggatggg | ggaggatgga | 1020 |
| gatggctctt | gcagtcgtga | ggaactcacc | tctggttgag | ttgagcagtt | ccacttttga | 1080 |
| tctgtgttgc | atgttggact | tttacataag | gttttatattg | aagaaagatc | catgtggtac | 1140 |
| catgaatatt | aataatatta | actaataata | atattaacta | ataatcaaaa | ggacttgaaa | 1200 |
| tccaatgatt | aaatacctaa | attttataaa | tgagggaact | gagcttcaga | gagagtaaat | 1260 |
| aacttgttta | acataacaca | gctagtaaaa | atttgcctga | attaattatt | cctaaaaatt | 1320 |
| aaaaggctat | gatcaactta | accaatattt | atcaaagtc | tacaaaggaa | aaggcactat | 1380 |
| gccaggcctc | cagggaataa | agaaaaatac | aacgtcacac | ttcaagtaaa | tggcagaatc | 1440 |
| aaggcatgaa | tctgaatctt | cttaccocaa | ctagagtga | ctttccacta | tagcagtag | 1500 |
| ccccgaattt | ggtttatgaa | cacctagtct | ttcaggatga | tcttccactt | gaaaaattca | 1560 |
| attgaagttt | tgcatatgaa | tttaaaaggt | atgagaagtc | ctatagtaaa | gaaatgtgct | 1620 |
| tagactacag | ttccaaaatt | tattaccatg | gaatactatt | catgtcatat | gtatttatat | 1680 |
| cccaaagaat | gcaatttgag | aaattctgca | ctatcatagc | tctctttatt | agatttgaaa | 1740 |
| ctgaaatcta | gcagagtctt | cttttaaaac | ctacatttcc | atgggtttaca | tgtaactgtg | 1800 |
| gcttgagaaa | tggttaatctc | agtgataagt | gttagactgg | atatactctt | tatatacaca | 1860 |
| tacacatatt | gtatgtggat | gttacagaaa | atactatgtc | caaataagca | taaattaaat | 1920 |
| gcagtatacc | aactgttaac | acagtattgt | gtttgcttta | aagttggctc | acacctgttt | 1980 |
| aatatttgct | atgaaaatct | caatgcagtt | tttattcctt | ctaaacattt | aatgccagca | 2040 |
| ttataattca | ctggaaatgt | tgaatttaatt | ttagtataaat | aaggactgat | taatacctct | 2100 |
| agtccaattt | tcagataaat | gactttctgc | aaaaggaaaa | tcaatttggt | tttcttataa | 2160 |
| aacacacaca | ccttgctgca | ttttgttttg | gtcagaatgt | taaatataaa | aaataataaa | 2220 |
| tagcaaaatt | ttccccaaaa | attcatttca | ggtacaccga | acaaaaataa | tgccataaac | 2280 |
| cttggaagag | aacctgttct | cgcaagacca | atttttcagg | atcaaaaagt | tttatataag | 2340 |
| ctcacaaaatt | ttggatcttt | ttcaagacta | aatctccaac | acttctgagg | ttggcccatc | 2400 |
| actaatattt | acaatatatt | tgacatttca | tgggatggac | tcacaggcct | ttatgatagg | 2460 |
| cagcttccta | atatgacaga | gtgccagaac | aatttgctgt | gtgtgcgaaa | ggcagtaaga | 2520 |
| tggcactggc | attttgcaac | aggaaggaaa | tgatctgtag | gtaccttgag | cctgttgtat | 2580 |
| cttgtcagtt | ttgttagttg | agaaatgtca | ttggggaccc | attagtacat | aggcagccgt | 2640 |
| gacgccacc | tgagaagcac | tccttgccaa | gtgatttgag | acctgtgttt | gggcaagggt | 2700 |
| ccttttcata | gtattacatt | gcattacaga | aaacttacag | aaacaggccc | tcacgcatgc | 2760 |
| tgattcttaa | ccctttataa | gacctagcat | gatgccttac | atttatttag | ccattataat | 2820 |
| aagatttttc | aacagggtca | tgcccttacc | aattgggtaca | tgtaaaaataa | agtgtattct | 2880 |
| gaactctttt | atcacacatc | cctctaatac | cccatttatc | actgtgttta | attttatttt | 2940 |
| tttccaaagt | ctggctcattc | ctaacaatgc | tcttgatgac | tgcataatta | ttacactttc | 3000 |
| attgccaaag | acagaaatag | ggtaaaagca | cagatgcttt | ttctcctgcc | aaattaggat | 3060 |
| tatctccaa | aatctgtatt | attacttggc | ctcttttaag | agaccatttt | ctttggattg | 3120 |
| tgagggttca | cacctgtaat | atgcatgctc | aaggttctgt | aatcttgaaa | gttgttttgt | 3180 |
| gagatcaagc | ataggaaact | ctattaggta | tgacaaagat | gggtaagata | tggatcctgc | 3240 |
| tctcggagag | cttcaatcta | attgtagatt | caagacaaac | acatatatcat | ggggcaccac | 3300 |
| ttaccaaaacc | ccaatgaaag | cactattgta | gaagtggttc | agatgcttgt | attgacccaa | 3360 |

aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 3420
 aaaaaaaaaa a 3431

<210> 26
 <211> 766
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (670)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (713)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (721)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (728)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (731)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (756)
 <223> n equals a,t,g, or c

<400> 26
 ggcgagacgc gttcaccacc atatgcactg ttgtcaactc ccctggagat gcgcccgaagc 60
 cccacaggaa gccttcctcc tctgcctcct cttcctcctc ctcgtcctcg ttctccttgg 120
 atgcagccgg ggccctccctg gccacactcc ctggaggctc cctcctgcag ccgcggccct 180
 ccttgccctc ctctcccaag atgcacttgg ggctgtgtgt ttccaaggcc ctgagtacct 240
 cttgccttgt ttgctgcctc tgccaaaacc cggccaactt caaggacctt ggggacctct 300
 gtggggcccta ctaccctgaa cactgcctcc ccaaaaagaa gccaaaactc aaggagaagg 360
 tgcggccaga aggcacctgt gaggaggcct cgctgccgct tgagagaaca ctcaaaggctc 420
 ccgagtgtgc agctgccgcc actgccggga agccccccag gcctgacggc ccagctgacc 480
 cggccaagca gggcccactg cgcaccagtg cccggggcct gtcccggagg ctgcagagct 540
 gctactgctg tgatggccgg gaggatgggg gcgaggaggc agccccagcc gacaagggtc 600
 gcaaacatga gtgcagcaag gaggtccgg cagagcccgg cggggaggcc caggaagcac 660
 tgggtgmatn aagcctgttm ccgtgttgga ccgggggggt tttaactggt ggnccggggaa 720
 ntcttttngg nttgcaagag gccaatataa gttggnccct ggaaaa 766

<210> 27
 <211> 6729
 <212> DNA

<213> Homo sapiens

<400> 27

| | | | | | | |
|-------------|-------------|------------|-------------|-------------|-------------|------|
| ccacgcgtcc | gggagggcaa | cggggcggcg | cggggcggcg | ggcgcagggt | cgcgaggagt | 60 |
| gacgcgcggc | gaggatggcg | gcgcggggcc | gggggtgctg | gctgctgacg | ctgtcggtgc | 120 |
| tggtggcggc | ggggccctcc | gccgctgcgg | ccaagctcaa | catcccaaaa | gtgctgctgc | 180 |
| ccttcacgcg | ggccacgcgc | gttaacttca | cgctggaggc | ctcgaggggc | tgctaccgct | 240 |
| ggttgtccac | ccggccggag | gtggccagca | tcgagccgct | gggcctggac | gagcagcagt | 300 |
| gctcccagaa | ggcagtgggtg | caggcccgcc | tgaccagcc | tgcccgcctc | accagcatca | 360 |
| tcttcgcaga | ggacatcacc | acaggccagg | tctgctgctg | tgatgccatt | gtggacctca | 420 |
| tccatgacat | ccagatcgtc | tccaccaccc | gcgagctcta | cctggaggac | tccccctgg | 480 |
| agctgaagat | ccaggccctg | gactccgaag | ggaacacctt | cagcactctg | gctggactgg | 540 |
| tcttcgagtg | gacgattgtg | aaggactccg | aggcggacag | gttctcagac | tcccacaatg | 600 |
| cgctgcgaat | cctcactttc | ttggagtcta | cgtaacatccc | tcttctttac | atctcagaga | 660 |
| tggaagaaggc | tgccaagcaa | ggggacacca | tcttggtgtc | tgggatgaag | accgggagct | 720 |
| ccaagctcaa | ggctcgatc | caggaggctg | tctacaagaa | tgtacgccct | gcagaagtca | 780 |
| ggctgctgat | tttgaaaaac | atccttctga | acccggccta | tgacgtctac | ctgatggtgg | 840 |
| gaacctccat | tactacaag | gtgcagaaga | tcaggcaagg | gaaaattaca | gaactctcca | 900 |
| tgcttccga | tcagtacgag | ttgcagcttc | agaacagcat | cccgggcccc | gaaggagacc | 960 |
| caacccggcg | ggtggctgtc | ttggcccagg | acacgtcgat | ggtcactgca | ctgcagctgg | 1020 |
| gacagagcac | cctcgctcctt | ggccacagga | gtattcgcat | gcaagggtgct | tctaggttac | 1080 |
| ccaacagcac | tatctacgtg | gtcgaacctg | gatacctagg | gttactgttt | cacctgggtg | 1140 |
| acagggtgggt | gctggagacc | ggccgcctgt | atgaaatcac | catcgaagtt | tttgacaagt | 1200 |
| tcagcaacaa | ggtctatgta | tctgacaaca | tccgaattga | aactgtgctt | cctgctgagt | 1260 |
| tcttcgaggt | gctctcgtcc | tcccagaatg | ggtcatacca | tcgcatcagg | gcactaaaga | 1320 |
| ggggacagac | ggccattgac | gcggccctca | cctctgtggt | ggaccaggat | ggagggggtcc | 1380 |
| acatactaca | ggtgcctgtg | tggaaccagc | aggagggtga | aattcacatc | ccgatcaccc | 1440 |
| tgtatcccag | catcttgaca | tttccgtggc | aaccaaaagc | gggcgcctat | cagtacacaa | 1500 |
| taaggggcca | cggtggcagt | gggaacttca | gctggtcttc | gtcaagccac | ctggttgcca | 1560 |
| cagttactgt | caagggcgtg | atgaccacag | gcagtgcacat | cgggttcagt | gtgatccagg | 1620 |
| cacatgatgt | gcagaaccca | ctccatttcg | gtgagatgaa | ggtgtatgtg | atcgagcccc | 1680 |
| acagcatgga | gtttgccccg | tgccagggtg | aggcagctgt | gggccaggcc | ctggagctgc | 1740 |
| ccctgaggat | cagtggcctc | atgcccgggc | gggccaagtga | ggtggtcacc | ttgagcgact | 1800 |
| gctcccactt | tgacttggtc | gtcgagggtg | agaaccaggg | tgtgttccag | ccactcccag | 1860 |
| ggaggctgcc | gccaggctct | gagcactgca | gcggcgctcg | ggtaaaggcc | gaggcccagg | 1920 |
| gctctaccac | gcttcttgtg | agctacagac | acggccacgt | ccacctgagt | gccaaagatca | 1980 |
| ccattgtctg | ctacctgcc | ctcaaggctg | tggatccctc | ctctgttgcc | ttggtaaccc | 2040 |
| tgggtctcctc | aaaggagatg | ctgtttgaag | gagggtcccag | accttggtac | ctcgagccgt | 2100 |
| ccaaattctt | ccagaacgtc | accgctgagg | acactgacag | catcggcctg | gctctctttg | 2160 |
| ccccccattc | ctcccgaat | tatcagcaac | actggatcct | tgtgacctgt | caggccttgg | 2220 |
| gtgagcaggt | catcgccctg | tcggtgggga | acaagcccag | cctcaccaac | ccctttcctg | 2280 |
| cggtggagcc | tgccgtgggtg | aagtctgtct | gcgccccacc | gtccaggctc | accctcgtgc | 2340 |
| ctgtctacac | cagccccag | ctggacatgt | cctgtccgct | gctgcagcag | aacaagcagg | 2400 |
| tggtcccagt | gtccagccac | cgcaaccccc | tgctggacct | ggctgcttac | gaccaggagg | 2460 |
| gccgcgggtt | cgacaacttc | agctctctga | gcatccagtg | ggagtccacc | aggccagtgt | 2520 |
| tggccagcat | cgagcctgag | ctgcccattg | agctggtgtc | ccaggacgat | gagagtggcc | 2580 |
| aaaagaagct | gcacggtttg | caggccattt | tggttcacga | ggcatcagga | accacagcca | 2640 |
| tactgccac | tgccactggc | taccaggagt | cccacctcag | ctctgccaga | acaaagcagc | 2700 |
| cgcattgacct | tctggtgcct | ctgtcggcct | ccatagagct | catcctggtg | gaggacgtga | 2760 |
| gggtgagccc | agaagagggtg | accatctaca | accaccctgg | catccaggca | gagctccgca | 2820 |
| tcagggaagg | ctcaggttac | ttcttctcca | acaccagcac | cgcagatgtt | gtcaagggtg | 2880 |
| cctaccagga | ggccagggggt | gtcgccatgg | tgcacctttt | gctcccgggc | tcatccacca | 2940 |
| tcgatattcca | tgacttgtgc | ctcgtcttcc | cggccccagc | caaggctgtc | gtttacgtgt | 3000 |
| cggacattca | ggagctgtac | atccgtgtgg | ttgacaagggt | ggagattggg | aagacagtga | 3060 |
| aggcatacgt | ccgcgtgctg | gacttgcaca | agaagccctt | ccttgccaaa | tacttcccct | 3120 |
| tcatggacct | gaagctccga | gcagcctccc | cgatcattac | attggtggcc | cttgatgaag | 3180 |
| cccttgacaa | ctacaccatc | acattcctca | tccgcgggtg | ggccatcggc | cagaccagtc | 3240 |
| taactgcaag | tgtgaccaat | aaagctggac | agagaatcaa | ctcagcccca | caacagattg | 3300 |

| | | | | | | |
|-------------|-------------|------------|------------|------------|------------|------|
| aagtcttttc | cccgttcagg | ctgatgccca | ggaaggtgac | actgcttata | ggggccacga | 3360 |
| tgcaggtcac | ctccgagggc | ggccccacgc | ctcagttcaa | catccttttc | tccatcagca | 3420 |
| atgagagcgt | tgcgctggtg | agcgtgctg | ggctggtaca | gggcctcgcc | atcggaacg | 3480 |
| gcactgtgtc | tgggctcgtg | caggcagtg | atgcagagac | cggcaagggt | gtcatcatct | 3540 |
| ctcaggacct | cgtgcagggt | gaagtgtgtc | tgctaagggt | cgtgaggatc | cgcccccca | 3600 |
| tcatgaggat | gaggacgggc | acccagatgc | ccatctatgt | caccggcatc | accaaccacc | 3660 |
| agaacccttt | ctccttttgc | aatgccgtgc | caggcctgac | cttccactgg | tctgtcacca | 3720 |
| agcgggacgt | cctggacctc | cgagggcggc | accacgaggc | gtcgatccga | ctcccgtcac | 3780 |
| agtacaactt | tgccatgaac | gtgctcggcc | gggtaaaagg | cgggaccggg | ctgagggttg | 3840 |
| tgggtcaaggc | tgtggacccc | acatcggggc | agctgtatgg | cctggccaga | gaactctcgg | 3900 |
| atgagatcca | agtccaggtg | tttgagaagc | tgcagctgct | caaccctgaa | atagaagcag | 3960 |
| aacaaatatt | aatgtcgccc | aactcatata | taaagctgca | gacaaacagg | gatggtgcag | 4020 |
| cctctctgag | ctaccgctgc | ctggatggac | ccgaaaagg | tccagttgtg | catgttgatg | 4080 |
| agaaaggctt | tctagcatca | gggtctatga | tcgggacatc | caccatcgga | gtgattgcac | 4140 |
| aagagccctt | tggggccaac | caaaccatca | ttgttgctgt | aaaggatctc | cctgtttcct | 4200 |
| acctgagggg | ttccatgagc | cctgtcctgc | acaccagaa | caaggaggcc | ctggtggccg | 4260 |
| tgcttttggg | aatgaccgtg | accttcaact | tccacttcca | cgacaactct | ggagatgtct | 4320 |
| tccatgtctc | cagttcggtc | ctcaactttg | ccactaacag | agacgacttt | gtgcagatcg | 4380 |
| ggaagggccc | caccaacaac | acctgtgttg | tccgcacagt | cagcgtgggc | ctgacactgc | 4440 |
| tccgtgtgtg | ggacgcagag | caccggggcc | tctcggactt | catgcccctg | cctgtectac | 4500 |
| aggccatctc | cccagagctg | tctggggcca | tgggtggtgg | ggacgtgctc | tgtctggcca | 4560 |
| ctgttctgac | cagcctggaa | ggcctctcag | gaacctggag | ctcctcggcc | aacagcatcc | 4620 |
| tccacatcga | ccccaaagac | ggtgtggctg | tggcccgggc | cgtgggatcc | gtgacggttt | 4680 |
| actatgaggt | cgctgggcac | ctgaggacct | acaaggaggt | ggtggtcagc | gtccctcaga | 4740 |
| ggatcatggc | ccgtcacctc | caccccatcc | agaccagctt | ccaggaggct | acagctcca | 4800 |
| aagtgtattgt | tgccgtggga | gacagaagct | ctaacttgag | aggcgagtgc | acccccacc | 4860 |
| agagggaaagt | catccaggcc | ttgcaaccag | agaccctcat | cagctgccag | tcccagttca | 4920 |
| agccggccgt | ctttgatttc | ccatctcaag | atgtgttcac | cgtggagcca | cagtttgaca | 4980 |
| ctgctctcgg | ccagtacttc | tgtcfaatca | caatgcacag | gctgacggac | aagcagcgga | 5040 |
| agcacctgag | catgaagaag | acagctctgg | tggctcagtc | ctccctctcc | agcagccact | 5100 |
| tctccacaga | gcagggtggg | gccgaggtgc | ccttcagccc | aggtctcttc | gccgaccagg | 5160 |
| ctgaaatcct | tttgagcaac | cactacacca | gttccgagat | cagggtcttt | ggtgccccgg | 5220 |
| aggttctgga | gaacttgag | gtgaaatccg | ggtccccggc | cgtgctggca | ttcgaaagg | 5280 |
| agaagtcttt | tgggtggccc | agcttcatca | catacacggt | cggcgtcttg | gacccccgg | 5340 |
| ctggcagcca | agggcctctg | tccactaccc | tgaccttctc | cagccccgtg | accaaccaag | 5400 |
| ccattgccat | cccagtgaca | gtggcttttg | tgggtggatc | cgtggggccc | ggtccttatg | 5460 |
| gagccagcct | cttccagcac | ttcctggatt | cctaccaggt | catgttcttc | acgtctctcg | 5520 |
| ccctgttggc | tgggacagcg | gtcatgatca | tagcctacca | cactgtctgc | acgccccggg | 5580 |
| atcttgctgt | gcctgcagcc | ctcacgcctc | gagccagccc | tggacacagc | ccccactatt | 5640 |
| tcgtgcctc | atcacccaca | tctcccaatg | cattgcctcc | tgtctgcaaa | gccagccctc | 5700 |
| cctcagggct | gtggagccca | gcctatgcct | cccactaggc | cgcgtgaagg | ttccccgagg | 5760 |
| atgggtctca | gccgagcctc | gtgcaccccc | aagatggaac | atccctgctg | cattcacact | 5820 |
| ggaacaagcc | cctccagatg | agtgcctcgg | ccccaggcca | gcttcaactg | cgtctcttca | 5880 |
| cacagagctg | tagtttcggc | tctgcccatt | agctcatttt | atgtaggagt | tttaaagtgt | 5940 |
| tgtttttttc | ctttcaagtc | ttacaaagct | aagacttttt | ggctcattcc | tttttgcatg | 6000 |
| gttgtctagg | gtttctggac | aatgtgctgt | tgcattttta | ttttcctagc | cttgctaaaa | 6060 |
| tctttccctt | ctcaagactt | tgagcagtta | gaagtgtctc | ttagaagttg | tctgtgggtg | 6120 |
| atgttactgt | agtggctctc | gggaaaggat | tgtccagtta | ctttaggggg | tttttggttg | 6180 |
| ggtttttccc | cctgtgaaaa | cttactttgc | ccctagtctg | gctgctgcta | ggacttctga | 6240 |
| ggagcaatgg | gacatgagt | tccctgtatc | tgcgccactg | cgcgaaggga | agcctcagga | 6300 |
| accagcacct | ggaggccagg | gctgctgccc | aacctggggc | cactgtgaac | agacttcagt | 6360 |
| cctctgtttt | tgtttcataa | gccgttgaga | catctgatgg | acttggttta | ggccctgctg | 6420 |
| ggacatccca | cgtgtgatcc | ctttcactcc | atcaggacac | caggactgtc | cttaggaaaa | 6480 |
| tgtccttgag | atggcagcag | gagtcataat | ttctgtgtgt | gtgtttcgga | aagccgctgt | 6540 |
| gtcctgcctc | agcacaaaaga | cccagtgtea | tttgctcctc | ctgttccctg | gccactccag | 6600 |
| aacctcagca | gatctgagcc | accgcctgcc | agtgtgagag | gcggccactt | tcatggcagc | 6660 |
| tcatcaggcg | cagggcccca | gacagcttcc | cagcaggccc | tagagcccgg | cctggggcaa | 6720 |
| tgatggagg | | | | | | 6729 |

<210> 28
 <211> 1242
 <212> DNA
 <213> Homo sapiens

<400> 28
 gcgtccgcac ctcaggccct ccaagcgcag gatgcaggcc gtggccaacg tgtccattgg 60
 ggccatgttc tgcattgtatg ggctcacagc aaccttttga tacctcacct tctacagtga 120
 gtggggcttg ggctagggct ggggggaggg ggaaggcctg gggcaggagc ctctgagctc 180
 tttccttctg tgaccacgga cctgtcaagt ttccaaacag aaagggtgtg ctcactttgtg 240
 tggattttgt cacttgtgca tgtatgtatg ggtttctggg gcattgggtc tgggtgctctc 300
 tccacatcct gcatcccgtat cctctgtct catggcccag gcagtgtgaa ggcggagatg 360
 ctgcacatgt acagccagaa ggaccgcctc atcctctgtg tgcgcctggc cgtgctgctc 420
 gcggtgacct tcaactgtgc agtcgtgctg ttccctatcc gccgggcccct gcagcagctg 480
 cttttccag gcaaggcctt cagctggcca cgacatgtgg ccatagctct gatcctgctt 540
 gttttggtca atgtccttgt catctgtgtg ccaaccatcc gggatatctt tggagttatc 600
 ggggtccacct cagccccag cctcatcttc atcctcccca gcattcttcta cctccgcat 660
 gtacctctg aggtggagcc tttcttatcc tggccaaga tccaggcccct gtgctttgga 720
 gtcctgggag tcctcttcat ggccgtcagt ctaggcttta tgtttgccaa ctgggccaca 780
 ggccagagcc gcatgtctgg acactgatca ggcctgctg gcccaggctc ctgtgcgcat 840
 gcacatggag gggtcagggc cgtccctag ggtccctcct gcccaacatg tggaggtggc 900
 tggttcccat gaacgtggtt gtcagaggcg ggggacaagc agaggcttgc agactggccc 960
 acttccctcc tccccaaggg atgccaaagct tggatcatgg ccctaatacc aaccccaacc 1020
 ccatgggag agggaggagga ggaagaagag gaggaggagg agggaggagga ggcggaggag 1080
 gaggaggcca ggtcctgggt gagcctttgc ccagcccagt cctctctgcc tctcctggc 1140
 tgaagctgtt tgtcaggatt accctcgggc taaagaggaa aaataaagat gttgagctac 1200
 caaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aa 1242

<210> 29
 <211> 2032
 <212> DNA
 <213> Homo sapiens

<400> 29
 ggcacgagag aagccatacc tgggtgaaga ggccggtttct tacaatgaac ttgactatgt 60
 gtcggttggc cttgaccagc agacggtgaa gctgggtgtgc accaaccgca ggaagcagtt 120
 tctgctggac acggtgatg tggcgctggc tgagttcttt ttggcttctt tgaagtcagc 180
 catgatcaaa ggctgtcgag aacctcccta cccagcatc ctgacggatg ccacatgga 240
 gaagctggca ctggccaaat ttgtggccca agaatcgaag tgtgaggcat ctgctgtcac 300
 cgtgcgcttc tacggccttg tgcactggga ggacccaca gacgagtcct tgggcccac 360
 gccctgccac tgctcaccac ccgagggcac catcacaaa gaaggcatgc tgcactaaa 420
 ggcgggcacc tcctacctgg gcaaggaaaca ctggaagacg tgcttcgtgg tgctcagcaa 480
 cgggatccct taccagtacc cggaccgcac cgacgtcatc cctctgctct cggatgaacat 540
 ggggggggag cagtgcggtg gctgccggag agccaacacc acggatcggc cccacgcctt 600
 ccaggtcatt ctctccgacc ggccctgcct ggagctaagt gccgagagcg aggcgagat 660
 ggccgagtg atgcagcatc tctgccaggc tgtgtccaaa ggggtcatcc cccagggcgt 720
 agtcccagc cctgtcatac cctgtgcct ggtcctcacg gatgaccgcc tctttacgtg 780
 ccatgaggat tgccagacca gcttcttccg ctctttgggc acagccaagc tggcgacat 840
 cagcgccgtc tccaccgagc cgggcaagga gtactgcgtc ttggagttct cccaggacag 900
 ccagcagctc ctcccgccct gggtcattcta cctagctgc acttctgaac tggaccgatt 960
 gctgtctgca ctgaactctg ggtgaaaaac catctatcag gtggacctcc cccacacggc 1020
 gatccaggaa gcctccaaca agaagaaatt cgaggatgcc ttgagcctca tccacagcgc 1080
 ctggcagcgg agcgacagtc tctgccgcgg ccgagcctcc cgagaccctc ggtgctgagg 1140
 cagagctggg tggcgctccct ggtgggcagg aaagggaagg acgccagccg gcaggcacac 1200
 tgtcacggct gttgtcatgc tgtcgggagc ctacagtcca cccctgcctt ggcggcaga 1260

| | | | | | | |
|------------|-------------|-------------|-------------|------------|-------------|------|
| accaccgagt | gtggcttaag | acaggggtccc | tccactccag | ggatccagat | caggtgcccg | 1320 |
| gcacccctgg | gcacctctgcc | cgacaggtag | cgaatggagg | tcgctggggg | cagaggggtcc | 1380 |
| gagccctgtg | ggctctgcgg | atgcacgccc | tctcccggg | cctccgcctc | agtctgcaga | 1440 |
| atctctgccc | agtggcaccg | agaacacccat | ccatctaagg | acgaacaaaa | gaaccaggag | 1500 |
| ggcgggaccc | cccctcttcc | tctcctgggt | tgggggctgg | ggccctgagt | gcccagccat | 1560 |
| ccttgttcgt | gtttgaacac | tctcctggcc | acgtggggaa | gcgggaacac | gggggtgtctg | 1620 |
| cgcattgttc | ctcctcctag | ctccatcact | gcgcacacag | ctgcctgcct | cgccagatgc | 1680 |
| agggggcgcg | gcagccctcc | ctggctgcca | ggaggctctg | catgcccaca | gtcctgcccct | 1740 |
| gcctgtcccc | tcaaccggc | agtgcctgta | gcaccgagga | gcaaaggggg | tggatggggg | 1800 |
| gcttgagaaa | gggcggagcc | caccagcctg | gcattccatgt | tgacatcttc | tgactgtccc | 1860 |
| ctgcttggtc | ggagccaggc | ccttccctag | agtttcgtca | agagcctcct | ggggaagggg | 1920 |
| tcaggtggtt | tgggttttgt | tttttaaaat | aaaatagaca | tgttatatgt | caaaaaaaaa | 1980 |
| aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | aa | 2032 |

<210> 30

<211> 3408

<212> DNA

<213> Homo sapiens

<400> 30

| | | | | | | |
|-------------|-------------|-------------|------------|-------------|-------------|------|
| ccacgcgtcc | ggaggcagga | ccttgtccta | ttcattaatc | ttgcccctca | acagttatatt | 60 |
| tcagaggggc | aagaagtgtt | tcaggggtct | tggcccttgt | ttgaccagtc | gtcctaacc | 120 |
| tcattgtctg | ggtcattgtt | gttataatct | ggggttacct | tttggaaagt | catgggggtac | 180 |
| ccttttgcaa | aagtatatgg | ccctctcctt | ggaaactgca | cacacaccat | gcagcttaca | 240 |
| attcaggag | ttcacaggtc | tacagaatcc | tggaaactct | catgtccggg | tctactcatt | 300 |
| gtagcttcag | tggaaacctt | tagcagtcct | ttccagctcc | tccccagctc | ctcagctctg | 360 |
| cttccctccg | cccatcaagc | cctcctcagc | ccataagggt | ggccaggtgg | cctgtgggga | 420 |
| taaatcagag | tgcccaaca | tgacagggcc | caaagacatc | ccagagcaaa | cccaagaatc | 480 |
| cctctacaag | ccccagccca | ctcagaagga | tgcattttgc | cccctctgtt | tatttgtttg | 540 |
| tttttaatta | tgaagtagg | gcattggtcat | tggtgagaat | gtgagaaatg | cagagaagtt | 600 |
| aaaatgatgc | tttttgttta | gggcgtgct | gcttctggct | taagatccta | aatcaaaagca | 660 |
| gctgccagat | ctggacctaa | gacttgcttc | ccatcacctt | acataaaaaga | aagagcactg | 720 |
| gactaggaat | caaagatctg | aattcccac | aaatctctgc | tattactagc | ttttatctct | 780 |
| atttatattat | ttatctgtct | atctatctag | taccttttgt | gaatatgagt | gtttctcacc | 840 |
| gaggccctcc | catctctctt | gcccccgatc | ctggaatgga | ccaaatacct | tgtttactga | 900 |
| aggatacaga | tggcatgtga | ctgttgagaa | tactcaccc | tcttagagcc | atggtttcct | 960 |
| cttctataaaa | atagggatgt | tcgtgcctat | ttgctaata | tgagtgacaa | tgacataagg | 1020 |
| tacataaagc | tgtactgcat | gatgcaaa | taacatcata | tttgtagggg | tggttgtaat | 1080 |
| aatactatcc | atcagcaaa | cagtcattca | tttactcagt | caaatactga | tgactgggta | 1140 |
| cattcagttt | cctcttttgt | aaaatgggga | taaaatagg | acttagctca | tagggatatt | 1200 |
| aagattcagt | gagttaatat | atataaaata | cttagagcag | tccttggaac | atattaagaa | 1260 |
| ctcaatacat | attagctagt | tgagcaggct | gtagtatttg | ttccagccaa | gaaaagactg | 1320 |
| ttctctaaca | gcacaggaaa | taaagatggg | gttaggcacg | acacggcagc | aggacttacc | 1380 |
| ttctgtctaa | ttcagctggc | agtcaaagaa | agaattatta | gaagcctatg | agcttgggtc | 1440 |
| cccaaagatc | tactgagtac | agggggatat | ttaaagaata | aaaatcccta | gaccacttta | 1500 |
| cagtacagga | gagacaagaa | gctgttcaca | caataacaag | tgctaaattc | cttgattttt | 1560 |
| atactgacag | ctgaagtgtt | agagaagaga | aggatcaata | aagaccggaa | tattaaagca | 1620 |
| gacaggccta | aaagaggatt | ttagatttga | taaagaattc | ctccagttct | cagagcaggg | 1680 |
| actttggagg | gtaacaactt | ggatttcagt | ccttgctgc | cacttactga | ttgtatgacc | 1740 |
| ttggaaaagt | tacttcactt | ctgtgagcaa | tgattctctc | atctggaaaa | caaaacaaaa | 1800 |
| caaaaaaact | aacaaggggtg | ataataatac | ctacctccct | acctcatagg | gctgatgaga | 1860 |
| agattaaaaa | gtacctacat | aaagcccttc | tctgcgtgcc | tgagcatgg | caagggctcc | 1920 |
| atgtgaacca | ctattttttt | ttttttttt | aatgaaaagt | catgggcagc | accaaagctc | 1980 |
| agaatttttg | cagtaaggaa | ttatgattct | acattgaaat | ttgccagaag | gggagctgac | 2040 |
| tgcctcatga | gacatttttg | aatgaggcca | aaaaaggaaa | caagtgtatc | ctgggatttt | 2100 |
| acgagatgct | aggtatgtcc | ttagcactta | atcctcatga | ctaccctatg | atgtaagtac | 2160 |
| tatctgttgt | ccctattttt | cagttggcca | aggtcacata | gctgaaacgg | acttctgtgt | 2220 |

| | | | | | | |
|------------|-------------|-------------|-------------|------------|------------|------|
| gtcttctaac | ttttgctttc | aggctctggaa | aaatgcaatg | taaacctaga | ccctctttga | 2280 |
| aatactgaag | atggtaatct | tatcccttcc | tcactttctg | ttttctaata | taacagtcct | 2340 |
| tggtccctac | aatgtctgtt | ttccagattc | ttagaagact | ttttgcttat | ttcccataac | 2400 |
| tctttacttg | tgatccctga | atgacaccgg | gggtatagca | gagaatgtcc | atttcctcaa | 2460 |
| agttcaaagg | tcctacaaaa | aatagttgct | agcctggcat | gatgggtgtg | gcctgtggcc | 2520 |
| tcagctacct | gggaggctga | ggcaggagga | ttgcttgagc | tgaggagttt | gaggcgcagc | 2580 |
| gtgatcgcg | ctcacgcgag | cctcccgggc | tcaagcattc | ctctcgccct | agcctcctga | 2640 |
| gtagctggga | ccacagctcc | actaattttg | aagttttttt | ggtagacatg | aagtctccct | 2700 |
| gtgttgcccg | ggctgggtctc | aaactcctga | cctcaagcag | tcctcctgtc | ttggcctctg | 2760 |
| gaaatgctgg | gattacaggc | gtgagccact | gtgctggcct | cttttttctt | tttctttttt | 2820 |
| tttaagggtt | ttatttggtta | aatgggaagt | ctgtgccatc | aactgagcat | tgtattttct | 2880 |
| ccttagtaag | agcctgggtg | ggccactggg | agagaactat | acattaaatg | taagtagcct | 2940 |
| ctgggtagag | agcccctggc | tggtttcctt | tcctttctct | ccttttctct | actttgggtg | 3000 |
| ctggaggcat | ttcccagact | ccagtttctt | accaccctca | cggattttgc | tattgtatta | 3060 |
| tcacctcctt | tatcattccc | aaaattgact | ttatggagac | tcattaaaag | aaagaatcat | 3120 |
| cggccgggag | cgggtggctca | cgccacgaag | gcgggcgaat | cacctgaggt | gcggagttcg | 3180 |
| tgaccagcct | gacaaaaaca | gagaaacccc | atctctacta | aacaatacaa | aattagctgg | 3240 |
| gcgtgggtgg | gcacgcctgt | aatcccagct | actggggagg | ctgggacggg | agaatcactt | 3300 |
| gaacccggga | ggcagagggt | gcagtgacca | aagatcgac | tattgcactc | cagcctgggc | 3360 |
| aacaagagca | aaactctatc | tcaaaaaaaaa | aaaaaaaaaaa | aaaaaaaaa | | 3408 |

<210> 31

<211> 1917

<212> DNA

<213> Homo sapiens

<400> 31

| | | | | | | |
|-------------|-------------|-------------|-------------|-------------|------------|------|
| acgcgtccgc | attctttttac | atgtagacat | ccagttttcc | caagtgtttg | ttgaagagac | 60 |
| tatccttgcc | tcattgtgtg | ttcctggtag | ccttggtgaa | aattagttga | acatatttgc | 120 |
| atgggtttat | ttctgagctc | tctattttgt | tccattgggt | atatgtctgt | ctttctgcca | 180 |
| gtaccgtact | gttttaatta | ctgtagcttt | gtaatatact | ttgaaatcag | gaaatatgaa | 240 |
| gcctccacct | ttgttcttct | ttctcaagat | tgtttttagct | atttgggggtc | ctttgtgggt | 300 |
| ccatatgaat | tttaggtttg | ttttttctat | ttccatgaaa | aatgccattg | ggattttgat | 360 |
| agggattgca | ttgaatctgt | agattgtctt | atgtattatg | gctattgtaa | tgatgttgaa | 420 |
| gtcttccaat | tcattgaacat | gggatgtctt | tttattttat | tgtgtctctt | aaaatttatt | 480 |
| tcacagtggt | tttatagttt | ttcatttgca | aatctttcat | ttccttgggtc | aatgttattc | 540 |
| ctaagtattt | tattcttttt | gatcttatag | taaatgagat | tgttttctta | atttgctttt | 600 |
| caaatagttt | gatgttattg | tatacagaca | cagctgattt | tgtatgttga | tttttatatc | 660 |
| ctacaacaaa | tttttctgaa | tttgtttagt | atgtctaata | aatttttgat | gcactcttta | 720 |
| gggttttcta | catataaggt | catgtcatct | gtgaacagaa | ataattttac | ttttcccttt | 780 |
| ctaatttgga | tgccctttat | ttctttttat | tgccgaattg | ctctgactgt | gacttccagt | 840 |
| actatatgga | atagaggtgg | taagaatagg | cattttggct | ttgtttctga | tcttaaggaa | 900 |
| agacttccag | tttttcacta | ttaagtctaa | tgtaactgtg | gagtttttca | tatatggcct | 960 |
| tcattatttt | gagatagtct | caccctgttt | gtagtttggt | gagatttttt | attatgaaaa | 1020 |
| agtgttgaat | tttgtcaaat | gcctttctgc | atctattaag | agatcttgaa | gaaagtgaag | 1080 |
| agtaaagagt | tactgaaatt | tcttgcttag | aagattagat | aaatgggtgat | gctactaatt | 1140 |
| aagggaaagt | agaaattttt | gaagaagata | aatagataag | tcttagacat | aatgtgtttt | 1200 |
| atttcttcaa | acattctggg | taaaaatca | aagtaacagt | tggttgcaat | ggaaaaata | 1260 |
| tgacagctctg | aggagacatt | ggaagcatcg | ataacttta | gaaaaacctt | acctaatgct | 1320 |
| gccccaaagca | ctttactctt | gtaaattcat | ctaattatca | ctgtaacacc | acaaagtagc | 1380 |
| tattagttgt | tatcttattt | tatagatgag | aaagctgagg | cacagagagg | ccatgtcact | 1440 |
| cgctgagggt | tccacggctt | ttaagtggca | gaacattgat | ttgagtcaat | gcattctgaa | 1500 |
| ctctgtgttc | atgttctcaa | ctattttatac | tattctgctt | cagaaataaa | gactcccat | 1560 |
| tactaagcat | tttaacatgc | tagacattgt | attgaacatt | atttgatca | tatcaattat | 1620 |
| ctaactctca | atacagctct | tttatttcac | agaataggaa | attgaagcaa | agagaggaaa | 1680 |
| agttgcatga | acgattttac | acagctgtta | ggctgttgaa | ctggattctg | aacctagcat | 1740 |
| tctggctgga | gagcccaagc | ttttatcagt | tgtgtatagc | agtgaagta | gattccccat | 1800 |

| | | | | | | |
|-------------|------------|------------|-------------|------------|------------|------|
| ggaaaggttaa | gagctgaata | cagagtccag | agggaaacacc | aatatgaaga | aggagttgtg | 1860 |
| aagcggcctg | aaaagaaatg | acaaagatag | gaagagagtt | acaaaaaaaa | aaaaaaa | 1917 |

<210> 32

<211> 2359

<212> DNA

<213> Homo sapiens

<400> 32

| | | | | | | |
|------------|-------------|-------------|-------------|-------------|-------------|------|
| cccgtgcgag | gtggc aaatg | caacatgctc | tccagtttgg | ggtgtctact | tctctgtgga | 60 |
| agtattacac | tagccctggg | aaatgcacag | aaattgccaa | aaggtaaaag | gccaaacctc | 120 |
| aaagtccaca | tcaataccac | aagtgactcc | atcctcttga | agttcttgcg | tccaagtcca | 180 |
| aatgtaaagc | ttgaaggtct | tctcctggga | tatggcagca | atgtatcacc | aaaccagtac | 240 |
| ttccctcttc | ccgctgaagg | gaaattcaca | gaagctatag | ttgatgcaga | gccgaaatat | 300 |
| ctgatagttg | tgcgacctgc | tccacctcca | agtcaaaaga | agtcatgttc | aggtaaaact | 360 |
| cgttctcgca | aacctctgca | gctggtgggt | ggcactctga | caccgagctc | ggtcttcctg | 420 |
| tcctgggggt | tcctcatcaa | cccacaccat | gactggacat | tgccaagtca | ctgtcccaat | 480 |
| gacagatttt | atacaattcg | ctatcgagaa | aaggataaag | aaaagaagtg | gatttttcaa | 540 |
| atctgtccag | ccactgaaac | aattgtggaa | aacctaaagc | ccaacacagt | ttatgaattt | 600 |
| ggagtgaag | acaatgtgga | aggtggaatt | tggagtaaga | ttttcaatca | caagactgtt | 660 |
| gttggaagta | aaaaagtaaa | tgggaaaatc | caaagtacct | atgaccaaga | ccacacagtg | 720 |
| ccagcatatg | tcccaaggaa | actaatccca | ataacaatca | tcaagcaagt | gattcagaat | 780 |
| gttactcaca | aggattcagc | taaatcccca | gaaaaagctc | cactgggagg | agtgatacta | 840 |
| gtccacctta | ttattccagg | tcttaatgaa | actactgtaa | aacttcctgc | atccctaattg | 900 |
| tttgagattt | cagatgcact | caagacacaa | ttagctaaga | atgaaacctt | ggcattacct | 960 |
| gccgaatcta | aaacaccaga | ggttgaaaaa | atctcagcac | gaccacacac | agtgactcct | 1020 |
| gaaacagttc | caagaagcac | taaacccact | acgtctagt | cattagatgt | ttcagaaaca | 1080 |
| acactgggtc | tcagcaaaag | gaccccgga | acattgcaaa | ctatttctaat | acctcagttt | 1140 |
| gaattgccac | tgagcactct | agctccaaaa | agccttccag | aatttctctga | ggcaaaaaaca | 1200 |
| cccttccctt | ttgagaaacc | taggggcaca | ttggcttcaa | gtgaaaagcc | atggattgtg | 1260 |
| cctacagcta | aaatatctga | agattccaaa | gttctgcagc | ctcaaactgc | aacttatgat | 1320 |
| gttttctcaa | gccctacaac | atcagatgag | cctgagatat | cagattccta | cacagcaaca | 1380 |
| agtgatcgta | ttctggattc | tatccacct | aaaacttcta | gaactcttga | acagccaagg | 1440 |
| gcaacactgg | ctccaagtga | aacaccattt | gttctcctcaa | aactggaaat | ctttaccagt | 1500 |
| ccagaaatgc | agcctacgac | acctgctccc | cagcaaaacta | catctatccc | ttctacacct | 1560 |
| aaacgacgcc | cccggcccaa | accgccaaga | accaaactg | aaagaaccac | aagtgccgga | 1620 |
| acaattacac | ctaaaatttc | taaaagccct | gaacctacat | ggacaacacc | ggctcccggg | 1680 |
| aaaacacaat | ttatttctct | gaaacctaaa | atccctctca | gccagaagt | gacacacacc | 1740 |
| aaacctgctc | ccaagcagac | accagtgct | cctcctaagc | caaaaacatc | accacgcca | 1800 |
| agaatcccac | aaacacaacc | agttcctaag | gtgccccagc | gtgttactgc | aaaacaaaaa | 1860 |
| acgtcaccaa | gtccagaagt | gtcatacacc | acacctgctc | caaaaagtgt | gtccttctct | 1920 |
| cataaaccat | acctgaggt | ctctcagagc | gaacctgctc | ctctagagac | acgaggcatc | 1980 |
| ccttttatac | ccatgatattc | cccaagtcct | agtcaagagg | aactacagac | cactctggaa | 2040 |
| gaaacagacc | aatccacca | agaacctttc | acaactaaga | ttccacgaac | aactgaacta | 2100 |
| gcaaagacaa | ctcaggcgcc | acacagattt | tatactactg | tgaggcccag | aacatctgac | 2160 |
| aagccacaca | tcagacctgt | tctgaatagg | acaactacaa | gacctactag | gcccacaccc | 2220 |
| agtggtgatc | ccagtgggaa | tggagtggga | acaggggtca | agcaagcacc | caggccatca | 2280 |
| ggtgctgata | gaaatgtatc | agtggtactct | acccacccca | ctaaaaagcc | agggactcgc | 2340 |
| cgccaccct | tgccaccca | | | | | 2359 |

<210> 33

<211> 527

<212> DNA

<213> Homo sapiens

<220>

<221> SITE
 <222> (284)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (292)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (422)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (514)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (527)
 <223> n equals a,t,g, or c

| | |
|---|-----|
| <400> 33 | |
| gggaggttgc ggtgagccga gatcacgcc a ttgcactcca gcctgggcga cagagtga aa | 60 |
| ctccgtctca aaaaaaaaaa aaaaaaaaaa attggccgta tgtgcgtggg tctgttcctg | 120 |
| agctctgttt ttttccatat atgtgttcat ccttttgcta atgccacact gtcttgctca | 180 |
| ctagaaattg gtaagtgtg tgaatccttc aactttgttc tttccaaat tgttttggct | 240 |
| attctggttc ctttaacttt catataaatc tttgaattag cttntagata anagcaaaat | 300 |
| gtcctgccaa gattttgtrt ctaaacttat aaatcagttt gaggagaatt agcatcataa | 360 |
| caatattgar tagtttwtac tgtgaacata gtgkaatgca ctatttaggt cttgatattt | 420 |
| tnatgggtgt tttgtatttt tcagtataaa gaccatgtgc atcatcttgt gtttagattt | 480 |
| ataccaaagc atttttgcag gtgttggaat tatnactggt taaaaan | 527 |

<210> 34
 <211> 1951
 <212> DNA
 <213> Homo sapiens

| | |
|--|-----|
| <400> 34 | |
| gcgtccgctt atctgaaatt gtaacctgcc tcctctcttt ggaactctca gtccctctga | 60 |
| tgttgttctt tctgtaattt acttatttat tctgtttatt gtgattatct attttgcccc | 120 |
| actctacact agaatacacac tccacgaggt cagggattag tgtttatttc gtttattgac | 180 |
| agatgccagg catttagaat agtgcctggc acataataaa ggcttattaa atagatgttg | 240 |
| aataaatgaa tcacctcttt atagtgcctg tgccctaaac tagatgatat cttgtctttg | 300 |
| taactttata gcacactgtg tagctctagt gatgcgtaca tgtatgctgg ttgtatcttc | 360 |
| caactttgct cttctcttcc tttaatatat aagggtctatt taggaattgt catctagcat | 420 |
| ttggatcagt gtccagtgat gttacatagt aggatttatc aaatttttaga aagagaaagg | 480 |
| aattttaccac actatcaaag aagtttaatt actatgtcct attttttggg ggaatcatat | 540 |
| ttaatgaaag agaattgtgaa attgctttgt cttatagttc agaaaccagt tttcaaaact | 600 |
| tccttcatgt agtaaatgga atcttctagt gaatgtcatg aaattttccaa attgtttttt | 660 |
| tttaattttt tttaaaattg tggtaagtac atataacata aaatttgcca ccttaactct | 720 |
| ttttaagtgt acacttcatt ggcattaagt acattctcac cgttgcataa gtgtcaccac | 780 |
| cgtccagttc cagaatattt ttaccttgca aacctgaaat tctgtaccca ttaaacagta | 840 |
| actcctcatt cctccccaac ccttagcaat caaagtttta ctttctgtac ctatgaattt | 900 |
| gatattctag gtacttttat ataagtga aa tcatacagtg tttgtccttt ttcactggct | 960 |

| | | | | | | |
|-------------|-------------|------------|------------|------------|------------|------|
| tattcactca | ccataatgtc | tgtaagacat | tatgttgtag | cacatatcag | aatttccttc | 1020 |
| ctttttaaag | gtgaataata | ttttattgta | tgtattgtgg | acatactttc | ttttcattac | 1080 |
| ctaaaaacct | tcctcatcct | acataaaata | tctgcagtg | gctgtctagt | atggatgaac | 1140 |
| cataaatcta | ccttctccaa | gattttctcc | cagtttttta | ctattttaaa | aagtgtctgc | 1200 |
| gggcgcggtg | gctcatgcct | gtaatcccag | cactttggga | ggccaaggca | ggtggatctc | 1260 |
| gagatcagga | gatcgagacc | agcctcgcca | atatggtgaa | accccatctc | tactaaaaac | 1320 |
| acacacaaaa | aaaaaagtgc | tataatacat | ttgcttgtaa | aaatatcctc | atcacttatc | 1380 |
| cacatatata | agccaggctt | ctaaaaatgg | catttctcac | ctgtagtccc | agctactcag | 1440 |
| gacgctgagg | caggagaagt | gcttgaacct | aggaggcaga | ggttgcagca | agctgagatc | 1500 |
| gcaccactgc | actctagcct | ggtcgataga | gcgagactcc | atctcaaaaa | aatgaataaa | 1560 |
| taaataaaaa | cagtatttct | gggacaaagg | tatacagacc | caaattttat | agcagttggt | 1620 |
| ttcattcatt | tacctataga | tattttaaaa | gcacctgcta | ggccgggcac | ggtggctcac | 1680 |
| gcctgtaatc | tcagcacttt | ggtagggcga | ggcgggcgga | tcacgaggtc | aagagatcaa | 1740 |
| gaccatcctg | gccaacgtgg | tgaaaccctg | tctctactaa | aaatacaaaa | attagctggg | 1800 |
| catggtgggtg | cgcgccctgta | gtcccagcta | ctcaggaggc | tgagaatcgc | ttgaacctgg | 1860 |
| gaggtagagg | ttgcagtgag | ctgagatcac | accactgtct | tccagcctgg | caacagagcg | 1920 |
| agactctgtc | ttaaaaaaaa | aaaaaaaaaa | a | | | 1951 |

<210> 35

<211> 2024

<212> DNA

<213> Homo sapiens

<400> 35

| | | | | | | |
|-------------|-------------|-------------|-------------|------------|------------|------|
| acgcgtccgt | tgtgtgggac | aagctgctcc | cgacagaagg | atgtcgtgct | tgagcctgcc | 60 |
| ctggctgggc | ctcagaccgg | tggaatgtc | cccattggcta | ctcctgctgc | tggttgtggg | 120 |
| ctcctggcta | ctcgcccgca | tcctggcttg | gacctatgcc | ttctataaca | actgccgccg | 180 |
| gctccagtgt | ttcccacagc | ccccaaaacg | gaactggttt | tggggtcacc | tgggcctgat | 240 |
| cactcctaca | gaggagggtc | tgaaggactc | gaccagatg | tcggccacct | attcccaggg | 300 |
| ctttacggta | tggtctgggtc | ccatcatccc | cttcatcggt | ttatgccacc | ctgacaccat | 360 |
| ccggtctatc | accaatgcct | cagctgccat | tgcacccaag | gataatctct | tcacaggtt | 420 |
| cctgaagccc | tggtctgggtg | agtacctgca | ggtgaaaggg | ggtggggaca | accttgccgg | 480 |
| gagggtaggg | gaagtgtctg | tcttgcccat | tgtccttggc | tgccctacta | ggagaaggga | 540 |
| tactgtctgag | tggcggtgac | aagtggagcc | gccaccgtcg | gatgctgacg | cccgccctcc | 600 |
| atttcaacat | cctgaagtcc | tataatacga | tcttcaacaa | gagtgcacaa | atcatgcttg | 660 |
| acaagtggca | gcacctggcc | tcagagggca | gcagtcgtct | ggacatgttt | gagcacatca | 720 |
| gcctcatgac | cttgagacgt | ctacagaaat | gcattcttcag | ctttgacagc | cattgtcagg | 780 |
| agaggcccag | tgaatatatt | gccaccatct | tgagctcag | tgcccttgta | gagaaaagaa | 840 |
| gccagcatat | cctccagcac | atggactttc | tgtattacct | ctcccatgac | gggcggcgct | 900 |
| tcacagggc | ctgcgcctg | gtgcatgact | tcacagacgc | tgtcatccgg | gagcggcgct | 960 |
| gcaccctccc | cactcagggt | attgatgatt | ttttcaaaga | caaagccaag | tccaagactt | 1020 |
| tggtattcat | tgatgtgctt | ctgctgagca | aggatgaaga | tggaaggca | ttgtcagatg | 1080 |
| aggatataag | agcagaggct | gacaccttca | tgtttggagg | ccatgacacc | acggccagtg | 1140 |
| gcctctcctg | ggtcctgtac | aaccttgcca | ggcaccacga | ataccaggag | cgctgccgac | 1200 |
| aggaggtgca | agagcttctg | aaggaccgcg | atcctaaaga | gattgaatgg | gacgacctgg | 1260 |
| cccagctgcc | cttcctgacc | atgtgcgtga | aggagagcct | gaggttacat | ccccagctc | 1320 |
| ccttcatctc | ccgatgctgc | acccaggaca | ttgttctccc | agatggccga | gtcatcccca | 1380 |
| aaggcattac | ctgcctcatc | gatattatag | gggtccatca | caacccaact | gtgtggccgg | 1440 |
| atcctgaggt | gctgccttcc | ccattcacca | ccaccacccc | catcctctac | ttttgtgtgt | 1500 |
| gtgtgtgtga | attccaaggt | tcctagtggg | gggggcaggg | ttttgatgag | gaaaatccaa | 1560 |
| catcacctca | ccccaaaaca | cacacaagtg | tctctccaag | gctgctggac | ataggaaatc | 1620 |
| ccactggcaa | accttctttg | tctcacctgc | aggtctacga | ccccttcgcg | tttgacccag | 1680 |
| agaacagcaa | ggggagggtca | cctctggctt | ttattccttt | ctccgcaggg | cccaggaact | 1740 |
| gcatcgggca | ggcggttcgcc | atggcgggaga | tgaaagtgg | cctggcggtg | atgctgctgc | 1800 |
| acttccggtt | cctgccagac | cacactgagc | ccgcagga | gctggaattg | atcatgcgcg | 1860 |
| ccgagggcgg | gctttggctg | cgggtggagc | ccctgaatgt | aggcttgacg | tgactttctg | 1920 |
| acccatccac | ctgttttttt | gcagattgtc | atgaataaaa | cgtgtctgtc | acctcaaaaa | 1980 |

aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaa

2024

<210> 36

<211> 3710

<212> DNA

<213> Homo sapiens

<400> 36

| | | | | | | |
|-------------|-------------|-------------|-------------|------------|------------|------|
| gtaccacaaa | ttttcaggtg | ggtgtgtaag | agctaattggg | gccacagatg | ctatgcatac | 60 |
| cactgcctag | ttgttgctag | agagcctttt | caacctcaga | gaacatagca | ttccatgcta | 120 |
| accatcattt | aaaattcctg | caataaatga | aaaaaataaa | cacagagctg | agagtataat | 180 |
| tcaactagcc | tcgctcccaa | atgaagctga | ttccagatag | aaagcagatt | ttcatattta | 240 |
| ctgtatgtga | cagctgcagg | aattaaatga | atcatctgtc | tatctctatt | gccttggtcc | 300 |
| tcctgtgctg | tgtgcatcta | tccttggggg | tgtctgtctt | cccattccag | gaagacagat | 360 |
| cagtgtagtt | tcaattatta | attccatggt | tggatcacag | taacaccagt | tccagttcag | 420 |
| tgacctgcag | ttcctattga | tttaagtgat | gaattatcta | cactgcaatg | tattattaac | 480 |
| tttattttgt | ctcttattcc | ttctgcattc | atgcatcaaa | ataataaagc | accattcaca | 540 |
| ggctaaaagg | acccgtttcc | cttctcacat | ttcccataag | ggcgaagcca | acacacacca | 600 |
| gggaggtaat | tacacagagc | tggggtgggg | gctggacatt | tattttacct | ctgaattatt | 660 |
| tatctcagct | gtgaatttgg | gggagggttc | ctcagtgagg | aacaaagggg | | 720 |
| cccaggaggg | aagttaatga | agacaagtga | cgattagaac | ctgggagatt | cataagaaat | 780 |
| gaaataacct | caacccaaaa | gtttaacttc | tgggagtcaa | tctgcccatt | ctgataatga | 840 |
| tgggggggatg | ggtgtaggag | tggtaacgac | agtggcacag | tgtgaaggga | aggaggcaag | 900 |
| agaagagagc | accatcagtt | tctcctgcca | tctcttttgg | attgctctgc | ctgccccatc | 960 |
| gcaagacctt | ctcagcaagg | tcagactcca | gtgagacatg | gatttgtgga | tgtttggcag | 1020 |
| aaatgtgctg | gccttatggc | taacctacca | ttgtcttcat | cttctttggg | taaaagcaca | 1080 |
| tcctcctatt | accttttgag | gaacttcctt | tccttgggta | caatctggta | ggactatcaa | 1140 |
| tcagatgcc | ctaccttgct | ctgaccaa | gcattgaccca | agctgggtca | gtgaggttct | 1200 |
| ccctccatga | catttgaatt | acagacaaag | ggcacaaaag | accgagggtt | agtggcatct | 1260 |
| gttcaactag | gctgatgcct | gggtctgact | gttgagcaat | tgctgctctc | aataccttct | 1320 |
| gaagatgatc | tgcaagcctt | gtttttcaac | cttccctttg | attctagatg | ccccacccca | 1380 |
| tagtctttca | ataaatgctt | ttttttttaa | gttaactaaa | tttgagtgtg | ttttgttgct | 1440 |
| ttcaacttag | agaactctgg | ctaagcagcc | aaaagacatg | gctgtatcta | aagtgtccct | 1500 |
| gagccctcag | agatggaaaa | ggtgaagagg | gaggggctag | atcttgtcta | cacctcttct | 1560 |
| gccaatcaag | caacaattac | atcaataggg | agatgggcag | gaggtgaagt | gaaggcattc | 1620 |
| ctctgtcctc | tcgctcccat | ttctagttag | aaactgaatc | aacctcccag | cctgcaggac | 1680 |
| tcctgttggg | gcaactccca | aagtcaagga | gtctgaagag | acacctcatc | tgacaatccc | 1740 |
| acacagaggg | aagacttctg | gaggggttgc | ctaggggtcag | gcaccccagg | catcccgcct | 1800 |
| atgagccaaa | cccaagagga | aggaaagaca | cccaggacct | cccagtttga | aagtgaagat | 1860 |
| aaagctgcta | agtgcgtaga | agtttccatg | tcagacaaat | tggaggatta | cctaaggctt | 1920 |
| catttctagt | gctattacac | tcagaaattg | cctcacctat | gactgctccc | attttgtgac | 1980 |
| tcacagctct | ctacgctgca | ctggatgaa | acagatgttg | tgaatgtttt | attgtagttt | 2040 |
| ccaattgcc | cccttcattc | atctgattca | gacctaaag | ccttgggctt | tcctttcttt | 2100 |
| tctttctcct | cctccaccct | ttgattttaa | gggcctagag | gtgaacatct | aacaggcaga | 2160 |
| acctgattct | acctgtcaac | tcaccccctt | atagcaatgg | agaatagccc | agcaatgcct | 2220 |
| gagaaatgag | ttctttttct | ttccagaact | tcaccacata | aggagagaag | gatcagaaaa | 2280 |
| ccatgtgatt | agattatatg | gaagaaggtc | aaagattaaa | aatccactgc | ctctcattat | 2340 |
| tagaggaggg | atgacagatt | cagaccagag | cacaaaagaa | aatacattat | ttcaaaacag | 2400 |
| gatatgaaac | ccagtcccag | gtgattagtc | tctgttaagt | aaatttacta | tctaattctt | 2460 |
| taactgtcct | ccgaaattgc | tgggtccaaca | ttctctccct | cctccattgg | aagcttaacc | 2520 |
| ctgggagggt | gctgtcatcc | accttaattg | aatggggagc | tcatttcagg | ggggtgtagg | 2580 |
| tgggggtggg | gtggctgggg | tggaaagggtc | tgtaaagtga | atgcaccctg | taataaaagc | 2640 |
| aggatgtgtg | ctttgcagac | cagcctgact | ggaggtagga | tggggatgga | ggttggggca | 2700 |
| attgtccttc | catttggtttc | agagctaggg | ccataaatga | ggagactatc | cttcaacctc | 2760 |
| actccaggct | catcctggga | aaactggact | ctcctctttc | cgcacagccc | caaggggccc | 2820 |
| acatgtcaaa | gaatctcatg | cttcgggggt | gctctcaaac | cagatcaacc | ttcaaagctt | 2880 |
| tgattttaaa | cgaatgctgc | tctgcaggct | aaatatcaca | ggcctgtgtt | ggggcccgaa | 2940 |

| | | | | | | |
|------------|------------|------------|------------|------------|-------------|------|
| gaggacaaga | tgtgcacttg | gaggccagac | aggcttgcaa | caccatccaa | gcaatgagaa | 3000 |
| aacaagacat | agtggaaaag | aggatctgtt | tctcagtatc | tgtcttggct | ggggtaccac | 3060 |
| tgtgaacatg | gcatgcaaca | atcagagagg | cagaggatac | caaacccttc | ggaactcttc | 3120 |
| ccctgtttac | caggaggaac | tccttttttt | ctgtacttca | ctatcttctc | gtctgtttag | 3180 |
| tctaaaaggc | tgaactatta | cacaaagaaa | gttctctcca | agaatgagtt | ttccagagaa | 3240 |
| agctcttcag | acaagggcag | tgtttgaacc | actctccaca | gcctggagaa | tgggtgtgtg | 3300 |
| gtttaacaga | agggggctgg | ggccaaacag | gactccctga | aggattccca | aggtgctctg | 3360 |
| tttgtgtgat | ttatcccttc | tggaaaatcc | aaacatacaa | atgtgcctgc | tgcaaaaagtc | 3420 |
| agggcagtct | ccaggactca | ggttccccca | caaagctaac | cacatatagc | ttgcctgccc | 3480 |
| tgaaggtttt | ctctatttta | accttcgtag | gttttcaacc | aaaacagatg | aaaggagggt | 3540 |
| gcatgtagtt | ttgctgggga | tcagcagacc | tcatttctct | ctctcttctt | agctcctagg | 3600 |
| ctcacttcac | aaaaaacctc | cggagaacag | gaactgggca | aaaaggggct | tcctcagct | 3660 |
| cctgggattt | cagaaccaga | cattgtcatc | ctgttcacca | agcagtcgac | | 3710 |

<210> 37

<211> 343

<212> DNA

<213> Homo sapiens

<400> 37

| | | | | | | |
|-------------|------------|------------|------------|------------|------------|-----|
| ccacgcgtcc | gcgagcgcgt | gggcgtgacc | ctgcgtgcaa | acatgtaccg | tggcctggta | 60 |
| tatgatagag | attgatatta | atgtaccatg | tatgttaatg | tgaatctgtg | ggcaggatac | 120 |
| ttttccatgg | caggaaatat | ccaagctgtt | gaaactggct | atgttttaat | atgcctcatt | 180 |
| gtgcctttac | tggtgtgtgg | actgcgtgag | ggacaagaag | ttccatttga | tgtcaataaa | 240 |
| gcaaaagtact | tgcctacttt | tttgaaaaaa | aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | 300 |
| aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | aaa | | 343 |

<210> 38

<211> 445

<212> DNA

<213> Homo sapiens

<400> 38

| | | | | | | |
|------------|------------|------------|------------|------------|-------------|-----|
| cgacccacgc | gtccgtggca | ggcttccaaa | tttgccaaaa | ttaagcggtc | agtattttca | 60 |
| acacatacgc | tttttactgg | tttatactga | actatcggat | gagaattcct | gtgttcccaa | 120 |
| agcaactgat | gttttaeagg | cttgtgtttc | tcctcctcct | ttctaaggat | gaggggaatcc | 180 |
| acaacagact | ttctctagaa | aacactaatg | atggacaact | ttttgggtgc | atcaaataag | 240 |
| ttggctacta | ccttgatggg | aaaaatttgt | aaggaaaaat | ttcaccattt | cgagtgggtca | 300 |
| agtgtatttt | taactgtctg | gtttgtactt | ttatgacttt | tgtactacca | aacgcggagt | 360 |
| taaaaataaa | aatgttaaac | gcaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | 420 |
| aaaaaaaaaa | aaaaaaaaaa | aaaaa | | | | 445 |

<210> 39

<211> 1312

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (16)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (18)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (19)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (21)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (25)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (1162)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (1181)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (1275)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (1288)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (1291)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (1294)

<223> n equals a,t,g, or c

<400> 39

| | | | | | | |
|-------------|-------------|------------|------------|-------------|------------|-----|
| gcggggggggg | aacggnngnnc | nccantcggg | ttcaagtggg | tggcgggatgc | ccacagttcg | 60 |
| gccatcctga | gcatgctccg | tggggccccg | gacagtgcc | cggcacaggt | gtgcacagcg | 120 |
| ctcagcctct | gtgagccgct | gcagaggcac | ctggccaccc | tgaggccact | ctccaaagag | 180 |
| gacacctttg | aggtctgtggc | tccgttcacg | gccaatgggc | cccttacctt | ccacccccgc | 240 |
| caggcgcctg | aaggagctct | gtgccaagac | tgtgtacggc | aggtctcccg | actccaggag | 300 |
| gctgtccggt | ccaacttgac | cttggccgac | ttgaacatcc | aggagcagtg | tgagtccttg | 360 |
| gggcctggcc | tggccgtcct | ctgcaagaac | tacctcttcc | agttttttgt | ccctgctgac | 420 |
| caagcactga | ggcttctccc | cccgcaggag | ctctgcagga | aggggggatt | ctgtgaggag | 480 |
| ctaggggcac | ctgcccgttt | gactcaagta | gtggccatgg | acgggggtccc | ctccctggag | 540 |
| ctgggggttg | caaggaaaca | gagcgagatg | cagatgaagg | ccggtgtgac | ctgtgaggtg | 600 |

30

| | | | | | | |
|------------|-------------|------------|------------|-------------|-------------|------|
| tgcatgaacg | tggtgcagaa | gctggaccac | tggtcatgt | ccaacagctc | tgagctcatg | 660 |
| atcacccatg | ccctggagcg | cgtgtgctcg | gtaatgcctg | cctctatcac | gaaggagtgc | 720 |
| atcatcttgg | tggaacaccta | cagcccctcc | ttggtgcagc | ttgtggccaa | aatcacccca | 780 |
| gagaaggtgt | gcaagttcat | ccgtctgtgt | ggcaaccgga | ggcgggcccg | ggcagtcctat | 840 |
| gatgcctatg | ccatcgtgcc | gtccccagag | tggtgatgcg | agaaccaggg | cagctttctgc | 900 |
| aatgggtgca | agaggctgct | caagggtggc | tcccacaacc | tgagagagcaa | gagcaccaag | 960 |
| cgagacatcc | tggtggcctt | caagggtggc | tgcagcatcc | tgccgctgcc | tatatgatcc | 1020 |
| aagtgcaagc | actttcgtca | cccaatacga | agcccgtgct | cattgagagt | ctcaaaggac | 1080 |
| atgatgggac | cccgtggctg | tgtgcaagaa | ggtggggggc | ctggcacggc | cccaagaccc | 1140 |
| cacttggttg | gcaccgacca | angtggcctg | gccccagctt | ntggtgcaag | aaccaagaag | 1200 |
| gccgccagct | ttggcaacgc | tttgcaacac | tggcagaaac | atggatggga | aagagatgcc | 1260 |
| cttcaccctt | ggggnaacac | gccgggancc | nggnttggca | aaaaaaccca | aa | 1312 |

<210> 40

<211> 1253

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (1188)

<223> n equals a,t,g, or c

<400> 40

| | | | | | | |
|-------------|------------|-------------|-------------|-------------|-------------|------|
| gacaacatgg | ctgacgtgtc | agccaaggac | agcagccagg | agacgctggt | gaacctggcg | 60 |
| gggtctcttg | tcagcctcct | gatgctccct | ctggtgtcag | gttgccctgg | cttcagcctt | 120 |
| ggatgtttct | tcttcctcac | tgccctccac | atctacgcca | actaccgcgc | ggtccgagcc | 180 |
| ctggtcatgg | agaccttgaa | cgaaggccgg | ctccggctgg | tcctgaagca | ctaccttcag | 240 |
| aggggagagg | tactcgaccc | aactgcagcc | aatcgcatgg | agccgctgtg | gacagggtttc | 300 |
| tggtccagctc | cgtctctatc | cctgggggtc | cccttacacc | gcttggtctc | cagtgtcttt | 360 |
| gagctgcagc | agctggttga | ggggcaccaa | gaatcctacc | tcctctgctg | ggaccagtca | 420 |
| caaaaccagg | tacaggtagt | tctgaaccag | aaggcaggcc | ccaagaccat | cctaaggggc | 480 |
| gccacacatg | ggctgatgct | tggggccctg | caggggagatg | gaccccttcc | agcagagctg | 540 |
| gaggagctga | ggaaccgggt | gcgggcagggt | cctaagaaag | agagctgggt | cgtcgtcaag | 600 |
| gagacacacg | aagtgttgga | catgctgttc | ccaaagtctt | tgaaaggact | gcaggatgcc | 660 |
| ggctggaaga | ccgagaagca | ccagctagag | gtggatgagt | ggaggggccac | atggcttctg | 720 |
| tctcccgaaa | agaaggctct | gtgagcagcc | cagacggagg | cccaagccca | gggcagggaac | 780 |
| ctggagcaag | gacacttttg | ccacagcagg | acaggggaaa | ggcagcttta | tttttcctta | 840 |
| gggcaactgc | agcgggtggg | ccaggccctc | atgggaagtg | actgccaatc | agatgcagtg | 900 |
| ggcccagggc | agaggaaggc | cgggagaagg | ggagccagga | ccttctcacc | ccactgcccc | 960 |
| ttcccctttt | ctggggagca | ccgcaggctc | ctcaccccca | cttcctgtga | ggctgtggct | 1020 |
| tatggtgtcc | aacgcagttg | tcttaggcac | agaagcccca | gaggaacacg | gccactgcca | 1080 |
| tcatgagcag | ggcattgagg | ttgaccacac | gggccagct | cgggtcctcg | ctgatgtcct | 1140 |
| ccagccgctt | ggctgctgcs | gctgcytctt | cctggktaag | gggcgranga | stgycacccc | 1200 |
| acctctgcty | attccacama | accagagcag | gmactggcgc | gaagaagggc | ttg | 1253 |

<210> 41

<211> 958

<212> DNA

<213> Homo sapiens

<400> 41

| | | | | | | |
|------------|------------|------------|------------|------------|-------------|-----|
| tttttttttt | tttttttttt | tttttttttt | tttttttttt | atttaaatac | atttatattga | 60 |
| accggccttg | gggaggcttg | atgttgatgg | gttggtgatt | aaagcctccc | aaagccaatc | 120 |
| cttggcatgg | cctttgggac | tcaaaacaca | ggatctgact | ggtgggcaca | ataccatctt | 180 |
| gaacgcccac | aaaaagggtt | ggttttgttg | ccaaagggca | ggtggctcca | ggcagggctg | 240 |

| | | | | | | |
|------------|-------------|-------------|------------|------------|------------|-----|
| atggtggcag | ggtgggggtga | ggacaggaca | agagatctgg | gtgtggaagg | atggccgggt | 300 |
| ttctgcagca | gcaggaggaa | aggggtgggag | cacacaggca | cagaacactg | tgagcaggac | 360 |
| tgccaggcca | gtgtcacaag | cgctaccatc | tgggtgtaga | cagacctgag | ctgacaaagc | 420 |
| tgggggagca | aggccaacgg | ctttaaacac | aagctcaggg | gcttgggggt | tatcccagg | 480 |
| gcacaggcca | gccatgtagg | gtggagttgg | catgagttaa | gcctgggctg | ggtgtgtaga | 540 |
| ctggacagag | gtgggtaggg | caggcagtga | cgagctggtg | tgcaagagac | tttagggcca | 600 |
| ggcatggggg | gacagaactc | ccacctcggt | cgtgctccca | ccctccccag | ctccaccgcc | 660 |
| tggtcttcag | tctccggcat | agcaaggagg | aacgggagga | cctggatgct | gatgagcagc | 720 |
| aaggctcctc | actccctgta | cagaatgggt | gcccagagag | tgccatggag | atgaatggta | 780 |
| gggcaccatg | ctgggaggtg | gggctggagg | agctgagttc | ccgcaaacta | actgcagggc | 840 |
| ctcaatttcc | ctcagagccc | caggccccgg | caccaagcct | cttcgccag | tgctgtctct | 900 |
| ggttttgtgg | aatgagsara | ggtgggggtg | gcagtcctcc | gccccctacc | caggagga | 958 |

<210> 42

<211> 2197

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (1975)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (2006)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (2047)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (2133)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (2177)

<223> n equals a,t,g, or c

<400> 42

| | | | | | | |
|------------|------------|------------|------------|------------|------------|-----|
| gctctgtcac | tggtgccttc | tggaagatgg | cttccccttg | gggagcattt | cctgcaaaga | 60 |
| ctcacaagtt | taaaataatt | taggatattt | ctctgtactt | gagagccagg | ataggatgag | 120 |
| agttggctct | gccctgatgc | tgtaaggcca | ctctgtcttc | agacctgag | gtcctcagga | 180 |
| gtggtttccc | aagtactggt | gtagtgtgat | ggcaagatca | ctcttgatta | ttttaggagc | 240 |
| tgattttact | tttctacct | cttttaactg | cttccaaaag | atgaatctgg | ccaagaagag | 300 |
| caggggctca | ttcactcatt | tactcactca | ttcctgggtg | ttgagtttat | tcctgaagga | 360 |
| agctgaccaa | gggttgagag | aaaacaactt | tgatttttct | catgtctgcc | ccagcaaacc | 420 |
| tcctttgtgg | actgattctc | cttctgtgcc | tggaagaaac | tggaataatc | ccaggacttt | 480 |
| tctggtcctt | tctaggtaat | tcagtgtctt | ccactaatgt | ggccccacac | aggctgggct | 540 |
| tgttgatagt | ataggtccag | caaaacagtt | aaaactctac | atgtcttcag | catgtgaacc | 600 |
| tcttcccaac | ctcacctggg | gctgtggcag | gtgggaggct | gggtgtcagg | ctggctgctg | 660 |
| ccccctgggt | tctgcccagc | tggtagctat | ggcttcaacc | ctycagtggg | aggaacaaga | 720 |
| gggataagag | gkaaggrcag | garaggttct | gtatggctgg | catggcagag | agctggcaca | 780 |

| | | | | | | |
|-------------|-------------|-------------|-------------|-------------|------------|------|
| tgaggctggy | aggaratcca | cagggctctg | ggggcattct | catggctctcc | taagtgacca | 840 |
| cactaccctg | gggtgtctct | gccacaactg | gccaaactcag | gaggtgtatc | ctctccagct | 900 |
| gggtcccttac | actcacttcc | ttagccctag | gtggccgtgg | tacctctagc | ctcttgccaa | 960 |
| ggctctgccct | cacatctcca | ggaacccttc | tgggcaggat | ttaagacatc | ccaggccagt | 1020 |
| gttcaggcac | tatgacccca | cctgggtccac | aggaaatgtg | cacctttgat | ccactggggt | 1080 |
| aggggtatatt | ccacatgcag | cctagtgtgc | tactgggagc | tccaggagtc | aacttttgcc | 1140 |
| cttttagattt | cttgagtgtg | agaggggatac | caggtctctg | cccaccacca | gcctcagaag | 1200 |
| gcacacagta | ggcacttgca | ggcctcgtga | aaccctctt | ggaatgaagt | aaggagaaag | 1260 |
| ctgccccatcc | ctctcatggt | acagggttggg | tggggacaga | ctgagagcac | tgccccctc | 1320 |
| tcccaaggaa | gccctcctgt | caaacttgag | cttctctcat | cttcaacacc | tcctctacct | 1380 |
| agatcaccca | gaatttcaca | tgggaacaak | tggcacttcy | tgtcttgggg | cctcagctaa | 1440 |
| aattcagact | ggaaaagcty | catattaaca | tcctgktcta | atgcaagtaa | aaaaagagct | 1500 |
| gctatggctct | ttgsttcttg | gwatttagag | tctagtggga | gaatattggc | tacttaatgc | 1560 |
| agtctcttcc | attttcaata | tgtaaactgag | gcaccttct | actgataaca | gccaaaactt | 1620 |
| caagaaaaaa | tattttaagc | ctttaaaaat | acatttgtgg | gctggtagga | aattaaggaa | 1680 |
| taatcagaga | ccaaaaacta | agtgaagctg | agaaccaga | gataacttta | ttaaaggaga | 1740 |
| atattaaaca | tatacttcac | gcagaaggaa | gagaattcca | ggtggaagat | ttgagatgca | 1800 |
| aggggggaatg | aaaaacaaga | aggtgggtgac | tatatgggta | aatcaaatca | aatcaaacac | 1860 |
| cttgtgtaaa | ataataatgg | tgatataaga | attttaaaaa | gagaaataag | tatatagtca | 1920 |
| taatatgata | caaatttgaa | gaggggtgatg | agttaaaaag | ttctaagatt | cttngttaa | 1980 |
| ctttggagga | tggttatcta | atttgnaata | aaattttcag | gataatcatc | aaccgtagaa | 2040 |
| atggggntga | aattttcaaac | taatggaaaag | aaaaaagaat | gagaaaatat | aatcaatcaa | 2100 |
| aagggaagaa | actagaaaag | gaaaggaaaa | aangaaacac | aaaagagggtg | ggaacccaaa | 2160 |
| agtcacaaat | atatcantaa | ctaccattaa | aacataa | | | 2197 |

<210> 43

<211> 403

<212> DNA

<213> Homo sapiens

<400> 43

| | | | | | | |
|------------|------------|-------------|------------|-------------|------------|-----|
| gaacctcttc | ccaacctcac | ctgggtgctgt | ggcaggtggg | aggctgggtg | tcaggctggc | 60 |
| tgctgcccc | tgggttctgc | ccagctggta | gctatggctt | ccaacctcc | agtgggagga | 120 |
| acaagaggga | taagaggtaa | ggacaggaga | ggttctgtat | ggctggcatg | gcagagagct | 180 |
| ggcacatgag | gctggcagga | aatccagagg | gctctggggg | cattctcatg | gtctcctaag | 240 |
| tgaccacact | accctgggg | gtctctgcca | caactggcca | actcaggagg | tgtatcctct | 300 |
| ccagctggtc | ccttacactc | acttccttag | ccctaggtgg | ccgtgggtacc | tctagcctct | 360 |
| tgccaaggtc | tgccctyaca | cttcaggaac | ccttctgggc | agc | | 403 |

<210> 44

<211> 440

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (430)

<223> n equals a,t,g, or c

<400> 44

| | | | | | | |
|------------|-------------|------------|-------------|------------|------------|-----|
| gcggccgccc | tttttttttt | tttttgcatt | gttcaaatgt | tctataacct | ttactggctt | 60 |
| ttatttactt | cttctataaa | atactgagtt | acagtattta | aaatatttct | ctctgatgat | 120 |
| gaatttatat | attcttcctt | atagttctgt | caatatttgc | tttacagatt | ttgaaattat | 180 |
| gttatggaaa | gaataaaaatt | caaaattata | ttatctgctt | gatgaaatga | atwgtttaat | 240 |
| caktacatag | tgattctctc | tatctccagt | gatgcytttt | accccaaaat | caawtttttt | 300 |
| acattaatat | agattcatca | tatttccttc | agttatatatt | ccccagtata | ccctttccca | 360 |

| | | | | | | |
|------------|------------|------------|------------|------------|------------|-----|
| tccttttact | ttcatccttt | ctgtaccctt | agagtgtaga | tgtgttttct | gtaaacagca | 420 |
| tatagtaatn | cactttaatc | | | | | 440 |

<210> 45

<211> 3288

<212> DNA

<213> Homo sapiens

<400> 45

| | | | | | | |
|-------------|------------|-------------|-------------|------------|-------------|------|
| ccacgcgtcc | ggggaaaagg | gcatgggttg | gaggttgagc | agtaagagac | cattaatact | 60 |
| aagtattttt | cctgaaatgc | ttctttataa | ttatttcagg | gaaggaaata | gataatctca | 120 |
| gacttactgg | ggtggggcct | ggggaatggt | tggatgttgc | tctttttctt | cttctctctt | 180 |
| tttttttttt | tttttttttg | gttgggtgtg | tttggcattt | ttttcttttc | ctttttaaag | 240 |
| aaaatgtttt | caggaaacat | gaacaaacat | actgccattt | atagtggggc | tgggaaggca | 300 |
| caggagctgg | cgacctctca | gctgcatagt | tgggacggca | aaccctgctg | tgagcttctc | 360 |
| agacttttca | cctacttcac | ctattaagaa | gccatgagga | gttgtaaact | cttgttcagg | 420 |
| gaagaaaaga | gggcaggagg | aattaaccca | atgcctttta | atacaaaaac | gatgaaaaac | 480 |
| aaaatactca | tttttctctt | tctattgtgg | gtttggggag | ccaaaccaag | tgtgactgtg | 540 |
| agcagtttca | ctgagaggca | tgacaacatt | attgaagagg | tctttccatt | atgtctgggt | 600 |
| ctcttatttt | cttaagtcaa | gccttaacta | tgaacatact | ttaagatgat | acgggtctgt | 660 |
| caccataaac | ctgtaaaagg | gcatcgacag | cttttcagat | aatccaggaa | tgttgagaca | 720 |
| tggctaagtt | tctagctgat | acttagtcat | tccctcttga | tattgttggg | atgggtagcc | 780 |
| aaaaggcagg | cttgtgggag | ttaagagcca | atgatacctg | gaaacatggg | aggtagctgg | 840 |
| gggtacaagt | ggatgttgag | actgaaggaa | gaggtggaaa | gaagacacag | aaggcaccca | 900 |
| gatgccaaaa | ggcctggcat | gaatgagagt | tggcatctgt | cctgttgtgg | tgttcccat | 960 |
| cttgccatac | ccatttgccc | agattgttct | gattctcatg | ttcaattgac | tatcccacat | 1020 |
| gagactacag | tttataggca | gtgaggaatt | gtttcttcc | gggaaatgtc | cttccagcaa | 1080 |
| cttcagagga | atgggaaagt | gttaatctta | cgtgaagcaa | aggagcctgg | actaggggca | 1140 |
| gagagaaaaga | agcagcagcc | caccttcttt | gtgtttgcac | atgtgcctat | gtcgtgtaca | 1200 |
| ttggtacctg | actgtggctc | agatctgcgt | cgcagcagcg | agagaagaaa | tcactccata | 1260 |
| tccgatgaga | ggaagggtgg | cacagagatg | gtgtctacaa | ttagagacat | ttctgacttc | 1320 |
| accttagcct | aagcaaactt | tatatactga | gtaacatttg | aagggtgctt | ttaatgggtg | 1380 |
| ggggtgtttt | ttccttttta | aactacagtg | cttgccacaag | agagggaggg | actcagaaaa | 1440 |
| ggttagggca | ggtgagggag | acagtagatg | gcctgggatg | acttgagtcc | atcatactat | 1500 |
| tgcttggcag | gtgtcctccc | ccatgtttga | ttcaaattcc | atgagtgacc | tacctttccc | 1560 |
| caggaatggg | actgagaggg | tagtctccag | caactcagtc | tgacaggggc | tccccgttca | 1620 |
| ggctgccttt | ggtggttgtg | cttttgtaag | tttctttctc | tgcaacttga | cttacctctg | 1680 |
| aatcagaaag | caagcccagc | agggtgaatga | gggatgtctg | cttggcattg | cccaatctaa | 1740 |
| ccagggaggc | tggctggcca | cccactgtcc | gctagagggg | agagctagca | ggtgttggtg | 1800 |
| tgaactcagg | aatagaaaca | cgaggccttt | ttaaatacga | gggagaagaa | tccatgatgc | 1860 |
| atactgttaa | ccccctagaa | cccaagtgcc | agaattccta | gatgctgctt | ctgtttgaac | 1920 |
| aaaatgtcac | tgctttacac | ttgaaaaaaa | cacactcgaa | aaatgttcaa | ctccatgaaa | 1980 |
| aataattttt | ggctttaaga | aattgtttgg | tgtttaactg | tttcctttga | ttgccattcc | 2040 |
| accagttaa | tggtggttga | tttgcaactgc | actctggggg | tgggggttgg | aggggagggt | 2100 |
| ccttatacag | agccgaacct | ggggttgctc | aggaaagtgg | ccagggaatg | tgggaagtcgt | 2160 |
| tgacattgcc | tgggccaaaa | gagtgggaga | tagtttttct | cccctcagcc | cactcctggg | 2220 |
| agcacctgtc | gccagcctgg | tacaaagcca | ggcctttttc | ttctgtgagc | atctcatcac | 2280 |
| tgtccagcag | cagggtgaaa | aaggggggaca | acaaccagac | ctattttttc | tccccatttt | 2340 |
| tttccaaatt | ttgctgtgcc | aaatgtttta | aattttataa | atatgaatct | attgaaattt | 2400 |
| ccttaatcaa | gagcttcttc | gtgtaaagtt | tgctttttta | gctatagaaa | aagaaaacag | 2460 |
| taaatatctc | ttaatggcat | ccagccttgc | tgagctcacc | ttttttcttg | aagaatgggt | 2520 |
| aggagtga | taattaatgt | aaacatttgc | ccaagtccct | ttacccta | tttgaagctg | 2580 |
| cattaaaacc | aactcactaa | cacagggaat | gattgcaccc | tagtctctct | tgggccaaga | 2640 |
| aactttcaga | agcattaaaa | aaatagttga | agtatatcac | ttctcaccaa | gtgggtagag | 2700 |
| tcagttggct | gtttgtccct | tgttttttat | ttattccata | attatgtttg | tgctttttgt | 2760 |
| tttgtaaaaca | gtaatggaac | gtacattttt | attttgttta | gaagacaact | ttgattcaat | 2820 |
| ctttcaagaa | ctgttccatt | cttgggtttct | tcttaggggg | ataaaaagtt | accagttaat | 2880 |

| | | | | | | |
|------------|-------------|-------------|------------|--------------|-------------|------|
| ttgttttgag | atattttaagc | attcttttgag | ttatagaatt | gtgatgcagg | gattttgtgaa | 2940 |
| tgagacattt | acatgtgaaa | ggtgacttca | ctagttacct | gcttgagcag | agtaaagtgt | 3000 |
| gtatatgtac | ataaaatgta | agtaatctta | actccattgt | gcagtgccct | ttagatgttc | 3060 |
| cgctttctat | aaagtcttca | aattttttgca | tatatattat | atatatatga | tgtaatgtta | 3120 |
| tagaaatata | tgtataatat | acatattttt | tccaggggta | tctgatatagct | ctgtattttg | 3180 |
| ttatggaagt | tgaaagaaaa | aagtattttt | cctcagaaat | taaataaaaa | aatactttta | 3240 |
| agtaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | | 3288 |

<210> 46
 <211> 1080
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (1078)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1079)
 <223> n equals a,t,g, or c

| | |
|-------------|---|
| <400> 46 | |
| gcgacgcgt | gggctagggg catagacttg tatttttatg ttttggggat gggatgggggt 60 |
| tatcttctgt | tgtaaattag gtgggatatg aacacactta ttttcttggg gagtaggaca 120 |
| cagggagggt | attgaactca aaggagaagg agcctctatt tcctgttgag gcttatactt 180 |
| tctgacaagg | aaactcccaa agggccaaga tacttttttg tcccatgata gtcagggtcc 240 |
| agacttcttt | cttttctatt gtatttgtat gtggaaagcc attaatgaat ttattttact 300 |
| ccaagagata | cacataaaaag ggaaaagggc atggttgagg ggttgagcag taagagacca 360 |
| ttaataactaa | gtatttttgcc tgaaatgctt ctttataatt atttcaggga aggaataaga 420 |
| taacttcaga | cttactgggg tggggcttgg ggaatgtttg gatgttgcyt ttttycttct 480 |
| tccttctttt | tttttttttt ttttgggtgg gttgtttggc atttttttct tttccttttt 540 |
| aaagaaaatg | ttttcaggaa acatgaacaa acatactgsc aattatagtg gggctgggaa 600 |
| ggcacaggag | ctggcgacct ctcagctgca tagttgggac ggcaaaccct gctgtgagct 660 |
| tctcagactt | ttcacctact tcacctatta agaagccatg aggagttgta aactcttggt 720 |
| caggggaagaa | aagagggcag gaggaattaa cccaatgcct ttaaatacaa aacagatgaa 780 |
| aaacaaaata | ctcatttttt cttttctatt gtgggttttg ggagccaaac caagtgtgac 840 |
| tgtgagcacg | ttcactgaga ggcattgaca cattattgaa gaggtctttc cattatgtct 900 |
| ggttctctta | ttttcttaag tcaagcctta actatgaaca tactttaaga tgatacgggt 960 |
| ctgcaccata | aacctgtaaa agggcatcga cagcttttca gataatccag gaatgttgag 1020 |
| acatggctaa | gtttctagct gatacttagt cattccctct tgatattggg gggatgggna 1080 |

<210> 47
 <211> 796
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (4)
 <223> n equals a,t,g, or c

| | |
|------------|---|
| <400> 47 | |
| ggantcctca | ggcggttcttt agaaggtccg cctgcaggta ccggtccgga attccccgggt 60 |
| cgaccacgc | gtycggcaga cacaggcact tattcattca tctcattgaa aagctacgag 120 |

35

| | | | | | | |
|-------------|-------------|------------|-------------|-------------|------------|-----|
| ttgggttcctt | attgcctctc | cataatagaa | aaactcttta | atgagctctc | tttttgtttt | 180 |
| tcaaatcaga | tatgcaaaga | agctcataac | aatttttttt | aaaaatgcaa | aacaagaatc | 240 |
| tccaattatg | ggagcaaaaat | cttcagcttc | tgggttcctg | tctcactgag | gaaatggatt | 300 |
| tgaaatggca | aggaggaaat | gaggaggcaa | actttcatgt | ctattttagt | tttccaatgc | 360 |
| agtcctattt | cctttggact | ttgtataaac | aaggaaagga | cagttgttag | ttcagttatt | 420 |
| acagataacc | tgtgtcttta | aagtaaatgt | atcttaaata | agtaggactc | ccataaatga | 480 |
| ctacactttt | tcaaaatatg | actccccagc | ttataacaag | aataatagca | aacatcactt | 540 |
| tattaagcaa | ttactatgta | aaagacactt | agtgccttagc | acacactgga | aatattgttg | 600 |
| actggctata | ttttccccag | aaatcccatt | tctgaaagcc | tattacaaag | aaataaaatc | 660 |
| atcagtataa | caaaggagtg | tgtgtgtgtg | tgggtgtgag | tgtgtgtggg | tgtgagtgtg | 720 |
| tggggggtgt | gagtgtgtgt | ggggtgtgag | agtgtgtgtg | tgagtgtaaag | tgtgtgtgtg | 780 |
| tgtaagtgca | cacacg | | | | | 796 |

<210> 48
 <211> 657
 <212> DNA
 <213> Homo sapiens

| | | | | | | |
|------------|------------|------------|-------------|-------------|------------|-----|
| <400> 48 | | | | | | |
| gctaacctgt | agataactat | agatgaaact | cacatthttcc | ttccccctggt | tcaccttgac | 60 |
| agcattgcag | ctctggtcag | ccacagaatg | ccaggcagtg | gtggacacca | tgatagccgt | 120 |
| gtggtcagaa | gggaagggca | cgggagtgag | ctgggaacct | tggctgttgg | ggaagttgca | 180 |
| aagctccagt | ttcctctaag | tgtgccttta | tgatccgcac | tggcsacagc | aggctttgtc | 240 |
| tttccttaga | aagactttgt | gcctgttctc | acttgaccag | atatcgtgta | cggagcactt | 300 |
| gctggcagta | ggccaagctc | tgacattca | tgaattgctt | tgtctcacia | caccctgagc | 360 |
| cgtaggatc | tttattgacc | atattttcca | gatgaagaga | cgaagccca | agagaaccct | 420 |
| gttaatgctc | ggccagatca | gaggagacaa | aacaagcacc | cctgccttca | tgggttgtgt | 480 |
| ggatgaatga | rggtcggcct | agggctgtcc | cctcgcggtc | tgtccggctc | gtcggtagac | 540 |
| gcascagatg | ggccacttgt | tcctggaatt | gcagaagcag | ataaccaagt | ktagaaaatg | 600 |
| tatgtatccc | tcttctaccc | caacagaagc | accgtctttw | tgtcttttct | tttttcc | 657 |

<210> 49
 <211> 1082
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (6)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (22)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (30)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (49)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (53)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (120)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (125)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1027)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1060)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1067)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1070)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1079)
 <223> n equals a,t,g, or c

<400> 49

| | | | | | | |
|-------------|-------------|-------------|-------------|-------------|-------------|------|
| cccanaggg | aggaaaaacc | cntcagttan | gtagaatgcc | caaatcagnc | ctnaagctga | 60 |
| agtctcaagg | gtgccaacga | gtggcctgcc | aaaagaaaga | gtgccaggaa | gggtagggtgn | 120 |
| ccagncagag | cctgaggagg | aataaccagt | gctgaggata | gccaaggggc | tcccggggggg | 180 |
| tctgagtcct | tgtcacacca | ggaagtgtctg | tgaccaaagc | tacagggggcc | tggctaagac | 240 |
| agcttactct | atTTTTtctt | gaggctttgc | agaattctca | gaagcactgc | cagaaaatta | 300 |
| aaaacagaca | cttacagcaa | agacctaatg | gttgaatggg | taacagggttc | ctctggaagt | 360 |
| actctttaca | aattgttcaa | aattataatg | gctcattata | ttaggaatct | acagaaacag | 420 |
| tctggacaaa | acaaaaaaac | ccatcccaag | tgcattctgga | caggcttcaa | ggacccccaa | 480 |
| cagccatcct | gcagctctga | ctccaggagc | tggacgggaa | ctgcggcctg | gacctcccag | 540 |
| caagcctctc | tgcgggcagg | cgggccacag | cagccgaagg | cggagctttg | cttccagatg | 600 |
| aggggtgcaca | gataccttta | acttaacggg | gttaaaactga | caccagtcaa | tctgacacca | 660 |
| ctgggcagct | cagggtgttg | gaagacagag | ttccacctgg | ggcagtgtct | tgtaaatgatg | 720 |
| gaaagtccaa | tgtaaaagtgt | tcattaaatg | agaaatactt | cctctaaaat | agctctgtaa | 780 |
| aatcaatata | gatacctcaat | caataatTTT | tttggtattt | cccacagaaa | aaaactacaa | 840 |
| aatggtaaagc | taaaatcagt | gtggcattta | ctaagtaagc | tttggccttc | taaacatcaa | 900 |
| aatcaattcc | atattctggg | atataacaca | tagccacaat | ggtgtgcaag | taattccagg | 960 |
| atgtgagaca | cacagacacc | gtgagtctgt | caatagtgtg | ttctcgagat | aggttcttca | 1020 |

gcaaggntcc gttttctcaa atcaaaagag ggaagatgtn caaaatnacn gacgttgng 1080
gt 1082

<210> 50
<211> 690
<212> DNA
<213> Homo sapiens

<220>
<221> SITE
<222> (36)
<223> n equals a,t,g, or c

<220>
<221> SITE
<222> (604)
<223> n equals a,t,g, or c

<220>
<221> SITE
<222> (630)
<223> n equals a,t,g, or c

<400> 50
ggcgaacaat cgcttcatct ccctttggtt agggcnccac caaccaagtg tagaaaatgt 60
atgtatccct cttctaccyt wcataagcac cgtctttttg ctttttcttt ttttttttag 120
ttacagatat gataggtagg tttttttcta tgggtgaagc atttctgctg ttggattttg 180
tttgctgttt gaagacctgc ctaactcgtc tcagtaggct gtataaatcg tttaatgaag 240
attggccctt ttctgaagtg agaaaaagga tgggtgctga agaacctatc tcgagaatac 300
actattgaca gactcacggt gtctgtgtgt ctcacatcct ggaattactt gcacaccatt 360
gtggctatgt gttatatccc agaatatgga attgattttg atgttttagaa ggccaaagct 420
tacttagtaa atgccacact gatttttagct taccattttg tagttttttt ctgtgggaaa 480
taacaaaaaa attattgatt gaggatctat attgatttta cagagctatt ttagaggaag 540
tattyctcat ttaatgaaca cttacattg gactttccat cttacaaga cactgsccca 600
ggtnggaact ctgtcttcac aacacctgan ctgcccagc ggtgtcaaga ttgactggtg 660
tcagtttaac cccgttaaag tttaaaagga 690

<210> 51
<211> 1618
<212> DNA
<213> Homo sapiens

<220>
<221> SITE
<222> (644)
<223> n equals a,t,g, or c

<220>
<221> SITE
<222> (647)
<223> n equals a,t,g, or c

<220>
<221> SITE
<222> (656)
<223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1439)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1468)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1500)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1527)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1617)
 <223> n equals a,t,g, or c

<400> 51
 gcggccgccc tttttttttt ttttttgaga tggagtctcg ctctgtcgcc caggctggag 60
 tgcagtggca cgatcttagc tcagcagtg agcctctgcc tcccagggtc aagcgattct 120
 cctgcctcag cctcttgagt agctgggatt acaggcgtgt gccaccacgc ccagctaatt 180
 ttttgtatth ttagtagaga cggggtttca ccatgttggt caggctgata tcgaactcct 240
 gacctgtga tccactggcc tcagcctccc aaagtgccgg gattacagga gtgagcaccg 300
 caaccggcca ttattacaat ttaacccct taatttcagc taccttctcc actgaatact 360
 acacaccatt cagtaactgg aatactgaaa gaataattagt cttgaatgtt catgtcccc 420
 caaaattcat aggtgaagc tctcactcta cggtaatggt gttaggaggc agggcctctg 480
 ggaggtgatt aggtcatgag agtggacaag cctcaagaat ggggttataa aagagacccc 540
 agagagctcc cctgcctctt ccaccacata aggatgcagt gagaagggtg tggctatgga 600
 ccaggaagcg gggccccacc agacatgcaa tctgccagtg cctngancct ggaccncag 660
 cctccagagc tgtgagcaac agactcctgc tgttgatgag ccacactcta tgggtgtttg 720
 tcatggcagc cccagtgggc taagatacac tagaagttca atcatggatc tctgtatctc 780
 atcaccaatg tgcactcagc ttgcctgtgc tgctttccct tcccacggcc ccgcctgcct 840
 ctggccgctc gccactccct ctccagcctc ccttcacct ctccctcctt cactgccaga 900
 caccttcaag tccctgccgg cagaaacaca cccaggcctt aagtgcctg ggggtagggtg 960
 gcgggacaaa agggcccaact tcacaaaggc cagacactga acatactccc acaggggggc 1020
 tcctggagat ggtgactggc ctggttaagag tcactaggac tattttttta aaattaaaag 1080
 agatggcacc tcaactatact cctaggctgg tcttgaacac gtggtcttaa gtaatcctcc 1140
 tcagtctccc aaagtgttag gattacaggt atgagccagt gtgcccagcc tgtggcaaaag 1200
 tgcagtggga caggaaaatg gctcaggttt aacttcagc ctcaatgaaa actaatatga 1260
 actctcatcc atgcaccaa ttaggaacac tcttacctg actcttagaa tgggkatttt 1320
 tagaaaaag cccaaaaagt agtttaggac aaagaacca gaatatgggc tcctagagca 1380
 agkgtctgg tgkccctyag gtcattgaaa gctgagaatc atcaatgaca ataaactgnt 1440
 ttatgggtta cgaaacactc ttcacatnca ctctttactg aaaatcttcc caaacctgnt 1500
 ggggtgcaaga cattgggtccc cttttgnaca ggattaaaga aaaatgggct ttggccaaaa 1560
 ggggaccagg aaccatggga cctttcctaa caattttaaa actcttggga tcccaanc 1618

<210> 52
 <211> 570

<212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (5)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (16)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (496)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (523)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (531)
 <223> n equals a,t,g, or c

| | |
|--|-----|
| <400> 52 | |
| ttccngtgcc actatnga | 60 |
| atgtacgcctgc aggtaccggt ccggaattcc cgggtcgacc | |
| cacgcgtccg ggcagtgggg tgagggcaca caagcagttc aggggccag caggaagtgg | 120 |
| ggctgcaggg ccgggggtggg tcctgggcct ggccatcagg cagcctagca ggttgttctg | 180 |
| ggcatggagg gggcctgggtg tggctgaggg catgcccagg gctccctgga ggatcccgt | 240 |
| ctgtgccctg cccaccctgt gcctggggag ccctctgccc tcacagccca cccaccccat | 300 |
| ttwctatgac cacagagctc cgacctggaa gatggctcac ccaggaggtc ccaggagctc | 360 |
| tcaactcccc aggacctgga rgacaccag ctctcagaca aargctgcct tgccggcggg | 420 |
| gggagcccga aacagccctt tgcagctctg caccaggagc aggttttgcg gaaccccat | 480 |
| gtaaggcttc cccgngtggt ggtcctccag ccgtgggcct canggtgacc natcacaggg | 540 |
| agagtggctc cctgtcctgg gcacccctg | 570 |

<210> 53
 <211> 355
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (10)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (19)
 <223> n equals a,t,g, or c

| | |
|---|----|
| <400> 53 | |
| acactatatn tagggacanc tgcccgtacc ggtccggaat tcccgggtcg acccacgcgt | 60 |

| | | | | | | |
|------------|------------|------------|------------|------------|------------|-----|
| ccgtactgcc | ctttttaa | ctcagatgtg | actttcatta | taggaagttc | tcaggcattt | 120 |
| tctcttgga | taatacctct | tctctcttct | ctttatgtcc | ttgtgccgca | ttctgggtta | 180 |
| ttcctttagc | tctaggttaa | gttcaactaa | tcttccttta | gctgtatttc | attattgttt | 240 |
| aagctgtcca | ttgcatttta | aactttcttt | caaatatctt | cccttccctt | cctttccctt | 300 |
| ctcttccctg | ccctgccctg | ccctgccctg | ccctgccctc | ccgtccctc | ccctc | 355 |

<210> 54

<211> 1149

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (146)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (147)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (148)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (149)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (927)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (1077)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (1083)

<223> n equals a,t,g, or c

<400> 54

| | | | | | | |
|------------|------------|-------------|-------------|------------|------------|-----|
| aaggtggatt | tccttccaac | tatctcagaa | agaagccgta | gccgactccc | tttctgccta | 60 |
| gagggcccg | tggtccttgc | tggtctttacc | cacgtattta | atagcctctt | taatttatca | 120 |
| ttctccacgc | agcttccaag | gcagannnnc | tctgggggtac | tgtgacttct | tgcttcccat | 180 |
| gggtggagag | tgagtttctg | acacctcgga | cgccccctgg | cctgatccct | cctatcacag | 240 |
| catcactgga | ggctcagaac | ccacagcctt | tgctttctgt | ccatgtcagc | atccctgcct | 300 |
| taagagaact | cctcctggcc | aatggcattg | ccaccagca | gtgggaccaa | gactctccaa | 360 |
| gacctccagg | actggatccc | attgcctgga | gaaactccag | caaggggtct | tcattggctt | 420 |
| gacatggcac | agtaaggggc | agccaaccca | gtccatgatg | acttttgctc | caacttcttc | 480 |
| atgtttctaa | aagcccagtg | gctttattca | ctcctcctaa | attgcctgct | accagaagga | 540 |
| acttcacct | gaagaaatgc | attccattac | cagttccagg | gaaagtgtcc | ccttccccaa | 600 |

| | | | | | | |
|------------|------------|------------|-------------|------------|------------|------|
| agcctcaggc | ccgtgggct | ctgagggtcc | actggatgcg | gctccccag | gagtgggct | 660 |
| ggagatccgc | tcagccgccc | tgccctccac | cttctatctt | gggaccgtgg | tcagccctga | 720 |
| agggtgggtc | cagcccgcg | tatgctgcac | tttgctgctg | caggctccgg | tcctycagg | 780 |
| gcttttcaat | gagagtcccc | accccaactt | ggagcatttc | attttgsct | acctwaagca | 840 |
| agaatctcaa | agtttggttg | aataaraagg | ccattcacaa | gtcgtgccct | tgtgaagcac | 900 |
| cccttttgyt | gaatgsctcg | tgggtgncca | aacactgkac | ttggcgcttt | gcattcatgc | 960 |
| tttgaccctc | acaacctccc | tttatagatg | aggaaattca | cgctccatgt | gccaaggac | 1020 |
| acaccagcca | ggaggaagag | ggaagaggat | ttgagtccag | gcctgcctga | cttcagngcc | 1080 |
| tgngccctcc | cctctgtaaa | agacaaacca | ggcaggggaag | agatagaagc | tggctaagtg | 1140 |
| gtggccaag | | | | | | 1149 |

<210> 55

<211> 1767

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (8)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (11)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (12)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (34)

<223> n equals a,t,g, or c

<400> 55

| | | | | | | |
|-------------|-------------|------------|-------------|------------|------------|------|
| aggatccnac | nncagggtga | acctttccgg | gaanctaccc | gaccgaggca | gccctgccct | 60 |
| gccctgcccc | gcccctcctc | accctgcyct | gccccagaga | tctccagttc | acagtgttca | 120 |
| tcaggcgaga | tacttttgga | cccacaaagc | cagttttttg | gggggatgct | cagtgtctac | 180 |
| ctgcagtgcc | aaagttttgc | ttgaacagcc | ctcttctgcc | gcgcctccca | gtccaccccc | 240 |
| ctatcaacgt | gggtgctggc | tgcttctgcg | tgaggagctt | cctctgatca | cctgtagcat | 300 |
| ctttagtgcc | ttcgtccagt | cggcctgcaa | atcccatttc | ccagagaggg | ttgggcaggg | 360 |
| cccacttgcc | acttgcccct | gtgcccagag | cgttgggggc | agatggcagc | caggggaaga | 420 |
| agggaggtgg | ggaaggggcg | ccaccgcttg | aggggtcccct | cccttgagag | gaggccccat | 480 |
| ggcagttttc | ctgagttctt | gatccacttt | ggttttaaca | acttttgtga | agctatgtga | 540 |
| gatttgactg | cattttcaaaa | gacaaaacac | atgttttttt | ctgatttgct | ctttgccttt | 600 |
| cagcgacgct | ttcagatact | ttgtgagtgt | ctactctgtg | ctaggtgctg | agtagactgg | 660 |
| gatgtgtgtg | gtcccatccc | tggttggaag | agctttgagt | cctgtctgct | gacttcactg | 720 |
| aaggagtttc | tctcttgctt | cattccttcc | tcccgggggt | cagctgggta | ggagcagact | 780 |
| tgccctgcct | tcctctaggg | agaaaagttt | cctgcaaaatg | aaagaatctg | ccattctctt | 840 |
| ttgctatgat | gatgatgatt | attctgacat | gaccgcgggg | tactgaagcc | acagtcctct | 900 |
| ctgttctactg | ttctctcctt | gagtcatatc | tatggcctga | ggacgtctgt | atccagctcc | 960 |
| ttttcctaca | atggtagaca | cttccccagg | cagtgggtgg | gtcgsggtgs | ccagtcactc | 1020 |
| tcctagctgt | ccttgccctg | ggacttagat | ttaccccatc | caggactgtg | gtcgctattc | 1080 |
| tgtgagcgat | tgatggcgca | tatgcagaac | tcccctggag | tcatggctct | agaaagtgtc | 1140 |
| caggccccatc | gtgacaatgg | tgccaagcac | cgggctggtg | accagtgcct | aaacacacac | 1200 |

| | | | | | | |
|------------|------------|-------------|------------|------------|------------|------|
| gctcccttaa | tccttacggc | aagcctgtga | ggcggatagg | accagccctg | atctatagtg | 1260 |
| agaaaattaa | ggttcagagg | gagaagggtcc | aaatagtggc | caagctggga | ctgagatctc | 1320 |
| gagtttggtt | ttgtttgggt | tttgttgttg | ttgttctttc | cctcgtgtcc | atgacttttc | 1380 |
| tctgccatta | gcarggaagt | gtgcggaggg | cttccagggg | aatcaagtga | ccctccctct | 1440 |
| gaatttcggt | gcctcttccc | cccgcctctt | gctcacacat | ttcagtaggt | cacaggccaa | 1500 |
| atggcagccc | aaaaggtggg | cagagatttt | tttttcttcc | cctcaccctt | rcttattaac | 1560 |
| acaattgtga | caactacttt | accttacatc | ccagggcaaa | cggacagctt | gcggtacctc | 1620 |
| caaaaatttg | gaaaaccctg | agtctaagac | cacctcttga | taaaagaggc | ctttgctcac | 1680 |
| atgttctgct | cggaatcttt | ctttgccatt | ggtgggtgta | tgtcgccag | ctgagactkc | 1740 |
| gggagagytg | agttcagctt | aaaccgc | | | | 1767 |

<210> 56

<211> 555

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (545)

<223> n equals a,t,g, or c

<400> 56

| | | | | | | |
|-------------|-------------|------------|------------|------------|------------|-----|
| tccttgettc | taagctgttg | ccccctgggg | aatagagctt | atggagcaac | gggagctgag | 60 |
| gtggcctcga | gggccagcct | ggagggctct | gagcatagca | tgcagcggtc | ccatagggag | 120 |
| gcaggttaacc | aagggccagg | cagagctgct | tcctgtgcct | ctccggcatt | tgtaatgktt | 180 |
| ttttcttttt | tcacgcactg | ccaaatctgt | tttctccctt | gaagaaaacc | agggcagatg | 240 |
| tggctcgtct | gtcaaactga | ggtctggaga | ggcatgacta | aatcactgca | gaagcaaaat | 300 |
| caggacccag | cattttctagc | tcccagccca | gagtaagaga | gcagagcaga | accccttccc | 360 |
| attctaccog | accgaggcag | cctgcctgcc | tgcccgccct | ccccaccctg | ccctgcccag | 420 |
| agatctcagt | tacagtgtca | tcaggcgaga | tctttggacc | cacaaagcag | ktttttgggg | 480 |
| ggatgctcag | tgtacactgc | agtgccaaag | ttttgcttga | acagcccttt | tttgccggct | 540 |
| ccagntcacc | cctat | | | | | 555 |

<210> 57

<211> 2694

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (258)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (925)

<223> n equals a,t,g, or c

<400> 57

| | | | | | | |
|------------|------------|------------|------------|------------|------------|-----|
| ggcgccgcgc | cgagcgaatc | tcggagtcgg | tgggtgcaga | tggcgccggc | agttgtggtg | 60 |
| gcgagggggg | acagcgactc | ccggccccga | caggagtgtg | tagtggcctg | gaacaccgtg | 120 |
| agcaccggcc | tgggtgccgc | ggctgcgctg | gggctggtgt | cttcccggac | cagcggtgca | 180 |
| gtcccgccaa | aggaagagga | gctccgggcg | gcggtggagg | ttctgarggg | ccacgggcta | 240 |
| cactcgggtc | tggargantg | gttcstggag | ktgctgcaga | acgatctgca | rgccaacatc | 300 |
| tyccctgart | tctggaatgc | catctcccaa | tgcgagaact | ctgcggatga | gccccagtgc | 360 |
| cttttgctac | tccttgacgc | ttttggcctg | ctggagagcc | gcctggatcc | ctacctgcgt | 420 |

| | | | | | | |
|------------|------------|------------|-------------|-------------|-------------|------|
| acctagagct | gctggagaaa | tggactcgcc | tgggcttgct | gatgggcact | ggtgctcagg | 480 |
| ggctgcgaga | agaagtccac | actatgttgc | gcggagtctt | gttctttagc | acccccagaa | 540 |
| ccttccaaga | gatgatccag | cgtctgtatg | ggtgcttctt | gagagtctat | atgcagagta | 600 |
| agaggaaggg | ggaagggggc | acagaccggg | aactggaagg | ggagctggac | agccggtatg | 660 |
| cccgtcgccg | gtactaccgg | ctcctgcaga | gcccgtgtg | tgcaggggtg | agcagtgaca | 720 |
| agcaacagt | ctgggtgtcg | caggctctgg | agcagttcca | tcagctcagc | caggctcttac | 780 |
| acaggctcag | tctgctggag | cgggtcagtg | ccgaggctgt | gaccaccacc | ctgcaccagg | 840 |
| tgacccgga | gaggatggag | gaccgttgcc | ggggcgagta | cgagcgctcc | ttcctgcgtg | 900 |
| agtwwcacia | gtggatcgag | cggngggtt | gctggctcgg | caagggtgtt | ctgcaggacg | 960 |
| gccccgccag | gcccgcac | cccaggccg | gcaacacct | gcgccgctgg | cgctgcacgt | 1020 |
| gcaaaaggtt | ttctaccgca | tctacgccag | cctgcgcac | gaggagctct | tcagcatcgt | 1080 |
| ccgagacttc | ccagactccc | ggccagccat | cgaggacctc | aagtactgcc | tggagaggac | 1140 |
| ggacyagagg | cagcagctgc | tcgtgtccct | caaggctgcc | ctggagactc | ggctcctgca | 1200 |
| tccaggcgct | aacacgtgtg | acatcatcac | cctctatata | tctgccatca | aggcgctgcg | 1260 |
| cgtgctggac | ccttccatgg | tcctcctgga | ggtggcctgt | gagcctatcc | gccgctacct | 1320 |
| gaggacgcgg | gaggacacag | tgcggcagat | tgtggctggg | ctgacggggg | actcggacgg | 1380 |
| gacaggggac | ctggctgttg | agctgtccaa | gaccgaccgg | gcgagcctgg | agacaggcca | 1440 |
| ggacagtga | gatgactcag | gcgagccaga | ggactgggtc | ccggaccctg | tggatgccga | 1500 |
| tccagggaag | tcgagctcca | agcggcgctt | atcggacatc | atcagcctgc | tggtcagcat | 1560 |
| ctacggcagc | aaggacctct | tcctcaatga | gtaccgctcg | ctgctggccg | accgcctgct | 1620 |
| gcaccagttc | agcttcagcc | ccgagcggga | gatccgcaac | gtggagctgc | tgaagctgcg | 1680 |
| ctttggcgag | gccccaatgc | acttctgtga | agtcagtctg | aaggacatgg | cggactcccg | 1740 |
| ccgcatcaat | gccaacatcc | gggaggagga | tgagaagcgg | ccagcagagg | agcagccacc | 1800 |
| gttcgggggt | tacgctgtca | tcctgtccag | tgagttctgg | ccgcccttca | aggacgagaa | 1860 |
| gctggagggt | cccgaggata | tcagggcagc | cctggagggt | tactgcaaga | agtatgagca | 1920 |
| gctcaaggcc | atgcggaccc | tcagttggaa | gcacacctg | ggcctggtga | ccatggacgt | 1980 |
| ggagctggcc | gaccgcacgc | tgtctgtggc | ggtcacccca | gtacaggcgg | tgatcttgct | 2040 |
| gtattttcag | gaccaagcca | gctggaccct | ggaggaaactg | agcaaggcgg | tgaagatgcc | 2100 |
| cgtggcgctg | ctgcggcggc | ggatgtccgt | gtggctgcag | cagggtgtgc | tgcgtgagga | 2160 |
| gccccccggc | accttctctg | tcattgagga | ggagcggcct | caggaccggg | acaacatggt | 2220 |
| gctcattgac | agtgcgacg | agagcgactc | cggcatggcc | tcccaggccg | accagaagga | 2280 |
| ggaggagctg | ctgctcttct | ggacgtacat | ccaggccatg | ctgaccaacc | tggagagcct | 2340 |
| ctcactggat | cgtatctaca | acatgctccg | catgtttgtg | gtgactgggc | ctgcactggc | 2400 |
| cgagattgac | ctgcaggagc | tgcagggtca | cctgcagaag | aagggtgcggg | accagcagct | 2460 |
| cgtctactcg | gcccggcgtc | accgcctgcc | caagaactgc | agctgacaca | tcgcccggcc | 2520 |
| gccccccgc | ccgccaaggc | ctgccctgca | ggtgctctcg | tcctcccgtg | ccagcccccg | 2580 |
| cccgcctg | tcccagaatg | cactgctgag | gagcatgccc | acccccaccc | ccgcagtgtg | 2640 |
| cagattaaag | caagtcagat | catcaaaaaa | aaaaaaaaaa | aaaaggcgcg | ccgc | 2694 |

<210> 58
 <211> 5811
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (2809)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (2827)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (5787)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (5803)

<223> n equals a,t,g, or c

<400> 58

| | | | | | | |
|-------------|------------|-------------|-------------|-------------|-------------|------|
| aaatTTTTTT | TTTTTTTTTT | TTTTTTTTTT | TTTTTTTTTT | TTTTTTTTTca | ccascacggc | 60 |
| ttctctccag | ccttctcttt | attttctcca | gggcaccagc | ccctctgcc | gcctgagtc | 120 |
| agggctccta | ggcactctac | cttgcccttg | ccattcccag | ggcctccccg | cggccccgca | 180 |
| ggggagggag | gctcagagca | cagtctcggc | agtgttaggg | tcacgagga | actggctgac | 240 |
| gcgtttgggg | tgctgcaggc | ctggcgcggg | gtccactccc | actcctgcct | ccttctgtct | 300 |
| ttcgggtgtcc | tcgtccttcc | cctgggtcctt | ggagaagtac | aagaataacct | cctccagcat | 360 |
| cgtctggctc | acggaaaagt | cctccacgcc | gtgctctgcg | ccgtgcaccg | ccagctctcc | 420 |
| aaagacgcgc | gccaggggcg | agcgccctcc | cggcggcagc | tggaaagcga | ggcggcctcc | 480 |
| atgtgcctcg | cgcagctccg | mcccaggga | ctcggccgcc | acgaaggccg | ctgccggctg | 540 |
| ggaccttgcg | gcggggcacc | gcagggtcag | tgtgtgaccc | gccgcgaatc | tgcccttgag | 600 |
| atgttgcggg | ctgcccaggc | agcggaaccg | cccattcacc | atgatggcca | ggcgcgagca | 660 |
| gagcgcttca | cactcctcca | tgctatggga | ggtgagcatc | actgaacggc | cctcccgcac | 720 |
| cacggccaaa | aggctgttcc | aaaggaagcg | ccgcgcgctg | gggtccatgc | ctgtggtcgg | 780 |
| ctcgtccaga | aacaccacgg | ctgggtcccc | aaccagcgcc | agggccgctg | ccagcttgcg | 840 |
| tttgttccct | ccgctgtagg | tgcttgacgg | ccggtctgcg | taccatgaga | gtcccagacg | 900 |
| cgccaggccc | gagccagcgg | tctgggcaac | ctgggcctcc | gggacaccgc | gcaggcgcg | 960 |
| aagcagctcc | agggtgctcg | ggcccgctag | cagctcaaa | atggcatcgg | attgagggca | 1020 |
| gtatccccatg | ctgaggtgcg | cagcactggg | ttcccgggcc | acgctgtggc | ctgccagcac | 1080 |
| agcctcgccc | ctgctggcca | atgtgtcccc | cgtcaccatg | cgaaacgtgg | acgtcttccc | 1140 |
| tgctccattc | acaccagca | gccccaaa | ctcaccaggg | ggaatcccc | ggcacaagcg | 1200 |
| gtcaacagct | ggcactcctt | gccccaggta | taccttggtc | aagttcctca | gcaccaacac | 1260 |
| atccccctgg | gtggctcctt | ggaccacccg | ctcccggtca | cgggctacat | cctcgtcctc | 1320 |
| ctctcccagg | agtggcagag | acctcaccct | gggctgtggc | aggagttggc | ttcgggtgctg | 1380 |
| cagcagtagt | gtgaagagaa | ggaagagggg | cccctgtatc | accatggcca | agaggttctt | 1440 |
| gccgaccacc | tcccagcgca | ggggtgactg | gaactgectg | tctcccagc | gctcaaaggc | 1500 |
| atcagccatg | gcctggttcc | gcaccatgtc | aatgagcccc | cggccccagc | agaagtgggg | 1560 |
| gaagataaag | aagacctggt | tcaagatccg | gctcacctcc | tgagcttctt | gatcagagaa | 1620 |
| gagctcaagc | acaaagggtg | ccatgcttcc | attgatgcca | ataaagaggt | ttatgcaggt | 1680 |
| gagcaccaca | taggctgtgc | tgggcacgga | gaagaagaag | gaggctgggt | acatgagcgg | 1740 |
| tgtgatcgac | cagccataca | gtagtagcaa | cagcaggaga | gcaggcaggt | tggcaggggc | 1800 |
| cacatatgcc | ctctgctgga | aggccagaaa | gatgagcacc | acgatgcatg | ctggcaccac | 1860 |
| gtagttagac | atgtcccaga | gaaagtgtgc | aagccagtag | aggggtgggg | acaggccccc | 1920 |
| catgagctgc | agggtgcttg | ctcgggtgac | tcgctcctca | atgaggacaa | gagtgaagct | 1980 |
| ggccgggaca | aaggacatgg | caaagaccac | acagatggag | acgaggacgt | ccaccgagga | 2040 |
| ggccatcagt | gcaccctcag | acagctgtct | ccttggtgag | ttcaaggggt | ggttgagtgt | 2100 |
| gggtgatgctg | tgggcgtggc | gggcccggcc | tgggggcagg | tgagcacgga | ggattgctgt | 2160 |
| gctggctcgg | ttgacaaagg | ccaccatgga | gtgccagcct | ttgttggtga | accagatctt | 2220 |
| gagactgtcc | tgagcatcca | ggctgtgagc | ccaggctgtg | aggtttttca | ggacacggtc | 2280 |
| gagggccccc | ccaggcaggg | gactcagcag | cggccacaac | tcctccactg | agcggcccaa | 2340 |
| ctcttgggcc | gagggcaggg | ctgggtctcg | gccccccagc | gagaagcctc | cgtatctgac | 2400 |
| ctcattcacc | cacttcttag | tcttcaggcc | ctggcgccac | agggcggggt | aggtcttgac | 2460 |
| caggaagtca | gacagggttc | ggcctgtcag | gttctgaacc | acttccccag | agccggtcac | 2520 |
| tgccctgggg | ggagggggac | caccagctgc | agccggcgag | tcgggcagca | ggcgccgggc | 2580 |
| accggggccg | ctacactggc | aggctgggga | tggagactct | gggggtccagt | tgccactggc | 2640 |
| caagaccttg | gccacttcag | caggaacttc | tgggtgccgag | aacctgtggg | agctatgctg | 2700 |
| cactgggggc | tcctccagtc | ctgcctcctg | cagcagcgcc | tcgagcagcc | gggcacgtcc | 2760 |
| agggctccct | ggggcgctct | cactggggca | acaggacgcg | gcgccccenc | cccccccccc | 2820 |
| ccccccnccg | cttgagaagc | agggcctgga | gctgctggcg | ggtcagtgcc | cagccctgta | 2880 |
| cccgccccac | ggcgtctggc | ccagagccct | ggtcagctct | tggggctgac | ccagctggtt | 2940 |
| ccccgttctc | cagcgtgtgc | tcctgtggcg | gcactcttag | ccgtakgggt | acgtctaggc | 3000 |

| | | | | | | |
|-------------|-------------|-------------|-------------|------------|-------------|------|
| cagcaatgcc | tgtgcatagg | tgtgccccgc | agctgccatc | ctccatatct | gtgtccgcag | 3060 |
| cacactcctc | caccaccttc | aggaagatct | cctcgaggct | ggtgtcggag | atcccgtagc | 3120 |
| cagtgcagct | cagctccgcc | agccgcgtgt | ctagctctcg | gaagagtgtg | gcgaagctgc | 3180 |
| cgtcatgggc | acccgtgtag | ggcagcacca | gcaccagctc | gtgtggcagc | tcctccacca | 3240 |
| gccgtgcccc | gggcacccag | tgtgtacca | gggccagcag | ctgaggagtg | ccgactctgc | 3300 |
| tgccctggct | gccattcttc | ttttcctgcc | tgggtgtccac | actgccctcc | atgtcagtgt | 3360 |
| cagccttctc | attggtggtc | aggggcaggg | gggccttcac | cagcgtcagg | tagtagccgg | 3420 |
| agcccagggt | acggcgcagg | aagagtgggg | agccacagca | gcacaagcgg | ccacctgcca | 3480 |
| ccacggccac | acggctctccc | agcagctctg | cctcatccag | gtggtgggtg | gagaggatca | 3540 |
| gcgtgcgacc | ttctcggtat | ttgagcagca | gctcccaaat | accgcggcgg | gaagcaggat | 3600 |
| ccacgccagc | cgtaggctcg | tccaggataa | caacttgggm | gmcgcccaca | aaggcaatgg | 3660 |
| ccacggacag | cttccggttc | atcccaccag | agaggtggcg | agtctgcaca | ctctgcttgg | 3720 |
| agaccagccc | cacatcctgc | agcagacggt | cctgctcggg | gcccactaca | gcggcactca | 3780 |
| gaccttccag | cgcgccatag | aaccagacgt | gctcgtccac | ggtcagcatg | tcaaacagca | 3840 |
| cgttgtactg | aggacagacg | cccagggtggg | gccggatggc | ggccatgctg | gagcggacgt | 3900 |
| cgtggcccag | gatgaaggca | gagccaccac | tgggtgggaa | gaggccactc | aagatggaca | 3960 |
| gggtgggtggt | cttgccggcc | ccgttgtggc | ccaggaaggc | ggtgatgtgg | ccctggtaga | 4020 |
| agtccaggct | gagccccgcg | agggtctggt | gcgggcttcc | aggaaagcgc | ttctccaggc | 4080 |
| tgcgaacgga | gacgccagga | ctcaggcccg | gcgggtgcctc | ttctaccagc | acctttgggt | 4140 |
| ccagcgggggt | ggggcaaggg | gctggactct | tggggggccg | aggtccgcac | cagtagctcc | 4200 |
| tccgaaaagg | aaaattccat | ggttcaggga | tcccgtagtg | gcctgggcac | acagcttcca | 4260 |
| ggtaccagggt | ggcgaggccg | tagagcgccg | cgtccagcag | cagaaggcca | gagacctggg | 4320 |
| ccaggctgaa | cagctctgcc | gtaggccggg | tgccacggtt | gtgccactgc | gcgccctcgc | 4380 |
| cctgytcttc | cagcagagcc | aggctcttcg | cagccgaagc | cgaaaggcca | cgggcgacag | 4440 |
| caggctcgcg | gccacgcggc | cacccgcggg | cagccggtcc | cgccaagcca | cacacagcac | 4500 |
| gtagggcagg | tagagggaga | agtagccag | gccgccgcag | gccgcagcca | ggttggcgcg | 4560 |
| ggagaagaag | gcgctgagca | ggaagctctg | ggtcaccgtg | gccaccgcga | aggctgccaa | 4620 |
| gaacaggaag | accacgcccc | ggtggctgta | ggggaggatg | tcycccagct | tgagcaccag | 4680 |
| aaccagcagy | gcggcgctga | gcaggaaggg | cccagggcag | ctgaggaacc | agcctagcca | 4740 |
| gagcaccgcg | cggctgagcc | ccatggcgcg | catgggtgtcc | cgcagccgcg | tctccttctc | 4800 |
| ccgcaccacg | gccttcaactg | tcagtgtcac | ggagtagatc | caggccagcg | tcaggaagag | 4860 |
| cggcagcgac | cggctcagca | cacgcaggaa | cacgtcgtcc | acatagcacg | gatagggcat | 4920 |
| ctgctgcagg | tagaggccgg | cccgggggtt | ggcgccgctg | agcacgcgga | cggctgcacg | 4980 |
| ctccaccagg | tcttgacaggt | acacgaagcc | gccccacacg | tagcgcaggt | cggctcagggg | 5040 |
| gtccgcggct | gggccagggt | cccaaaacct | gtccctgate | ttattgggtc | tcgtgaccac | 5100 |
| gtcaatgtcc | atgcggtatt | tgatgcgcac | gtgccggggc | ccaggtctgg | ggttgggtgc | 5160 |
| tctgtgggggt | ctgaagagtc | ctcaggtccc | agaagacga | cgccggccca | gaatcgatgt | 5220 |
| tccgcgagca | gttgacaggc | ccgcgacacc | agggtctgct | ctgagggtgc | cgctccagc | 5280 |
| ttgtccaagg | acaggcactc | cgtcactcgg | cccagcgtgc | ccaccagggt | ccccacatca | 5340 |
| gcgtgtgcgt | cctgccagct | gtagccaccg | ctcccagggt | ccagaaagga | tcgcagggcc | 5400 |
| tccatgtgggt | cccggcctcc | aggctctggg | tgccttcttc | cttcacctcg | catctgcagg | 5460 |
| agccgctgca | gcatggccmc | attggaactg | tcgttcatra | aggtgaagaw | ccgggggtccc | 5520 |
| agcatctccc | acacctcccc | gacatcccy | agcaggggtga | gctcctcgaa | ggtccgggtc | 5580 |
| acctggggcca | tgagcttccg | ggtaaaaggt | gtatctgggtg | caaagagtag | cttcccagg | 5640 |
| atcagaggct | tcaggcgtct | ccagagcagg | cgggacagcg | ggtggctgtc | cagggtcca | 5700 |
| atcagctccg | agcagggcgg | gctcaggctg | ctgtctggca | gggcggattc | tggctcctgc | 5760 |
| cccaccagct | ccatcaggtc | actagntcgt | taccagtga | gtnttccttt | c | 5811 |

<210> 59

<211> 1012

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (19)

<223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (956)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (970)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (981)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (989)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (999)
 <223> n equals a,t,g, or c

<400> 59
 gccatggtag catatgtant acatthttctt tatccagttc accgttgatg ggcacctggg 60
 ttactccat gtctttgcta ttgtgaatag tgctgcgata aacataggag tacagggtgtc 120
 tttttgatag aacaatttgt ttctctttga gtatatacct agtaatggga ttgctgggat 180
 gaatggtagt tctattttta gttctktgag aaatctccaa actgctggaa aacaagttht 240
 aatacacaag caccctcttt ccccaacctac agaaccttga cctcctttct agattgatct 300
 ctgctaataca cctcccaact ccccaacctc tggcttgac tagacttggt tgctattccc 360
 tggtaawtct catgagcatt ggctgaagca gttctctkt tctgaaatac cctgaccctc 420
 cgtacctagc taggcttttc tccaggcagc atttatgagg ccggtggtgt ctgagtgtcc 480
 tttgtggcca ggcaggacag agcaagggca agatgcttga tatatgctga caagacacca 540
 ttctagccgc acagatatct cagcaaattg gaagaccagc gtcgtccatt ctgacacctt 600
 gcacctthtt ggtacctttg tcccttggtg ctggctttgg agttaggatg ggtctgtctg 660
 tcctccacgt agggagttag agcgagtttc atggtgcacc ccacagagcc ctgcacgggg 720
 atttgagaat ggtaggcctc cagggaatgt tttgtatcca actacatgat ttagggggca 780
 caaaaatttc tccctggccc tgtgagagtc agagctgaga agcatcgaga aagtgtagct 840
 ctgggcgtgg ataactaatc agactgcata aatgacaaaa gccctctcat cccacagcag 900
 gttattggga gtcgtacctt gacttcagtt cattttgctc acatagacca aaaagnaaag 960
 cccaagaaan gacaatttgg ngaacttcng tagaaggtnc tgtaagaaga tc 1012

<210> 60
 <211> 1680
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (306)
 <223> n equals a,t,g, or c

<400> 60
 gccccggaaa cgthttctttc ctacgcagcc gctcctgccg ccgtggctgc tggagctttg 60

| | | | | | | |
|------------|-------------|-------------|-------------|-------------|-------------|------|
| cctctctagg | ccggcagcgc | ctctcctcca | tggctcgtgc | tgtcagcgct | gttttgggag | 120 |
| cccgcgggtg | aggccggggc | acgctcagac | acttcgatcg | tcgagtctgt | cactggggcat | 180 |
| ggcgggtcag | ttccgcagct | acgtgtggga | cccgcgtgctg | atcctgtcgc | agatcgtcct | 240 |
| catgcagacc | gtgtattacg | gctcgtctggg | cctgtggctg | gcgctgggtgg | acgggctagt | 300 |
| gcgaanagcc | cctcgtctgga | ccagatgttc | gacgccgaga | tcctgggctt | ttccaccctt | 360 |
| ccaggccggc | tctccatgat | gtccttcac | ctcaacgccc | tcacctgtgc | cctgggcttg | 420 |
| ctgtacttca | ttccggcgagg | aaagcagtg | ctggatttca | ctgtcactgt | ccatttcttt | 480 |
| cacctcctgg | gctgctgggt | ctacagctcc | cgtttcccct | cggcgctgac | ctgggtggctg | 540 |
| gtccaagccg | tgtgcattgc | actcatggct | gtcatcgggg | agtacctgtg | catgcggacg | 600 |
| gagctcaagg | agataccctt | caactcagcc | cctaaatcca | atgtctagaa | tcaggccctt | 660 |
| tggacatcct | gctgacactt | gggcccctta | acaccttggtg | ctgtcagac | cctccagatg | 720 |
| aggtccagcc | cagatctgag | aggaaccctg | gaaatgtgaa | gtctctgttg | gtgtgggaga | 780 |
| gatagtgagg | gcctgtcaaa | gaaggcaggt | agcagtcagc | atgacagctg | caagaatgac | 840 |
| ctctgtctgt | tgaagccttg | gtatctgaga | ggtcaggaag | gggacctctt | tgagggtaat | 900 |
| aacakaattg | gaaccatgcc | actcttgagc | cacaatacct | gtcaccagcc | tgtgttttta | 960 |
| agagagaaaa | aaaatcaagg | atatctgatt | ggagcaaacc | acttcttttag | tcactctgtct | 1020 |
| taccyccctg | ggacagctgt | tacctttgca | gtgttgccga | atcacagcag | ttacctttgc | 1080 |
| artgttgccg | aatcacagca | gttctgttgg | agaaacgctt | ggtttccgga | tccagagcca | 1140 |
| cagaaagaaa | tgtaggtgtg | aagtattagg | ctgctgtcag | ggagaggatg | gcagatggag | 1200 |
| gcatcaagca | caaggaaaat | gcacaacctg | tgccctgtta | tacacacggt | catgtgcacc | 1260 |
| caagaaccta | tgaactttctt | ccagttcctt | ctaccaggtc | cccacctctg | tgccagctct | 1320 |
| caacatagca | ggccatagga | cccagagaag | aatcccagyg | ttgctcaaag | tctaaccatc | 1380 |
| ataaagacac | tgcctgtctt | ctaggaatga | ccaggcacc | agctcccact | ggactccaat | 1440 |
| tttttttctt | gccttattta | gaattctttg | gcgggaagg | tatgatgggt | tcccagagac | 1500 |
| aagaagccca | accttctggc | ctgggctgtg | ctgatagtgc | tgaggagat | aggaatttgc | 1560 |
| tgctaagatt | tttctttggg | gtggagtctt | ctctgtgagg | ggcttgacgc | tatccttctt | 1620 |
| gtgtatacaa | atacagtatt | ttccatgaaa | aaaaaaaaa | aaaaaaaaa | aaaaaaaaa | 1680 |

<210> 61
 <211> 1386
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (1313)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1329)
 <223> n equals a,t,g, or c

| | | | | | | |
|-------------|------------|-------------|------------|-------------|------------|-----|
| <400> 61 | | | | | | |
| aggccccctgg | ttctgtgaaa | gaagggtgaaa | accttccaga | gagagggtgga | ggcgagcccc | 60 |
| aagagctcac | accctcccc | tccacttccc | ctctcctgca | ccaactgctc | agaggcagga | 120 |
| agttgagggg | gtgtgtgagc | ccccaccaca | gacacacaca | caacaaacca | aaacacataa | 180 |
| acacaaatac | ctacacacat | aagtagatac | acaccactgc | actccagcct | gggcaacaga | 240 |
| gccagactcc | gtctcaaaaa | awtaaaaaata | aaatatgcac | atctgtgccc | tcctagtctt | 300 |
| aaacagctac | tgaattatgc | tgtataataat | gggctttcca | gtgctaatta | tttctgcctc | 360 |
| tataccaaaag | tgccacaggg | catgcttctc | ttggcttctg | ggctctcttc | tgcttccctt | 420 |
| tatgactccc | tgggatttac | gctttccatg | ctgctgtttt | tcgagagaaa | caaatcacgt | 480 |
| gttgaagtcc | tggctaagga | gccgtcagct | cccagcagtt | attgggactc | tgagaaccgt | 540 |
| ggttgctcagc | tgtgaggtgc | aggttccttg | ttctcagaag | cgttctagct | ggcgggagct | 600 |
| ggacagggtt | cctctggaac | acaattttac | ctggtttcac | atctgactgg | caggaaaaag | 660 |
| ggaagcaggg | caagtgaggt | tagggctacc | cgggacagac | ggggatggac | ggagccatgt | 720 |
| cgctcagcgcc | cagtggagtt | ctgactgtaa | aggaacacac | cggtagcgcc | tccttgtaga | 780 |

| | | | | | | |
|------------|-------------|-------------|-------------|------------|------------|------|
| ggatgctatc | ccttccccac | caccgggtgat | tttgtctctg | ggataaatgt | gaattttcac | 840 |
| acctgttctc | ttagggtagg | agtcagcaaa | ctatggctca | caggccacat | ccagcctgct | 900 |
| gcctgttttt | tagggccatg | agctaagaac | taagaatggc | ttttatattt | gtaattgggt | 960 |
| gggaaaaaaa | aaaaaaaaarg | gcggccgstc | taraggatcc | ctcgaggggc | ccaagcttac | 1020 |
| gcstgcatgc | gacgkcatar | ctctctccct | atagggagtc | gaattataag | ctaggcactg | 1080 |
| gccgtcgttt | tacaacgtcg | tgactgggag | atctgctagc | ttgggatctt | tgtgaaggaa | 1140 |
| ccttacttct | gtggtgtgac | ataattggac | aaactaccta | cagagattta | aagctctaag | 1200 |
| gtaaatataa | aatttttaag | tgtataatgt | gtttaaactag | ctgcatatgc | ttgctgcttg | 1260 |
| agagttttgc | ttactgagta | tgattttatga | aaatattata | cacaggagct | agngattcta | 1320 |
| attgtttgng | tatttttagat | tcacagtccc | aaggctcatt | tcaggcccct | cagtcctcac | 1380 |
| agtctg | | | | | | 1386 |

<210> 62

<211> 716

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (630)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (710)

<223> n equals a,t,g, or c

<400> 62

| | | | | | | |
|------------|-------------|------------|------------|------------|------------|-----|
| gcctcagcgg | ccggggccac | ggccccgagc | agccatgctg | ggcgcgcggg | cctggttggg | 60 |
| ccgcgtcctt | ctgctgcccc | gcgcgggtgc | aggcctcgcc | gcragccgca | ggtgtcctgg | 120 |
| agtctggccc | aggacctggc | cccacaggag | tcccagcagg | ggtagctcct | cccgggacaa | 180 |
| ggaccgaagt | gcgacgggtca | gtagttcagt | gcccattgct | gctggaggga | aaggaagcca | 240 |
| tccttcattc | acaccccaga | gggtcccca | ccgcctgata | cacgagaagt | caccatacct | 300 |
| cctacaacat | gcctacaatc | ctgtggactg | gtacccctgg | ggacaggaag | ccttcgacaa | 360 |
| ggccaggaag | gaaaacaagc | cgattttcct | ctcagtcggg | tactccacct | gccactggtg | 420 |
| ccacatgatg | gaagaggagt | ccttccagaa | tgaggagatt | ggccgcctgc | tcagtgagga | 480 |
| ctttgtgagt | gtgaaggtag | accgtgagga | gcggcctgac | gtggacaagg | tgtacatgac | 540 |
| gttcgtgcag | gccaccagca | gcggcggggg | ctggcccatg | aatgtgtggc | tgactcccaa | 600 |
| cctccagccc | tttgtcgggg | gcactatttn | cctcctgaag | gatggcttga | mccgagtsgg | 660 |
| ttccgcacag | tgttkctgag | aatacgagaa | cartggaaac | agaacaagan | caccct | 716 |

<210> 63

<211> 2716

<212> DNA

<213> Homo sapiens

<400> 63

| | | | | | | |
|-------------|------------|------------|------------|------------|------------|-----|
| ggccggggccc | acggcmccga | gcagccatgc | tgggcgcgcg | ggcctgggtg | ggcgcgtccc | 60 |
| ttctgctgcc | ccgcgcgggt | gcaggcctcg | ccgcgagccg | caggctctgc | tgcagtccca | 120 |
| cttccaggct | gaactccctg | aggtctctga | ttccctaggt | gtcctggagt | ctggcccagg | 180 |
| acctggcccc | acaggagtcc | cagcaggggt | agctcctccc | gggacaagga | ccgaagtgcg | 240 |
| acggtcagta | gttcagtgcc | catgcctgct | ggagggaag | gaagccatcc | ttcatctaca | 300 |
| ccccagaggk | tccccaaccc | cctgatccac | gagaagtcac | catacctcct | acaacatgcc | 360 |
| tacaatcctg | tggactggta | cccctgggga | saggaagcct | tygacaaggc | caggaaggaa | 420 |
| aacaagccga | ttttcctctc | agtcgggtac | tccacctgcc | actggtgcca | catgatggaa | 480 |
| gaggagtcct | tccagaatga | ggagattggc | cgctgctca | gtgaggactt | tgtgagtgtg | 540 |

| | | | | | | |
|-------------|------------|------------|------------|-------------|-------------|------|
| aaggtagacc | gtgaggagcg | gcctgacgtg | gacaagggtg | acatgacgtt | cgtgcaggcc | 600 |
| accagcagcg | gcgggggctg | gcccataaat | gtgtggctga | ctcccaacct | ccagcccttt | 660 |
| gtcgggggca | cctatttccc | tcctgaggat | ggcttgaccc | gagtcggctt | ccgcacagt | 720 |
| ttgctgagaa | tacgagaaca | gtggaaacag | aacaagaaca | ccctgctaga | aaatagccag | 780 |
| cgtgtcacca | ctgccctgct | ggcccgatca | gagatcagcg | tgggtgaccg | ccagctgccg | 840 |
| ccctctgccg | ccaccgtgaa | caatcgctgc | ttccagcagc | tggatgaggg | ctatgatgag | 900 |
| gaatacggtg | gcttcgctga | ggcccccaag | tttcccacgc | cggatgatcct | gagcttcctg | 960 |
| ttctcctact | ggctcagcca | tcgactgact | caggatggct | ctcggggcca | gcagatggcc | 1020 |
| ttgcataccc | tgaatatgat | ggctaacggg | ggcatccggg | accatgtggg | gcagggcttt | 1080 |
| caccgctact | ccacagaccg | ccagtggcac | gtccctcact | ttgagaagat | gctctatgac | 1140 |
| caggcacagc | tcgctgtggc | ctattcgca | gccttcacgc | tctctggtga | tgaattctac | 1200 |
| tctgacgtgg | ccaaaggcat | cctgcagtac | gtggctcgga | gcctgagcca | ccggtccgga | 1260 |
| ggcttctata | gcgcagarga | tgcagactcg | ccccagagc | ggggccagcg | gccccaaagag | 1320 |
| ggcgcctact | atgtgtggac | ggtcaaagag | gttcagc | tcctcccgga | gcctgtgttg | 1380 |
| gggtgccaccg | agccgctgac | ctcaggccag | ctcctcacya | agcactacgg | cctcacagag | 1440 |
| gctggttaaca | tcagccccag | tcaggacccc | aagggggagc | tgcagggcca | gaatgtgctg | 1500 |
| accgtccggg | actcgctgga | gctgactgct | gcccgccttg | gcttggatgt | ggaggccgtg | 1560 |
| cggaccttgc | tcaattcagg | gctggagaag | ctcttcacag | cccgaagca | tcggcccaag | 1620 |
| ccgcacctgg | acagcaagat | gctggctgcc | tggaaatggc | tgatgggtgc | aggctatgct | 1680 |
| gtgactgggg | ctgtcctggg | ccaagacagg | ctgatcaact | atgccacca | tgggtgccaag | 1740 |
| ttcctgaagc | ggcacatggt | tgatgtggcc | agtggccgcc | tgatgcgga | ctgctacacc | 1800 |
| ggccctgggg | ggactgtgga | gcacagcaac | ccaccctgct | ggggcttcct | ggaggactac | 1860 |
| gccttcgtgg | tgcggggcct | gctggacctg | tatgaggcct | cacaggagag | tgcgtggctc | 1920 |
| gagtgggctc | tgcggctgca | ggacacacag | gacaggctct | tttgggactc | ccaggggtggc | 1980 |
| ggctacttct | gcagtgaggg | tgagctgggg | gctggcctgc | ccctgcgtct | gaaggacgac | 2040 |
| caggatggag | cagagcccag | cgccaattcc | gtgtcagccc | acaacctgct | ccggctgcat | 2100 |
| ggcttcacgg | gccacaagga | ctggatggac | aagtgtgtgt | gcctattgac | cgcttttcc | 2160 |
| gagcgcacgt | gtcgtgtccc | ggtggcgttg | cccagatgg | tcgcgcacct | ctcagcccag | 2220 |
| cagcagaccc | tcaagcagat | cgtgatctgt | ggagaccgtc | aggccaagga | caccaaggcc | 2280 |
| ctgggtgacg | gcgtccactc | tgtctacatt | cctaacaagg | tgtgtattct | ggctgatggg | 2340 |
| gacccctcga | gcttcctgtc | ccgccagctg | cctttcctga | gtaccctccg | acggttgga | 2400 |
| gaccaggcca | ctgcatatgt | gtgtgagaat | caagcctgct | cagtgcccat | cactgatccc | 2460 |
| tgcgaattac | gaaaactact | acatccatga | ctgccccaac | ccccttgggg | tggggcagaa | 2520 |
| ggtgaagcat | cccaactgac | tagagactca | ggccctgcag | ggccctatag | aacctgtggc | 2580 |
| catccctgag | cacctgcca | ccaggtgacc | tcggccatac | tcactgcccc | ccttgggcac | 2640 |
| ccactcacc | tagaataaac | ttaacagtgt | cccgtggtaa | aaaaaaaaaa | aaaaaaaaaa | 2700 |
| aaaaaagggc | ggccgc | | | | | 2716 |

<210> 64
 <211> 1011
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (68)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (950)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (993)
 <223> n equals a,t,g, or c

<220>

<221> SITE

<222> (1008)

<223> n equals a,t,g, or c

<400> 64

| | | | | | | |
|-------------|------------|-------------|------------|------------|------------|------|
| ctttgttgac | tcagtggggc | tcattgattgg | ttgtggagta | tcctgagaag | caggagtcac | 60 |
| agtaagggnag | gagaacaccc | ctcttattgt | cttatgccca | atttctgcct | ccaaagaaaa | 120 |
| aagaattaaa | aactgaagg | cagaaatgaa | atccacaggc | aggcagccca | gcaacgtgcc | 180 |
| ttgggcctag | tagttaacaa | tcaactcctg | acttaactgc | ttgtgtccca | gacattgtat | 240 |
| ggaaaagcat | cgtgaaaatc | cctgtcytgt | tctgttccat | tctgattacc | ggtgcatgca | 300 |
| gccccagtc | aggtaccccc | tgcttgctca | atcgatcatg | accctctcac | tcgacctcct | 360 |
| tagagttgta | agcccttaaa | agggacagga | attgctcact | aggggagctc | ggtttttttg | 420 |
| agacataggt | ctgctgatgc | tcctggccaa | taaagccctt | tccttcctca | actcggtgtc | 480 |
| tgagggggtt | tgtctgtggc | ttgtcctgct | acatttcttg | gttccttgat | ctggaagcaa | 540 |
| ggtgataagc | ggatggtcga | ggcagccct | taggcggctt | aggcctgccc | agtggagcat | 600 |
| ccctgtggga | gactctggcc | agcttttagcg | atgcgaatcc | tgagagcgct | ccccggcagg | 660 |
| caattgcccc | agtggaacgc | ctccagagga | gtgcacggta | ggcccccgcg | gaggattaat | 720 |
| gcagcggtg | aacaccggga | aggaattggc | gcttgagtc | cagacatctg | gaacacggta | 780 |
| agaccgggtc | tcgaacttgc | cactccatct | gagtggaaat | gtggcctgat | cacctatggc | 840 |
| gtgccttttt | ggcaccttgg | ttttgggttt | gattttgact | agatttgac | tggtttgggt | 900 |
| taratttcaa | gkattgactt | ttgggtttga | actggtttgg | ctttgattcn | ggtctgactt | 960 |
| ggctcaaaat | gcttgatgaa | taaagtaact | tentatccat | actttggntt | a | 1011 |

<210> 65

<211> 940

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (940)

<223> n equals a,t,g, or c

<400> 65

| | | | | | | |
|------------|-------------|------------|------------|-------------|------------|-----|
| gaawttaata | agctcctaac | ataaatttga | tgtcatttgc | agatgagaaa | atggggcttg | 60 |
| atgaagttga | ttgcttcaat | gatgcaacca | gtcctgcttg | aactcctctc | tgtctggagg | 120 |
| aaggaaggca | gagactctag | aaacatacat | gattcccaca | gcatgtatgt | gctgaggaag | 180 |
| agactgagtg | ggtcttggct | tcagcaggtc | tgtactctac | tgtgatataca | gcagactca | 240 |
| tccttgaatc | tccaagagct | tgtcttccct | caaaaggctg | attcagacac | agagtctaga | 300 |
| cttgtttctc | ttttcttaaa | agtacacatc | aggaagaaaa | cctaasgcct | cttgtaaaaa | 360 |
| catcactgac | atgcttccctg | gtgatttytc | agcagctatt | tgtccccag | atatattttt | 420 |
| tcactgccat | ttcacccaat | gttaattgaa | atgctaaact | aaatgttttag | ccatccttgc | 480 |
| atgcatggat | aattaataaa | ctgattataa | aattatagtg | atgagggatt | actaagccat | 540 |
| gataaattac | aatcattatt | tcagtcttga | ctataaagat | tattataagg | atgatagagg | 600 |
| kttcttatgg | attttttagtc | atttacacat | aatgaagtaa | aaatgkcttt | ttcaattttc | 660 |
| aaatatgtca | ccgaagccaa | aattggatct | tgtcattttt | aagtttttga | tatgcttgat | 720 |
| ttctgttgta | tttgwttcag | attgwyttg | cttcatgaaa | aggatgtttt | cttttttcag | 780 |
| gtagtgaggg | gtttaactac | attttatttc | tcccaaaagc | ttattaatct | tggcaataaa | 840 |
| aattctatat | ttctccagca | aaatctaaga | tgcttacctc | aaagcagctt | ttgaagttat | 900 |
| ttctatcagt | attggtctga | ttatgctgtg | aggagcctgn | | | 940 |

<210> 66

<211> 490

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (15)

<223> n equals a,t,g, or c

<400> 66

| | | | | | | |
|------------|------------|------------|------------|------------|------------|-----|
| gcatgcacgg | taagntgggc | ccttcggagg | gatctttaga | gcgccgccct | tttttttttt | 60 |
| ttcactttta | aatcatcct | aaagtgaag | ttattaaaag | tgaaaaaaag | aaaagaaaat | 120 |
| cctctccacc | tgttccacag | aatcttcagg | taagtctaaa | tttctcacca | aagtgtgtgc | 180 |
| tattcatgaa | gagttcccga | atcctgcata | catcagcaca | gggaaaaaga | aggactctaa | 240 |
| gaaaagcacc | aaacagctgg | gttttagaag | tatggcatgt | gtgcaaataa | ccacaccaag | 300 |
| acaaaaatat | aaaccacgca | tacaaaagag | agaaatagaa | catgtgccac | tggatttcag | 360 |
| agggaaaagg | gagtattgct | ggatagggtt | aaaattgtta | acttgacact | taaggaggaa | 420 |
| aaccattaga | aagcattcaa | aaagggcggc | cgctcgcat | ctagaactag | tccggacgcg | 480 |
| tgggtcgacc | | | | | | 490 |

<210> 67

<211> 2409

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (694)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (716)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (755)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (761)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (791)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (808)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (880)

<223> n equals a,t,g, or c

<400> 67

```

ccacgcgtcc ggcgagacgc gttcaccacc atatgcactg ttgtcaactc ccctggagat      60
gcgcccgaagc cccacaggaa gccttcctcc tctgcctcct cttcctcatc ctctgcctcg      120
ttctccttgg atgcagccgg ggcctccctg gccacactcc ctggaggctc catcctgcag      180
ccgcggccct ccttgccctc ctccctccacg atgcacttgg ggcctgtggt ttccaaggcc      240
ctgagtacct cttgccttgt ttgctgcctc tgccaaaacc cggccaactt caaggacctt      300
ggggacctct gtggggcccta ctaccctgaa cactgcctcc ccaaaaagaa gccaaaactc      360
aaggagaagg tgcggccaga aggcacctgt gaggaggcct cgctgccgct tgagagaaca      420
ctcaaaggtc ccgagtgtgc agctgccgcc actgccggga agccccccag gcctgacggc      480
ccagctgacc cggccaagca gggccactg cgcaccagtg cccggggcct gtcccggagg      540
ctgcagagct gctactgctg tgatggccgg gaggatgggg gcgaggaggc agccccagcc      600
gacaagggtc gcaaacatga gtgcagcaag gaggctccgg cagagcccgg cggggaggcc      660
caggagcact ggggtgcatga ggcctgtgcc gtgnggaccg gcggcgtcta cctgngggcc      720
gggaagctct ttgggctgca ggaggccatg aaggnggccg nggacatgat gtgttccagc      780
tgccaagaag ncggggccac catcggnngc tgccacaaag gatgcctcca cacctaccac      840
taccctgtgt ccagcgatgc aggttgcata ttcatcgaan agaacttttc tttgaaatgt      900
cccaaacata agaggctgcc gtagtaatcc accccaacgg ccggaggagc cgccggagcc      960
cgctgccccg cccgccgccg aaggagagga gccgcctgcg cagcccccg gcttttgagc     1020
tgctcccagc gctggtccag agccgatcct tgatccgggt cccggatcgt ggatccggcc     1080
gcctagggct cagacttgcg gccccgggtt gggagggaaa cccgttccgg agccgcctgc     1140
tcccgaacc ggacggcaca gggcggttctt gcccaccca gggggcaggc ttgcggaggg     1200
ggagcccgcg gagcgccag actccccggg gcgctcagcc tccggcgagg gtgggagacg     1260
gctttgtcct ggggacactt tccctctgga atctcaagac gacgtggcac acattccacg     1320
tgggtgctgc cgccaccca gtcggtcgtg gcgtgcagct gggagccctg ggcttggggg     1380
tgggggtcga aacagtactg gaagaggcgg agggcggtc ctagctccgt ggactaggcg     1440
ggggagaaa gaaacctttc tgagagcggg ctaggccggc actggagagg ccggagcctt     1500
tggaacaaac cgtgcggaac gcgtccaggg gccttcccgc ccagcctttg ccagatctct     1560
cgtgcggttc gggcaaagcc ggggtagacc tgggctatgc tcagttaggg gttgcgggat     1620
ccccgagtgt gggcgggact gggacacctt ttggcctctg tttgtccctt ttccagtcct     1680
ccaccccacc cctggagccc agcctgggag cgcaaaaccc aagaagcggc cagaacgcac     1740
ctccggctcc ggcggacgcg cgaccgttgt gcaccaccag ggaccgccgc gcctactctg     1800
cacgggagca gggacagcgc tagatttcgt gtacaaaacc tgtgtacccc tctatatata     1860
tgttacatag aatgtatata tgttgggaac atgctcgtt ctcctgtgtg tcgccgcgt     1920
gcgtcgtcgc cccgcaacag agccccaacc gggcctttgc cgggtaaggg gctaccgca     1980
cgccacttgt ccacgcagcc accaccggcc cgggccagtc cctgccagtc cgtccgcctg     2040
tccgtccgtg tcctcagctc tgtccacgct tcgataggcc tgacgcagcc cccagcccag     2100
ggcgcgccct gcaacttcct gtacatatga ctgtaaaatg gtaaacgtgt gtattatata     2160
tggcctcggt atatagtgtg tatatatgta tacatatata tatatataat atatatgaag     2220
actgtaaatg ttaagacgac tagtggttctt attagtatat tgcttcacac tgaagattgt     2280
gtgtatcgag ctgtttctaa aagatgttta ttttccttaa gagtaaaaaa cagtcattgc     2340
attcagaaaa aaaaaaaaaa aaaaagtc aaagatata acgattgttt tggaaaaaaa     2400
aaaaaaaaa

```

<210> 68

<211> 737

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (1)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (21)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (369)

<223> n equals a,t,g, or c

<400> 68

| | | | | | | |
|-------------|------------|------------|------------|------------|------------|-----|
| nagagggaag | agagaaaaag | ntacaatgct | tgcgcgtycg | sctktmcgkc | csygtcctca | 60 |
| gctctgtcca | cgcttcgata | ggcctgacgc | agccccagc | ccagggccgc | cctagcaact | 120 |
| tcctgtacat | atgactgtaa | aatggtaaac | gtgtgtatta | tatctggcct | cgttatatag | 180 |
| tgtatatata | tgtatacata | tacatatata | taatatatat | gaagactgta | aatgttaaga | 240 |
| cgactagtgt | tcttattagt | atattgcttc | acactgaaga | ttgtgtgtat | cgagctgttt | 300 |
| ctaaaagatg | tttattttcc | ttaagagtaa | aaaacagtca | ttgcattcag | aaaaaaaaaa | 360 |
| aaaaaaaaang | tcaataaaga | tacaacgatt | gttttggaag | atctgcagcc | cgtggattcc | 420 |
| gaccagattc | agctgggagc | cgggccaggc | tttaggttgg | ggaatgggaa | tgaagggagg | 480 |
| ggctgggggg | gggggcatga | atggagtcag | ggagtgcggc | tttcacagaa | caggaaacct | 540 |
| ccccgcgcc | tgtgccccct | ctccagtgtg | gcggcaggtc | gggagggagg | aggcttcttt | 600 |
| gctgtgagat | gaccaggggc | cgggatgggg | gaggtgagac | gtgccagact | tcttgcaggg | 660 |
| agacccaagc | tgtagctcct | gtcccacaac | aggctctgga | agtcagtcca | tcctcccgtg | 720 |
| ccaccagggg | acctaact | | | | | 737 |

<210> 69

<211> 1471

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (798)

<223> n equals a,t,g, or c

<400> 69

| | | | | | | |
|-------------|------------|-------------|-------------|-------------|------------|------|
| tttctgaatg | caatgactgt | tttttactct | taaggaaaat | aaacatcttt | tagaaacagc | 60 |
| tcgatacaca | caatcttcag | tgtgaagcaa | tatactaata | agaacactag | tcgtcttaac | 120 |
| atttacagtc | ttcatatata | ttatatatat | gtatatgtat | acatatatat | acactatata | 180 |
| acgaggccag | atataataca | cacgtttacc | attttacagt | catatgtaca | ggaagtgtgt | 240 |
| aggggggccc | tgggctgggg | gctgcgtcag | gcctatcgaa | gcgtggacag | agctgaggac | 300 |
| acggacggac | aggcggacgg | actggcaggg | actggccccg | gccggtgggtg | gctgcgtgga | 360 |
| caagtggcgt | cgcggtagcc | ccttaccggg | caaaggcccc | gttggggctc | tgttgcgggc | 420 |
| gcacgacgca | cggcgggcag | acacgggaga | agcagcatg | ttcccaacat | atatacatte | 480 |
| tatgtaacat | atatatagag | gggtacacag | gttttgtaca | cgaatctag | cgctgtccct | 540 |
| gctcccgtgc | agagtgggag | cggcggtccc | tgggtgtgca | caacggtcgc | gcgtccggcg | 600 |
| gagccggagg | tgcgttctgg | ccgcttcttg | ggttttgcgc | tcccaggctg | ggctccaggg | 660 |
| gtgggggtgga | ggactggaaa | ggggacaaac | agaggccaaa | gggtgtccca | gtcccggcca | 720 |
| cactcgggga | tcccgcgaac | cctaactgag | catagcccar | gtctaccccg | gctttgcccg | 780 |
| aaccgcacga | gagatctngc | aaargctggg | cgggaaggcc | cctgracgcg | ttccgcacgg | 840 |
| tttgttccaa | aggctccggc | ctytccagt | ccggcctagc | ccgctctcag | aaaggcttcc | 900 |
| tttctccyc | gcctagtcca | cggagctagg | agccgccttc | cgctcttcc | agtactgttt | 960 |
| cgacccccac | ccccaaagcc | agggtctcca | gctgcacgcc | acgaccgact | ggggtggcgg | 1020 |
| cagcaccac | gtggaatgtg | tgccacgtcg | tcttgagatt | ccagagggaa | agtgtcccca | 1080 |
| ggacaaagcc | gtctcccacc | ctcgccggag | gctgagcgcc | ccggggagtc | tggccgctcc | 1140 |
| gcgggctccc | cctccgcaag | cctggcccct | gggggtgggca | agaacgccct | gtgccgtccg | 1200 |
| gttccgggag | cagkcggtc | cggaaacgggt | tttctcccca | acccggggcc | gcaagtctga | 1260 |
| gccctaggcg | gccggatcca | cgatccggga | cccggatcaa | ggatcggtc | tggaccagcg | 1320 |
| ctgggagcag | ctcaaaggcc | cgggggctgc | gcaggcggtc | cctctccttc | ggcggcgsgc | 1380 |
| gggcaggcgg | gctccggcgg | ctcctmcgrc | cgttgggggtg | gattactacg | gcagcctctt | 1440 |

atgtttggga catttcaaag gggaattctc t.

1471

<210> 70

<211> 3599

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (2675)

<223> n equals a,t,g, or c

<400> 70

| | | | | | | |
|-------------|-------------|-------------|-------------|-------------|-------------|------|
| gggagggcaa | cggggcggcg | cgggcggccg | ggcgagggt | ckcgggargt | gacrcgcggc | 60 |
| gaggatggcg | gcgcggggcc | gggggctgct | gctgctgacg | ctgtcggtgc | tggtggcggc | 120 |
| gggcccctcc | gccgtgcg | ycaagctcaa | catcccaaa | gtgctgctgc | ccttcacrcg | 180 |
| ggccacgcgc | gttaacttca | cgctggaggc | ctcggagggc | tgctaccgct | ggttgtccac | 240 |
| ccggccggag | gtggccagca | tcgagccgct | gggcctggac | gagcagcagt | gctcccagaa | 300 |
| ggcagtgtgtg | caggcccgc | tgaccagcc | tgcccgcctc | accagcatca | tcttcgcaga | 360 |
| ggacatcacc | acaggccagg | tcctgcgctg | tgatgccatt | gtggacctca | tccatgacat | 420 |
| ccagatcgtc | tccaccaccc | gcgagctcta | cctggaggac | tccccctgg | agctgaagat | 480 |
| ccaggccctg | gactccgaag | ggaacacctt | cagcactctg | gctggactgg | tcttcgagtg | 540 |
| gacgattgtg | aaggactccg | aggcggacag | gttctcagac | tcccacaatg | cgctgcgaat | 600 |
| cctcactttc | ttggagtcta | cgtacatccc | tccttcttac | atctcagaga | tggagaaggc | 660 |
| tgccaagcaa | ggggacacca | tcctggtgtc | tgggatgaag | accgggagct | ycaagctcaa | 720 |
| ggctcgcatc | caggaggctg | tctacaagaa | tgtacgccct | gcakaagtca | ggctgtgat | 780 |
| tttgaaaaac | atccttctga | acccggccta | tgacgtctac | ctgatggtgg | gaacctccat | 840 |
| tcactacaag | gtgcagaaga | tcaggcaagg | gaaaattaca | gaactctyca | tgcttccga | 900 |
| tcagtacgag | ttgcagcttc | agaacagcat | cccgggcccc | gaaggagacc | caaccggcc | 960 |
| ggtggctgtc | ttggcccagg | acacgtcgat | ggtcactgca | ctgcagctgg | gacagagcag | 1020 |
| cctcgtcctt | ggccacagga | gtattcgcat | gcaagggtgct | tctagggttac | ccaacagcac | 1080 |
| tatctacgtg | gtcgaacctg | gatacctagg | gttctactgtt | caccctggtg | acagggtgggt | 1140 |
| gctggagacc | ggccgcctgt | atgaaatcac | catcgaagtt | tttgacaagt | tcagcaacaa | 1200 |
| ggctctatgta | tctgacaaca | tccgaattga | aactgtgctt | cctgctgagt | tcttcgaggt | 1260 |
| gctctcgtcc | tcccagaatg | ggtcatacca | tcgcatcagg | gcactaaaga | ggggacagac | 1320 |
| ggccattgac | gcggccctca | cctctgtggt | ggaccaggat | ggaggggtcc | acatactaca | 1380 |
| ggtgcctgtg | tggaaccagc | aggaggtgga | aattcacatc | ccgatcacc | tgtatcccag | 1440 |
| catcttgaca | tttccgtggc | aaccaaagac | gggcgcctat | cagtacacaa | taaggggcca | 1500 |
| cgggtggcagt | gggaacttca | gctggtcttc | gtcaagccac | ctggttgcca | cagttactgt | 1560 |
| caagggcgctg | atgaccacag | gcagtgcacat | cgggttcagt | gtgatccagg | cacatgatgt | 1620 |
| gcagaaccca | ctccatttcg | gtgagatgaa | ggtgtatgtg | atcgagcccc | acagcatgga | 1680 |
| gtttgccccg | tgccagggtg | aggcacgtgt | gggccaggcc | ctggagctgc | ccctgaggat | 1740 |
| cagtggcctc | atgcccggcg | gggccagtga | ggtggtcacc | ttgagcgact | gctcccactt | 1800 |
| tgacttggtc | gtcgagggtg | agaaccaggg | tgtgttccag | ccactcccag | ggaggctgcc | 1860 |
| gccaggctct | gagcactgca | gcggcgtccg | ggtaaaggcc | gaggcccagg | gctctaccac | 1920 |
| gcttcttggtg | agctacagac | acggccacgt | ccacctgagt | gccaagatca | ccattgctgc | 1980 |
| ctacctgccc | ctcaaggctg | tggatccctc | ctctgttgcc | ttggtaaccc | tgggctcctc | 2040 |
| aaaggagatg | ctgtttgaag | gaggtcccag | accttggtac | ctcgagccgt | ccaaattctt | 2100 |
| ccagaagctc | accgctgagg | acactgacag | catcggcctg | gctctctttg | cccccatc | 2160 |
| ctcccggaat | tatcagcaac | actggatcct | tgtgacctgt | caggccttgg | gtgagcaggt | 2220 |
| catcgccctg | tcgggtgggga | acaagcccag | cctcaccaac | ccctttcctg | cgggtggagcc | 2280 |
| tgccgtggtg | aagttcgtct | gcgccccacc | gtccaggctc | accctcgtgc | ctgtctacac | 2340 |
| cagccccag | ctggacatgt | cctgtccgct | gctgcagcag | aacaagcagg | tggtcccagt | 2400 |
| gtccagccac | cgcaaccccc | tgctggacct | ggctgcttac | gaccaggagg | gccgcccgtt | 2460 |
| cgacaacttc | agctctctga | gcattccagt | ggagtccacc | aggccagtgt | tggccagcat | 2520 |
| cgagcctgag | ctgcccattgc | agctggtgtc | ccaggacgat | gagagtggcc | aaaagaagct | 2580 |
| gcacggtttg | caggccattt | tggttcacga | ggcatcagga | accacagcat | cactgccact | 2640 |

| | | | | | | |
|-------------|------------|-------------|-------------|-------------|-------------|------|
| gccactggct | accaggagtc | ccacctcagc | tctgncagaa | caaagcagcc | gcatgaccct | 2700 |
| ctgggtgcctc | tgtcggcctc | catagagctc | atcctgggtg | aggacgtgag | ggtgagccca | 2760 |
| gaagagggtga | ccatctacaa | ccacctggc | atccaggcag | agctccgcat | caggggaaggc | 2820 |
| tcagggttact | tcttcctcaa | caccagcacc | gcagatggtg | tcaagggtggc | ctaccaggag | 2880 |
| gccagggggtg | tcgccatggt | gcaccctttg | ctcccggtg | catccaccat | catgatccat | 2940 |
| gacttgtgcc | tcgtcttccc | ggccccagcc | aaggctgtcg | tttacgtgtc | ggacattcag | 3000 |
| gagctgtaca | tccgtgtggt | tgacaagggtg | gagattggga | agacagtga | ggcatacgtc | 3060 |
| cgcgtgtcg | acttgacaaa | gaagcccttc | cttgccaaat | acttccccct | catggacctg | 3120 |
| aagctccgag | cagcctcccc | gatcattaca | ttggtggccc | ttgatgaagc | ccttgacaac | 3180 |
| tacaccatca | cattcctcat | ccgcgggtgtg | gccatcgccc | agaccagtct | aactgcaagt | 3240 |
| gtgaccaata | aagctggaca | gagaatcaac | tcarccccac | aacagattga | agtctttccc | 3300 |
| ccgttcaggc | gaaggtgaca | ctgcttatcg | ggggccacgat | gcaggtcacc | | 3360 |
| tccaggggcg | gccccagcc | tcagtccaac | atccttttct | ccatcagcaa | tgagagcggt | 3420 |
| gcgctggtga | gcgctgtgtg | gctggtacag | ggcctcgcca | tcgggaacgg | cactgtgtct | 3480 |
| gggctcgtgc | aggcagtggg | tgagagacc | ggcaagggtg | tcacatctc | tcaggacctc | 3540 |
| gtgcagggtg | aagtgtgtgt | gctaaggggc | tgtctcttat | acacatctca | acctgcagg | 3599 |

<210> 71

<211> 2924

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (2892)

<223> n equals a,t,g, or c

<400> 71

| | | | | | | |
|-------------|-------------|-------------|------------|-------------|------------|------|
| gaggccaaga | ttcgcacgag | tttttttttt | ttttttaaaa | aatcacatag | aactttatta | 60 |
| aataaaccag | tagaaaggct | cctattttaca | tccagggtgt | cagtttaagc | tcaagtaata | 120 |
| tgcaaggcac | aattcttctt | ttctccagct | ttagaaaaat | caaccacatg | gcccagccct | 180 |
| gaaggactga | agggagcttg | tgcgggatcc | cgtcatcctg | gactctccac | acagcgtcca | 240 |
| cgcattggatg | gccatgggca | tgagtctctg | gatctccctg | agggtagccc | agcctcggcc | 300 |
| tggagcacag | gggcctgtga | tctgagaatc | tcagatccag | aatcaggagc | tgggaagaga | 360 |
| ggacttttcc | aagggtctcc | tccccaggg | cctggggagt | cccttctgga | tgggcaggcc | 420 |
| ctgggctggg | ggccgcccct | catcattggc | ccaggccggg | ctctaggggc | kgctgggaag | 480 |
| ctgtctgggg | ccctgcgcct | gatgagctgc | catgaaagtg | gccgcctctc | acactggcag | 540 |
| gcggtggctc | agatctgctg | aggttctgga | gtggcacagg | aacaggagga | gcaaatgaca | 600 |
| ctgggtcttt | gtgctgaggg | aggacacagc | ggctttccga | aacacacaca | cagaaaatat | 660 |
| gactcctgct | gccatctcaa | ggacattttc | ctaaggacag | tcctggtgtc | ctgatggagt | 720 |
| gaaagggatc | acacgtggga | tgtcccagca | gggcctaagc | caagtccatc | agatgtctca | 780 |
| acggcttatg | aaacaaaaac | agaggactga | agtctgttca | cagtggccca | tggttgggca | 840 |
| gcagccctgg | cctccagggtg | ctggttctctg | aggcttccct | tgcggcagtg | gcgcagatac | 900 |
| agggacactc | atgtcccatt | gtctctcaga | agctctagca | gcagccagac | taggggcaaa | 960 |
| gtaagttttc | acagggggaa | aaaccccacc | aaaaaccccc | taaagtaact | ggacaatcct | 1020 |
| ttccctgaga | ccactacagt | aacatcaccc | acagacaact | tctaaagagc | acttctaact | 1080 |
| gctcaaagtc | ttgagaaggg | aaagatttta | gcaaggctag | gaaaataaaa | atgcaacagc | 1140 |
| acattgtcca | gaaaccctag | acaaccatgc | aaaaaggaat | gagccaaaaa | gtcttagctt | 1200 |
| tgtaagactt | gaaaggaaaa | aaacacacat | ttaaaactcc | tacataaaaat | gagctaattg | 1260 |
| gcagagccga | aactacagct | ctgtgtgaag | agacggcagt | gaagctggcc | tggggccggg | 1320 |
| gcaatcatct | ggaggggctt | gttccagtgt | gaatgcagca | gggatgttcc | atcttggggg | 1380 |
| tgcacgaggc | tcggctgaga | cccatectcc | gggaaccttc | acgcggccta | gtgggaggca | 1440 |
| taggctgggc | tccacagccc | tgagggaggg | ctggctttgc | gagcaggagg | caatgcattg | 1500 |
| ggagatgtgg | ctgatgaggc | agcgaaatag | tgggggctgt | gtccagggct | ggctcgaggc | 1560 |
| gtgagggctg | caggcacagc | aagatcccgg | ggcgtgcaga | cagtgtggta | ggctatgatc | 1620 |
| atgaccgctg | tcccagccaa | cagggcgaa | agcgtgaaga | acatgacctg | gtaggaatcc | 1680 |
| aggaagtgtc | ggaagaggct | ggctccataa | ggaccggggc | cacggcgatc | caycacaaaa | 1740 |

| | | | | | | |
|------------|-------------|------------|-------------|------------|-------------|------|
| gccactgtca | ctgggatggc | aatggcttgg | ttgggtcacgg | ggctggagaa | ggtcagggta | 1800 |
| gtggacagag | gcccttggct | gccagccgcg | gggtccaaga | cgccgaccgt | gtatgtgatg | 1860 |
| aagctggggc | acccaaaaga | cttctccttt | gcgaatgcca | gcacggcccg | ggacccggat | 1920 |
| ttcacctcca | agttctccag | aacctccggg | gcaccaaaga | ccctgatctc | ggaactgggtg | 1980 |
| tagtggttgc | tcaaaaggat | ttcagcctgg | tcggcgaaaga | gacctgggct | gaagggcacc | 2040 |
| tcggccccc | cctgctctgt | ggagaagtgg | ctgctggaga | gggaggcact | gaccaccaga | 2100 |
| gctgtcttct | tcattgctcag | gtgcttccgc | tgcttgctccg | tcagcctgtg | cattgtgatt | 2160 |
| gagcagaagt | actggccgag | agcagtgtca | aactgtggct | ccacggtgaa | cacatcttga | 2220 |
| gatgggaaat | caaagacggc | cggcttgaac | tgggactggc | agctgatgag | ggtctctggg | 2280 |
| tgcaaggcct | ggatgacttc | cctctgggtg | ggggtgcact | cgctctcag | gtagagctt | 2340 |
| ctgtctccca | cggcaacaat | cactttggag | gctgtagcct | cctggaagct | ggtctggatg | 2400 |
| gggtggaggt | gacgggccat | gatcctctga | gggacgtga | ccaccacctc | cttgtaggtc | 2460 |
| ctcaggtgcc | cagcgacctc | atagtaaacc | gtcacggatc | ccacggcccc | ggccacagcc | 2520 |
| acacccgtct | tggggtcgat | gtggaggatg | ctgttgcccg | aggagctcca | ggttcctgag | 2580 |
| aggccttcca | ggctggtcag | aacagtggcc | agacagagca | cgtccccac | caccatggcc | 2640 |
| ccagacagct | ctggggagat | ggcctgtagg | acaggcaggg | gcatgaagtc | cgagaggccc | 2700 |
| gggtgctctg | cgtcccacac | acggagcagt | gtcaggccca | cgtgactgt | gcggacaaca | 2760 |
| caggtgttgt | tgggtggggc | cttcccgatc | tgcacaaagt | cgtctctgtt | agtggcaaag | 2820 |
| ttgaggaccg | aactgtgagc | atggaagaca | tctccagagt | tgtcgyggaa | gtggacagtg | 2880 |
| aaggtcaccg | gncattccca | aaggcaccgg | ccaccaaggg | cctc | | 2924 |

<210> 72

<211> 628

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (137)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (450)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (465)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (488)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (585)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (619)

<223> n equals a,t,g, or c

<400> 72

57

| | | | | | | |
|------------|------------|-------------|------------|------------|------------|-----|
| gcacctcagg | ccctccaagc | gcaggatgca | ggccgtggcc | aacgtgtcca | ttggggccat | 60 |
| gttctgcatg | tatgggctca | cagcaacctt | tggatacctc | accttctaca | gtragtgggg | 120 |
| ctggggctag | ggctggnngg | agggggaagg | cctggggcag | gagcctctga | gctctttcct | 180 |
| tctgtgacca | cggacctgtc | agttttccaaa | cagaargtgt | gcctcacttg | tgtggatttt | 240 |
| gtcactgtgc | atgtatgtat | gggtttcttg | ggcattggtc | ctggtgctct | ctccacatcc | 300 |
| tgcaccccg | acctctgtc | tcatggscce | rgcartgtga | aggcggagat | gctgcacatg | 360 |
| tacagccaga | aggacctgct | catcctctgt | gtgcgcctgg | ccgtgctgct | cgcggtgacc | 420 |
| ctcactgtgc | cagtcgtgct | gttccctatn | cgccgggccc | tgcanagct | gcttttccca | 480 |
| ggcaaggnc | tcagctggcc | acgacatgtg | gccataagct | ctgatctgct | tgtttggtca | 540 |
| atgtccttgt | catctgtgtg | ccaacatccg | ggatatcttt | ggganttaat | cgggtcaact | 600 |
| caagccccag | ctcatcttna | tctcccag | | | | 628 |

<210> 73
 <211> 923
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (9)
 <223> n equals a,t,g, or c

| | |
|-------------|-------------|
| <400> 73 | |
| gggaacggna | aactcgtttg |
| gcacacatgt | ggccatagct |
| tgccaacat | ccgggatata |
| tcacacatgt | ggccatagct |
| ctcactgtgc | atgcacatgg |
| aggggtccctc | ctgcccacaa |
| cgggggacag | cagaggctgc |
| ggatcatggc | cctaataccca |
| ggaggaggag | gaggaggagga |
| agcccagtc | tctctgcctc |
| aagaggaaaa | ataaagatgt |
| aaaaaaaaaa | aaaaagggag |
| | gcc |

<210> 74
 <211> 1297
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (1104)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1131)
 <223> n equals a,t,g, or c

<220>

<221> SITE
 <222> (1183)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1249)
 <223> n equals a,t,g, or c

<400> 74
 gagaagccat acctgggtgga agaggccggt tcttacaatg aacttgacta tgtgtcgggt 60
 ggccttgacc agcagacggt gaagctgggtg tgcaccaacc gcaggaagca gtttctgctg 120
 gacacggctg atgtggcgct ggctgagttc tttttggctt ctttgaagtc agccatgatc 180
 aaaggctgtc gagaacctcc ctacccacgc atcctgacgg atgccaccat ggagaagctg 240
 gcactggcca aatttgtggc ccaagaatcg aagtgtgagg catctgctgt caccgtgcgc 300
 ttctacggcc ttgtgactg ggaggacccc acagacgagt ccctggggccc cagccctgc 360
 cactgctcac cccccgaggg caccatcacc aaagaaggca tgctgcacta caaggcgggc 420
 acctcctacc tgggcaagga aacttggaag acgtgcttcg tgggtgctcag caacgggatc 480
 ctctaccagt acccggaccg caccgacgtc atccctctgc tctcggtgaa catggggggg 540
 gagcagtgcg gtggctgccg gagagccaac accacggatc ggccccacgc cttccaggtc 600
 attctctccg accggccctg cctggagcta agtgccgaga gcgaggccga gatggccgag 660
 tggatgcagc atctctgccg ggctgtgtcc aaaggggtca tccccagggg cgtagctccc 720
 agccctgca taccctgctg cctggtcctc acggatgacc gcctctttac gtgccatgag 780
 gattgccaga ccagcttctt ccgctctttg ggcacagcca agctgggcca catcagcgcc 840
 gtctccaccg agccgggcaa ggagtactgc gtcttgagg tctcccagga cagccagcag 900
 ctctcccgcc cctgggtcat ctacctgagc tgcacttctg aactggaccg attgctgkct 960
 gcactgaact ctgggtggaa aaccatctat cagggtggacc tccccacac ggcatccag 1020
 gaagccttca acaagaagaa attcaggat gccttgagcc tcatccacag cgcctggcag 1080
 cggagcgaca gtctctgccg cggncgagcc tccgagacc ctggtgctga ngcagaactg 1140
 gttggcgctc ctgggtggcag gaaaggaagc acgcagccg cangcacact gcacggctgt 1200
 gcattgctgc ggagctaagt cacctgcctg ggcggaaaac accgagtng gttaagaagg 1260
 tcctcatcag ggatcagaca gtgccggacc tggatct 1297

<210> 75
 <211> 2550
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (1190)
 <223> n equals a,t,g, or c

<400> 75
 cgaccagctc aacgggcagc tggaccccag cacctgggtgc tcccgctgctg agccccaga 60
 ccagtccttt cggaccgggt ctcccgggga tgcccggag aggcgcgcg tttgcgactt 120
 tagtgagggg ctttcagccc aatggacttc taccgcttta ccgtcgagag tccaagcact 180
 gttacatcag gtggcgccca ccatgacct gcagggttg gccaaccgct gcatttctc 240
 agtgccctg aggmtgctgg ccaagaagaa gagggaggag gagagaggg acagacgctc 300
 cggccctag aggataccac gagggaggct caggagctgg aggccagct gtccctggtc 360
 agggaggggc ctgtgtctga gccagagcct gggaccagg aggttctctg ccagctcaag 420
 cgagaccagc ccagcccggt tctgagtarc gctgargatt ctggggtgga tgaggacag 480
 gggagccctt cggagatggt ccattcctcg gagttcagag tagacaacaa tcacctgctc 540
 ctgctcatga tccacgtgtt ccgagaaaac gargagcagc tgttcaaaat gatccggatg 600
 agcaccgggc acatggaggg caacctgcag ctgctgtacg tgctgtcac agactgctat 660
 gtctacctgc tccggaaagg ggccacagag aagccatacc tgggtggaaga ggccgtttct 720
 tacaatgaac ttgactatgt gtcggttggc cttgaccagc agacggtgaa gctgggtgtg 780

| | | | | | | |
|-------------|------------|-------------|------------|-------------|-------------|------|
| accaaccgca | ggaagcagtt | tctgctggac | acggctgatg | tggcgctggc | tgagttcttt | 840 |
| ttggcttctt | tgaagtcagc | catgatcaaa | ggctgtcgag | aacctcccta | ccccagcatc | 900 |
| ctgacrgatg | ccaccatgga | gaagctggca | ctggccaaat | ttgtggccca | agaatcgaag | 960 |
| tgtgagggcat | ctgctgtcac | cgtgcgcttc | tacggccttg | tgcactggga | ggacccca | 1020 |
| gacgagtc | tggtgcccac | gccctgccac | tgctacccc | ccgagggcac | catcaccaaa | 1080 |
| gaaggcatgc | tgcactacaa | ggcgggcacc | tcctaccttg | gcaaggaaca | ctggaagacg | 1140 |
| tgcttcgtgg | tgctcagcaa | cgggatcctc | taccagtacc | cggaccgcan | cgacgtcatc | 1200 |
| cctctgctct | csgtgawcat | gggggtcatcc | cccagggcgt | agctcccagc | ccctgcatac | 1260 |
| cctgctgcct | ggctctcacg | gatgaccgcc | tctttacgtg | ccatgaggat | tgccagacca | 1320 |
| gcttctccg | ctctttgggc | acagccaagc | tgggcgacat | cagcgccgtc | tccaccgagc | 1380 |
| cgggcaaggga | gtactgcgtc | ttggagttct | cccaggacag | ccagcagctc | ctcccgcct | 1440 |
| gggtcatcta | cctgagctgc | acttctgaac | tggaccgatt | gctgtctgca | ctgaactctg | 1500 |
| ggtggaac | catctatcag | gtggacctcc | cccacacggc | gatccaggaa | gcctccaaca | 1560 |
| agaagaaatt | cgaggatgcc | ttgagcctca | tccacagcgc | ctggcagcgg | agcgacagtc | 1620 |
| tctgcccg | ccgagcctcc | cgagaccctc | ggtgctgagg | cagagctggg | tggcgctcct | 1680 |
| ggtggcgagg | aaaggaaggc | acgccagccg | gcaggcacac | tgtcacggct | gttgtcatgc | 1740 |
| tgtcgggagc | ctacagtcca | cccctgccct | gggcggcaga | accaccgagt | gtggcttaag | 1800 |
| acagggctcc | tccactccag | ggatccagat | caggtgcccg | gcacccctgg | gcacccctgcc | 1860 |
| cgacaggtag | cgaatggagg | tcgctggggg | cagagggctc | gagccctgtg | ggctctgcgg | 1920 |
| atgcacgccc | tcctcccg | cctccgcctc | agtctgcaga | atttctgccg | agtggcaccg | 1980 |
| agaacaccat | ccatctaagg | acgaacaaaa | gaaccaggag | ggcgggaccc | ccctcttctc | 2040 |
| ctcctgggtt | gggggctggg | gccctgagtg | cccagccatc | cttgttcgtg | tttgaacact | 2100 |
| ctcctggcca | cgtggggaag | cgggaacacg | gggtgtctgc | gcatgtttcc | tcctcctagc | 2160 |
| tccatcactg | cgcacacagc | tgctgcctc | gccagatgca | ggggggcggg | cagccctccc | 2220 |
| tggctgccag | gaggtctctg | atgcccacag | tcctgcctg | cctgtcccct | caacccggca | 2280 |
| gtgctgtag | caccgaggag | caaagggggg | ggatgggggg | cttgagagaag | gcggagccca | 2340 |
| ccagctggc | atccatgttg | acatcttctg | actgtcccct | gcttggtggg | agccaggccc | 2400 |
| ttccctagag | tttcgtcaag | agcctcctgg | ggaaggggtc | agggtggttg | ggttttgttt | 2460 |
| tttaaaataa | aatagacatg | ttatattgcc | aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | 2520 |
| aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | | | | 2550 |

<210> 76

<211> 308

<212> DNA

<213> Homo sapiens

<400> 76

| | | | | | | |
|-------------|------------|------------|------------|-------------|-------------|-----|
| ggagggcagga | ccttgtccta | ttcattaatc | ttgcccctca | acagttat | tcagaggggc | 60 |
| aagaagtgtt | tcagggttct | tggcccttgt | ttgaccagtc | gtcctaacc | tcrtgtcttg | 120 |
| ggatcattgt | gttrtaatct | gggtttacct | tttggaaggt | catgggggtac | ccttttgcaa | 180 |
| aagttatggg | ccctmtcctt | ggaaactgca | cacacaccat | gcagcttaca | attcaggggag | 240 |
| ttcacaggtc | tacagaatcc | tgggaaactc | tccatgtccg | gttctaattc | attgtagctt | 300 |
| cagtggga | | | | | | 308 |

<210> 77

<211> 1568

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (1550)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (1564)

<223> n equals a,t,g, or c

<400> 77

| | | | | | | |
|-------------|-------------|------------|-------------|-------------|------------|------|
| ctcatcagcc | ctatgaggta | gggaggtagg | tattattatc | acccttggtta | gtttttttgt | 60 |
| tttggtttgt | tttccagatg | agagaatcat | tgctcacaga | agtgaagtaa | cttttccaag | 120 |
| gtcatacaat | cagtaagtgg | caggcaagga | ctgaaatcca | agttggtacc | ctccaaagtc | 180 |
| cctgctctga | gaactggagg | aattctttat | caaattctaaa | atcctctttt | agscctgtct | 240 |
| gctttaatat | tccggtcttt | attgatcctt | ctcttctcta | aamcttcagc | tgtcagtata | 300 |
| aaaaatcaagg | aatttagcmc | ttgttattgt | gtgamcagct | tcttgtctct | cctgtactgt | 360 |
| aagtgggtct | agggattttt | attctttaaa | tatccccctg | tactcagtag | atctttggga | 420 |
| gamcaagctc | ataggcttct | aataattctt | tctttgactg | ccagctgaat | tagacagaag | 480 |
| gtaagtccctg | ctgccgtgtc | gtgcctaacc | ccatctttat | ttcctgtgct | gttagagaac | 540 |
| agtcttttct | tggtctggaac | aaatactaca | gcctgctcaa | ctagctaata | tgtattgagt | 600 |
| tcttartatg | ttccaaggac | tgctctaagt | attttatata | tattaactca | ctgaatctta | 660 |
| aataccctat | gagctaagtc | ctatttttat | ccccatttta | caaaagagga | aactgaatgt | 720 |
| accagtgcac | cagtatttga | ctgagtaaat | gaatgactgc | tttgctgatg | gatagtatta | 780 |
| ttagcaacaa | ccctacaaat | atgatgttat | gtttgcatca | tgcagtacag | ctttatgtac | 840 |
| cttatgtcat | tgtcactcat | gattagcaaa | taggcacgaa | catccctatt | ttatagaaga | 900 |
| ggaamccatg | gctctaagag | ggtgagtgat | tctcaacagt | cacatgccat | ctgtatcctt | 960 |
| cagtaaacaa | ggatatttgg | ccattccagg | atcgggggca | agagagatgg | gagggcctcg | 1020 |
| gtgagaaaca | ctcatattca | caaaaggtag | tagatagata | gacagataaa | taataaaata | 1080 |
| gagataaaaag | ctagtaatag | cagagatttg | atgggaattc | agatctttga | ttcctagtcc | 1140 |
| agtgtctctt | cttttatgta | aggtgatggg | aagcaagtct | taggtccaga | tctggcagct | 1200 |
| gctttgat | aggatcttaa | gccagaagca | gcagcgccct | aaacaaaaag | catcatttta | 1260 |
| acttctctgc | atttctcaca | ttctcaacaa | tgaccatgcc | ctactttcat | aattaaaaac | 1320 |
| aaacaaataa | acagaggggg | caaaatgcat | ccttctgagt | gggctggggc | ttgtagaggg | 1380 |
| attcttgggt | ttgtctctgg | atgtctttgg | gcccctgcac | ttgtgggcac | tctgatttat | 1440 |
| ccccacaggc | cacctggccc | accttatggg | ctgaggaggg | cttgatgggc | ggaggsaagc | 1500 |
| agagctgagg | agctggggag | gagctggaaa | ggactgctag | aagggtccan | tgaagctaca | 1560 |
| atgnntag | | | | | | 1568 |

<210> 78

<211> 865

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (846)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (853)

<223> n equals a,t,g, or c

<400> 78

| | | | | | | |
|------------|------------|------------|-------------|------------|------------|-----|
| catgcacgcc | taagcttggg | cccctcgagg | gatcctctag | agcggccgcc | cttttttttt | 60 |
| tttttttttt | tttttttgag | atagagtttt | gctcttggtg | cccaggctgg | agtgcaatag | 120 |
| tgcgatcttt | ggctactgca | acctctgcct | cccgggttca | agtgattctc | ccgtcccagc | 180 |
| ctccccagta | gctgggatta | caggcgtgca | ccaccacgcc | cagctaattt | tgtattgttt | 240 |
| agtagagatg | gggtttctct | gttttggcca | ggctgggtcac | gaactccgca | cctcaggtga | 300 |
| ttcgcgccgc | ttcgtggcgt | gagcmacgct | cccggccgat | gattctttct | tttaatgagt | 360 |
| ctccataaag | tcaatttttg | gaatgataaa | ggaggtgata | atacaatagc | aaaatccgtg | 420 |
| aggggtgtaa | gaaactggag | tctgggaaat | gcctccagac | accaaagtag | agaaaaggag | 480 |
| agaaaggaag | ggaaaccagc | caggggctct | ctaccagag | gctacttaca | tttaatgtat | 540 |

| | | | | | | |
|------------|------------|------------|------------|------------|------------|-----|
| agttctctcc | cagtgGCCCA | cccaggctct | tactaaggag | aaaatacaat | gctcagttga | 600 |
| tgGCacagac | ttcccatTTA | acaaataaaa | accttaaaaa | aaaagaaaaa | gaaaaaagag | 660 |
| gcCagcacag | tggtcCagcC | ctgtaatccc | agcatttcca | gaggccaaga | caggaggact | 720 |
| gcttgagggt | aggagtttga | gaccagcccg | ggcaacacag | ggagacttca | tgtctaccma | 780 |
| aaaaacttca | aaattagtgg | arctgkggtc | ccagctamtc | aggaggctga | rgcsaragga | 840 |
| atgctngagc | ccngcaggct | gcggt | | | | 865 |

```
<210> 79
<211> 1182
<212> DNA
<213> Homo sapiens
```

```
<220>  
<221> SITE  
<222> (1134)  
<223> n equals a,t,g, or c
```

```
<220>
<221> SITE
<222> (1160)
<223> n equals a,t,g, or c
```

[illegible]

```
<210> 80
<211> 1074
<212> DNA
<213> Homo sapiens
```

```
<220>
<221> SITE
<222> (1)
<223> n equals a,t,g, or c
```

<220>
<221> SITE

<222> (2)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (11)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (33)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (694)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (972)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1053)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1055)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1058)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1060)
 <223> n equals a,t,g, or c

<400> 80
 nnggactaac ngaaaagctc ctatgacttc gcntgcacgc gcaagcttgg gccctcagag 60
 ggatcctcta gagcgccgc cctttttttt tttttttgta actctcttcc tatctttgtc 120
 atttcttttc aggccgcttc acaactcctt cttcatattg gtgttccctc tggactctgt 180
 attcagctct tacctttcca tggggaatct actctcactg ctatacacia ctgataaaaag 240
 cttgggctct ccagccagaa tgctagggtc agaatccagt tcaacgacct aacagctgtg 300
 taaaatcggt catgcaactt ttcctctctt tgcttcaatt tcctattctg tgaaataaaa 360
 gagctgtatt gaggattaga taattgatat gatacaata atgttcaata caatgtctag 420
 catgttaaaa tgcttagtaa atgggagtct ttatttctga agcagaatag tataaatagt 480
 tgagaacatg aacacagagt tcagaatgca ttgactcaaa tcaatgttct gccacttaaa 540
 agccgtggaa cctcaggcga gtgacatggc ctctctgtgc ctgagctttc tcatctataa 600
 aataagataa caactaatag ctactttgtg gtgktacagt gataattaga tgaatttaca 660
 agagtaaagt gcttggggca gcattaggta agnnttttct aagtattatc gatgctttcca 720
 atgkctyctc agagctgcat attttttcca tttgcaacaa ctgktacttt gatattttta 780
 ccagaatggt tgaagaaata aaacmcattw tgtktaagac ttatctattt atcttcttca 840

| | | | | | | |
|------------|------------|------------|------------|-------------|------------|------|
| aaaawttcta | cttttcctta | attagtaagc | atcaccattt | atctaattctt | ctaagcaaga | 900 |
| aatttcagta | actctttact | cttcactttc | ttcaagatct | cttaatagat | gcagaaaggc | 960 |
| atgtgacaaa | antcaacact | ttttcataaa | taaaaaatct | tcaacaaact | tccaaaacag | 1020 |
| ggtggagact | atcttcaaaa | ttaatggaaa | ggnctntntn | tggaaaaaac | ctca | 1074 |

<210> 81

<211> 546

<212> DNA

<213> Homo sapiens

<400> 81

| | | | | | | |
|------------|------------|------------|------------|------------|------------|-----|
| tcagctccag | ggttgcgagg | caccaccag | catcattccc | catgcgaggt | ggcaaatgca | 60 |
| acatgctctc | cagtttgagg | tgtctacttc | tctgtggaag | tattacacta | gccctgggaa | 120 |
| atgcacagaa | attgccaaaa | ggtaaaaggc | caaacctcaa | agtccacatc | aataccacaa | 180 |
| gtgactccat | cctcttgaag | ttcttgcgtc | caagtccaaa | tgtaaagctt | gaaggctctc | 240 |
| tcctgggata | tggcagcaat | gtatcaccaa | accagtactt | ccctcttccc | gctgaaggga | 300 |
| aattcacaga | agctatagtt | gatgcagagc | cgaaatatct | gatagttgtg | cgacctgctc | 360 |
| cacctccaag | tcaaaagaag | tcatgttcag | gtaaaactcg | ttctcgcaaa | cctctgcagc | 420 |
| tggtggttgg | cactctgaca | ccgagctcag | tcttctgtgc | ctgggggttc | ctcatcaacc | 480 |
| cacaccatga | ctggacattg | ccaagtcact | gtcccaatga | cagattttat | acaattcgct | 540 |
| atcgag | | | | | | 546 |

<210> 82

<211> 557

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (73)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (320)

<223> n equals a,t,g, or c

<400> 82

| | | | | | | |
|-------------|------------|------------|------------|------------|------------|-----|
| gcggccgcgc | actagtgagc | tcgtcgaccc | gggaattcgc | ggccgcgctc | actgggtggc | 60 |
| aagggtgggc | gngagtcctc | tggcttttta | gtgggggtgg | tagagwccac | tgatacattt | 120 |
| ctatcagcac | ctgatggcct | gggtgcttgc | ttgacctctg | ttcccactcc | attcccactg | 180 |
| ggcatcccac | tgggtttggg | cctagttagt | cttgtagttg | tcctattcag | aacaggtctg | 240 |
| atgtgtggct | tgtcagatgt | tctgggcctc | acagtagtat | aaaatctgtg | tggcgcctga | 300 |
| gttgtctttg | ctagttcagn | tgttcgtgga | atcttagttg | tgaaagggtc | ttgggtggat | 360 |
| tggctctgtt | cttcagaggt | ggctctgtag | tcctcttgac | taggacttgg | ggaaatcatg | 420 |
| ggtataaaaag | ggatgcctcg | tgtctctaga | ggagcaggtt | cgctctgaga | gacctcaggg | 480 |
| tatgggtttat | gaggaaggag | cacatctttt | ggagcaggtg | tgggtgatga | cacttctgga | 540 |
| cttgggtgacg | tttttgg | | | | | 557 |

<210> 83

<211> 932

<212> DNA

<213> Homo sapiens

<220>

<221> SITE
 <222> (675)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (920)
 <223> n equals a,t,g, or c

<400> 83
 gcttatctga aattgtaacc tgcctcctct ctttgggaact ctcagtcacct ctgatgttgt 60
 tctttctgta atttacttat ttattctggt tattgtgatt atctatcttg cccactcta 120
 cactagaatc acactccacg aggtcagggg ttagtggttta tttcgtttat tgacagatgc 180
 caggcattta gaatagtgcc tggcacataa taaaggctta ttaaatagat gttgaataaa 240
 tgaatcacct ctttatagtg cctgtgccct aaactagatg atatcttgtc tttgtaactt 300
 tatagcacac tgtgtagctc tagtgatgcg tacatgtatg ctggttgat cttccaactt 360
 tgctccttct ttcctttaat atataagggtc tatttaggaa ttgtcatcta gcatttggat 420
 cagtgtccag tgatgttaca tagtaggatt tatcaaattt tagaaagaga aaggaattta 480
 ccacactatc aaagaagttt aattactatg tcttattttt tgggtggaatc atatttaatg 540
 aaagagaatg tgaaattgct ttgtcttata gttcagaaac cagttttcaa aacttccttc 600
 atgtagtaaa tgggaatcttc tagtgaatgy catgaaattt ccaaattgkt tttttttaat 660
 tttttttaaa attngngtaa gtacatataa catamaattt gccaccttaa ctctttttaa 720
 gtgtacactt cattggcatt aagtacattc tcaccgttgc ataagygtta ccaccgacca 780
 gttccagaat atttttacct tgcaaacctg aaattctgta cccattaaac agtaactcct 840
 cattcctccc caacccttag caatcaaagt tttactttct gtacctatga atttgatatt 900
 ctaggtactt ttatataagn gaaatcatac ag 932

<210> 84
 <211> 661
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (2)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (615)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (640)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (655)
 <223> n equals a,t,g, or c

<400> 84
 gnggccgccc tttttttttt ttttttttta agacagagtc tcgctctggt gccaggctgg 60
 agagcagtgg tgtgatctca gtcactgca acctctacct cccagggttca agcgattctc 120
 agcctcctga gtagctggga ctacaggcgc gcaccaccat gccagctaa tttttgtatt 180
 tttagtagag acagggtttc accacgttgg ccaggatggt cttgatctct tgacctcgtg 240

| | | | | | | |
|------------|------------|------------|-------------|-------------|------------|-----|
| atccgccccg | ctcggcctac | caaagtgctg | agattacagg | cgtgagccac | cgtgccccgc | 300 |
| ctagcaggtg | cttttaaaat | atctataggt | aaatgaatga | aaacaactgc | tataaaattt | 360 |
| gggtctgtat | acctttgtcc | cagaaatact | gtttttat | atatttcat | ttttttgrga | 420 |
| wggagtctcg | ctctatcgac | caggctagag | tgcagtgggtg | cgatctcagc | ttgctgcaac | 480 |
| ctctgcctcc | tgggttcaag | cactttctct | gcctcagcgt | cctgagtagc | tgggactaca | 540 |
| ggtgagaaat | gccattttta | gaagcctggc | ttatatatgt | ggataagtga | tgaggatatt | 600 |
| tttacaagca | aatgnattat | agcacttttt | ttttttgggn | gggggttttta | agtanaaaac | 660 |
| g | | | | | | 661 |

<210> 85

<211> 2224

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (2186)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (2202)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (2218)

<223> n equals a,t,g, or c

<400> 85

| | | | | | | |
|-------------|-------------|------------|-------------|------------|------------|------|
| accacgcgt | ccggcagaag | aggagaagag | gttgtgtggg | acaagctgct | cccacagaa | 60 |
| ggatgtcgct | gctgagcctg | ccctggctgg | gcctcagacc | ggtggcaatg | tccccatggc | 120 |
| tactcctgct | gctgggttgtg | ggctcctggc | tactcgcccc | catcctggct | tggacctatg | 180 |
| ccttctataa | caactgccgc | cggctccagt | gtttcccaca | gccccaaaaa | cggaaactgg | 240 |
| tttgggggtc | cctgggcctg | atcactccta | cagaggaggg | cttgaaggac | tgcaccaga | 300 |
| tgtcggccac | ctattcccag | ggctttacgg | tatggctggg | tcccatcatc | cccttcatcg | 360 |
| ttttatgcc | ccctgacacc | atccggtcta | tcaccaatgc | ctcagctgcc | attgcacca | 420 |
| aggataatct | cttcatcagg | ttcctgaagc | cctggctggg | araaagggat | actgctgagt | 480 |
| ggcggtgaca | agtggagccg | camcgtcgg | tgtgacgcc | cgcttccat | ttcaacatyc | 540 |
| tgaagtccta | tataacgatc | ttcaacaaga | gtgcaaacat | catgcttgac | aagtggcagc | 600 |
| acctggcctc | agagggcagc | agtcgtctgg | acatgtttga | gcacatcagc | ctcatgacct | 660 |
| tggacagtct | acagaaatgc | atcttcagct | ttgacagcca | ttgtcaggag | aggcccagtg | 720 |
| aatatattgc | caccatcttg | gagctcagtg | cccttgtaga | gaaaagaagc | cagcatatcc | 780 |
| tccagcacat | ggactttctg | tattacctct | cccattgacg | gcggcgcttc | cacagggcct | 840 |
| gcgcctgggt | gcatgacttc | acagacgctg | tcattcggga | gcggcgctgc | accctcccca | 900 |
| ctcaggggat | tgatgatatt | ttcaaagaca | aagccaagtc | caagactttg | gatttcattg | 960 |
| atgtgcttct | gctgagcaag | gatgaagatg | ggaaggcatt | gtcagatgag | gatataagag | 1020 |
| cagaggctga | caccttcatg | tttgagggcc | atgacaccac | ggccagtggc | ctctcctggg | 1080 |
| tcctgtacaa | ccttgcgagg | caccagaat | accaggagcg | ctgccgacag | gaggtgcaag | 1140 |
| agcttctgaa | ggaccgcgat | cctaaagaga | ttgaatgggtg | agtgcgaagt | cttctggcct | 1200 |
| gttctctgagc | ccatcattgg | ttctgtctcc | caagttagga | ggggtggagg | agttgttttg | 1260 |
| tggattcttc | cactattgct | tagtggtgat | aaaagcagag | gaccacaggc | agggcttgat | 1320 |
| accaagcctg | gctgtcaggg | gaaaagttgc | aggccttttag | gaccgaaaga | cccgggtgc | 1380 |
| taagagaatt | ggtgacaatg | tgtraggaca | gataatgcta | ctttgcacat | gttcagcaaa | 1440 |
| tgaacttccc | ctccactctg | tttctgttct | cccagaaata | tcagtttttt | cgaaaatcct | 1500 |
| cacttcttga | atgtttgctc | aggtgggrag | ggagaagttg | ttttgtcaa | ttcttccatt | 1560 |
| attgcttagt | agtactagga | gcagagcaca | gaggcagaac | ttggtatcca | acctggctag | 1620 |

| | | | | | | |
|-------------|-------------|------------|------------|------------|------------|------|
| caggggaaag | ttttcagact | tttgggacag | agagtcctag | cccgtgaga | gaactaagga | 1680 |
| caatatatga | aggcagacga | tgacatctag | acatgttcag | tcaatgaatt | ttcccttcac | 1740 |
| tcctttcttg | tttactctga | attatcatat | ttttttcaaa | catctttact | tcctgaatgt | 1800 |
| ttactcttct | gttccctaata | tcctaccctc | cgggccagtc | cagggagtcc | taagggagac | 1860 |
| ctgggggttaa | aagtcaccta | tcttttgtct | agaacacctg | catcgttcat | ccattcagtc | 1920 |
| ttcattcagc | aaactcttcc | acatgccttt | ccattgccag | ttctgttccc | acacctttct | 1980 |
| gtkctctgga | gacctggaaa | gaacctgtcc | tatggatgtg | ttttccaggt | actcccagcc | 2040 |
| tagtaggaac | agtccccac | aagacattcc | cagtcacat | aggcaaggat | ggagttgaga | 2100 |
| gcaagagaag | catggctggg | agwacagaaa | gtgtgataga | aaatgttttc | aagccaaatc | 2160 |
| ttggttggt | gttcgcttgt | tgatanccaa | gttaccgaag | anggcttgaa | atttgggnta | 2220 |
| aatc | | | | | | 2224 |

<210> 86

<211> 1669

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (1162)

<223> n equals a,t,g, or c

<400> 86

| | | | | | | |
|-------------|------------|-------------|-------------|-------------|-------------|------|
| ctcgagtttt | tttttttttt | tttttttttt | tttgacagca | ccgtttttatt | catgacaatc | 60 |
| tgcaaaaaaa | caggtggatg | ggtcagaaaag | tcactgcaag | cytacattca | ggggctccac | 120 |
| ccgcagccaa | agcccgccct | cggcgcgcat | gatcaattcc | agcttcctgc | ggggctcagt | 180 |
| gtggtctggc | aggaaccgga | agtgcagcag | catcaacgcc | aggaccactt | tcctctccgc | 240 |
| catggcgaac | gcctgcccga | tgcaagttct | gggccctgcg | gagaaaggaa | taaaagccag | 300 |
| aggtgacctc | cccttgctgt | tctctgggtc | aaagcggaaag | gggtcgtaga | cctgcagggtg | 360 |
| agacaaagaa | ggtttgmcag | tgggattttcc | tatgtccagc | agccttgagg | agacacttgt | 420 |
| gtgtgtkktg | gggtgagggt | atgttggatt | ttcctcatca | aawccctgcc | ccctccacta | 480 |
| ggaaccttgg | aattcacaca | cacacacaca | aaagtagagg | atgggggtgg | tgggtggtgaa | 540 |
| tggggaaggc | agcacctcag | gatccggcca | cacagttggg | ttgtgatgga | cccctataat | 600 |
| atcgatgagg | caggtaatgc | ctttggggat | gactcggcca | tctgggagaa | caatgtcctg | 660 |
| gggtgcagcat | cgggagatga | agggagctgg | gggatgtaac | ctcaggctct | ccttcacgca | 720 |
| catggtcagg | aagggcagct | gggccagggtc | gtcccattca | atctcttttag | gatcgcggtc | 780 |
| cttcagaagc | tcttgcacct | cctgtcgcca | gcgtcctctg | tattctgggt | gcctcgcaag | 840 |
| gttgtacagg | accaggaga | ggccactggc | cgtggtgtca | tggcctccaa | acatgaagggt | 900 |
| gtcagcctct | gctcttatat | cctcatctga | caatgccttc | ccatcttcat | ccttgctcag | 960 |
| caaaagcaca | tcaatgaaat | ccaaagtctt | ggacttggct | ttgtctttga | aaraatcrtc | 1020 |
| aataccctra | gtrgggaggg | tgcgacgccg | stcccggwtg | acagcgtytg | tgaagtcatg | 1080 |
| caccaggcgg | caggccctgt | ggaagcgccg | cccgtcatgg | gagaggtaat | acagaaagtc | 1140 |
| catgtgctgg | aggatatgct | gncttctttt | ctctacaagg | gcactgagct | ccaagatgggt | 1200 |
| ggcaatatat | tactgggcc | tctcctgaca | atggctgtca | aagctgaaga | tgcatttctg | 1260 |
| tagactgtcc | aaggtcatga | ggctgatgtg | ctcaaacatg | tccagacgac | tgctgccttc | 1320 |
| tgaggccagg | tgtgccact | tgtcaagcat | gatgtttgca | ctcttggtga | agatcgttat | 1380 |
| ataggacttc | aggatgttga | aatggaaggc | gggcgtcagc | atccgacgggt | ggcggctcca | 1440 |
| cttgtcaccg | ccactcagca | gtatcccttc | tcctagttag | gcagccaagg | acaatgggca | 1500 |
| agagcagcac | ttcccctacc | ctccccgcaa | ggttgtcccc | aacccctttc | acctgcagggt | 1560 |
| actcaccag | ccagggtctc | aggaaacctga | tgaaragatt | atccttgggt | gcaatggcag | 1620 |
| ctgaggcatt | gggtgatagg | ggatgggggtc | aaaggcgtct | ctatcggcc | | 1669 |

<210> 87

<211> 1158

<212> DNA

<213> Homo sapiens

<220>
 <221> SITE
 <222> (668)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1020)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1043)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1096)
 <223> n equals a,t,g, or c

<400> 87
 tacgcctgca ggtaycggyc cggaattcgc ggccgcgctcg acgtacycac aatttttcagg 60
 tgggtgtgta agagctaata gggccacaga tgctatgcat accactgcct agttgttgct 120
 agagagcctt ttcaacctca gagaacatag cattccatgc taacctatcat ttaaaattcc 180
 tgcaataaat gaaaaaata aacacagagc tgagagtata attcaactag cctcgtcccc 240
 aaatgaagct gattccagat agaaagcaga ttttcatatt tactgtatgt gacagctgca 300
 ggaattaaat gaatcatctg tctatctcta ttgccttggt cctcctgtgc tgtgtgcatc 360
 tatccctggg gttgtctgtc ttcccatcc aggaagacag atcagtgtag tttcaattat 420
 taattccatg tttggatcac agtaacacca gttccagttc agtgacctgc agttyctatt 480
 gatttaagt atgaattatc tacttgcaa tgtattatta actttatatt gtctcttayt 540
 ccttctgcat tcatgcatca aaataataaa gcaccattca caggctaaaa ggacccgttt 600
 cccttctcac atttcccata agggcgaagc caacacacac caggaggagta attacacaga 660
 gctggggngg gggctggaca tttattttac ctctgaatta tttatctcag ctgtgaattt 720
 gggggagggg ttgggggagg tcctcagtgg ggaacaaagg ggcccaggag ggaagttaat 780
 gaagacaagt gacgattaga acctgggaga ttcataagaa atgaaataac ctcaacccaa 840
 aagtttaact tctgggagtc aatctgcca tctgataat gatgggggga tgggtgtagg 900
 agtggtaacg acagtggcac agtgtaagg gaaggaggca agagaagaga gcaccatcag 960
 tttctcctgc catctctttt ggattgctct gcctgcccc tcgcaagacc tactcagcan 1020
 ggtcagactc cagttagaca tgnatttgcc gatgtttgcc agaaatgtgc tggccttatg 1080
 gctaacctac cattgnettc atcttctttg ggtaaaagca catcctccta ttaccttttg 1140
 aggaacttcc tttccctg 1158

<210> 88
 <211> 963
 <212> DNA
 <213> Homo sapiens

<400> 88
 gtggccgccg actagtgagc tcgtcgaccc gggaattcgc ggccgcgctcg actgcttggt 60
 gaacaggatg acaatgtctg gttctgaaat cccaggagct gagggagagc cctttttgcc 120
 cagttcctgt tctccgaggg tttttgtga agtgagccta ggagctaaga agagagagag 180
 aaatgaggtc tgctgatccc cagcaaaact acatgcacct ccctttcatc tgttttggtt 240
 gaaaacctac gaagggtaaa atagagaaaa ccttcagggc aggcaagcta tatgtggtta 300
 gctttgtgtg ggaacctgag tcctggagac tgccctgact tttgcagcag gcacatttgt 360
 atgtttggat tttccagaag ggataaatca cacaacaga gcacctggg aatccttcag 420
 ggagtcctgt ttggccccag ccccttctg ttaaamcac accccattct ccaggctgtg 480

| | | | | | | |
|-------------|------------|------------|------------|------------|------------|-----|
| gagagtgggtt | caaacactgc | ccttgtctga | agagctttct | ctggaaaact | cattcttggg | 540 |
| gagaactttc | tttgtgtaat | agttcagcct | tttagactaa | acagacgaga | aaatagtga | 600 |
| gtacagaaaa | aaaggagttc | ctcctggtaa | acaggggaag | agttcctttg | ggtttggat | 660 |
| cctctgcctc | tctgattgtt | gcatgccatg | ttcacagtgg | taccccagcc | aagacagata | 720 |
| ctgagaaaca | gatcctcttt | tccactatgt | cttgktttct | cattgcttgg | atggtgttgc | 780 |
| aagcctgtct | ggcctccaag | tgcacatctt | gtcctcttcg | ggccccaaca | caggcctgtg | 840 |
| atatttagcc | tgcagagcag | cattcgttta | aaatcaaagc | tttgaagggt | gatctgggtt | 900 |
| gagagcaacc | ccgaagcatg | agattctttg | acatgtgggc | cccttggggc | tgtgcggaaa | 960 |
| gag | | | | | | 963 |

<210> 89

<211> 579

<212> DNA

<213> Homo sapiens

<400> 89

| | | | | | | |
|------------|------------|------------|-------------|-------------|-------------|-----|
| gcactgctca | cctgcgcctt | tcccagctct | cattttwtgtt | agggtgatttg | ggatagggat | 60 |
| agtgttttgg | ggtatggggg | gagtgtttct | gacctgcttt | gcagacgtgc | ctccgcacct | 120 |
| cagcagtttg | gggtgtggcc | ccagggcggt | tcttggtatg | aaaagatgtg | gccatctagc | 180 |
| ctcgtaaact | cactgtcacc | tgtgtcccat | agggtgcctt | ctgaatactg | ttattagaat | 240 |
| aagtttgttg | cagaacgtga | ccctgcgtgc | aaacatgtac | cgtggcctgg | tatatgatag | 300 |
| agattgatat | taatgtacca | tgtatgttaa | tgtgaatctg | tgggcaggat | acttttccat | 360 |
| ggcaggaaat | atccaagctg | ttgaaactgg | ctatgtttta | atatgcctca | ttgtgccttt | 420 |
| actgttgtgt | ggactgcgtg | agggacaaga | agttccattt | gatgtcaata | aagcaaaagta | 480 |
| cttgcttact | tttttgaaaa | aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | 540 |
| aaaaaaaaaa | aaaaaaaaaa | aaaaaaaaaa | aaaaataaaa | | | 579 |

<210> 90

<211> 1316

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (157)

<223> n-equals a,t,g, or c

<400> 90

| | | | | | | |
|------------|-------------|-------------|------------|------------|-------------|------|
| cctgcttctc | atataagatg | ggcctgcacc | tacctctggc | atgtttttct | tttgttattt | 60 |
| cctgtttttc | atcctgtaac | taatgctcat | tatttaaaac | actccagtta | cttttccctt | 120 |
| taggcctggc | aaacttttct | ttttcttttt | ttttttnata | actgtacatt | ttacaaaaca | 180 |
| cctgaaagga | cccagaggca | taaatagtat | tacatgattt | atggtagctc | tggggtgagc | 240 |
| tgtgggggag | gccagttaca | aggtctccaa | gacccgcctc | catgtcagtg | ttgggggggtg | 300 |
| ccttgagcac | ttcttttatt | tcatatagga | aaatgcacat | ctttcatttg | tctcccaaaa | 360 |
| gatgatttcc | ataaaaaatat | acttcaaagt | atttactgta | catagacttc | aatgatttta | 420 |
| tttccagcat | caaactaaac | atttctaacc | ttttaaataa | atggggaatt | tagtttcatg | 480 |
| agaattttta | ctgacaacta | ataaactctg | tatgtgtaca | tattcatata | cactgtggac | 540 |
| tggagaaagt | cagtatggaa | ataaatttca | ctgagatcca | gaaagtagca | atataggact | 600 |
| tttaggggta | tgtattagtg | ttaaataatat | atgtgcagtc | taggctaata | attttaatta | 660 |
| aaaacaaata | agcatacttt | tttgctttta | aaatatgccc | tcaaaatggg | gggggataaa | 720 |
| atgtaagata | ttcttttagct | ttaattattt | ttccatctta | ttcaagtcag | agcacttttt | 780 |
| tttaaatagc | aatacaacag | gcaagagaat | actcaaaaat | atttaaaact | gtattgatag | 840 |
| caagagtatg | ttttaaatat | tttctaataa | atacttgagc | ggtttttgtc | tggcaggcctt | 900 |
| ccaaatttgc | caaaattaag | cgttcagtat | tttcaacaca | tacgcttttt | actggtttat | 960 |
| actgaactat | ctgatgagaa | ttcctgtgtt | cccaaagcaa | ctgatgttta | caggctctgt | 1020 |
| gtttctcttc | ctcctttcta | aggatgaggg | aatccacaac | agactttctc | tagaaaacac | 1080 |

69

```

taatgatgga caactttttg gtgtcatcaa tgagttggct actaccttga tgtaaaaatt    1140
tgtaaggaaa attttcacca ttctgagtggt caagtgtatt ttttaactgtc tggtttgtac    1200
ttttatgact tttgtactac caaagcggag ttaaaaataa aaatgtttaa cgcaaaaaaa    1260
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaa    1316

```

<210> 91

<211> 531

<212> PRT

<213> Homo sapiens

<400> 91

```

Met Leu Cys Ala Leu Leu Leu Leu Pro Ser Leu Leu Gly Ala Thr Arg
 1          5          10          15

Ala Ser Pro Thr Ser Gly Pro Gln Glu Cys Ala Lys Gly Ser Thr Val
          20          25          30

Trp Cys Gln Asp Leu Gln Thr Ala Ala Arg Cys Gly Ala Val Gly Tyr
          35          40          45

Cys Gln Gly Ala Val Trp Asn Lys Pro Thr Ala Lys Ser Leu Pro Cys
          50          55          60

Asp Val Cys Gln Asp Ile Ala Ala Ala Ala Gly Asn Gly Leu Asn Pro
 65          70          75          80

Asp Ala Thr Glu Ser Asp Ile Leu Ala Leu Val Met Lys Thr Cys Glu
          85          90          95

Trp Leu Pro Ser Gln Glu Ser Ser Ala Gly Cys Lys Trp Met Val Asp
          100          105          110

Ala His Ser Ser Ala Ile Leu Ser Met Leu Arg Gly Ala Pro Asp Ser
          115          120          125

Ala Pro Ala Gln Val Cys Thr Ala Leu Ser Leu Cys Glu Pro Leu Gln
          130          135          140

Arg His Leu Ala Thr Leu Arg Pro Leu Ser Lys Glu Asp Thr Phe Glu
          145          150          155          160

Ala Val Ala Pro Phe Met Ala Asn Gly Pro Leu Thr Phe His Pro Arg
          165          170          175

Gln Ala Pro Glu Gly Ala Leu Cys Gln Asp Cys Val Arg Gln Val Ser
          180          185          190

Arg Leu Gln Glu Ala Val Arg Ser Asn Leu Thr Leu Ala Asp Leu Asn
          195          200          205

Ile Gln Glu Gln Cys Glu Ser Leu Gly Pro Gly Leu Ala Val Leu Cys
          210          215          220

Lys Asn Tyr Leu Phe Gln Phe Phe Val Pro Ala Asp Gln Ala Leu Arg
          225          230          235          240

Leu Leu Pro Pro Gln Glu Leu Cys Arg Lys Gly Gly Phe Cys Glu Glu

```


70

| | | | | | |
|---|-----|-----|-----|--|-----|
| | 245 | | 250 | | 255 |
| Leu Gly Ala Pro Ala Arg Leu Thr Gln Val Val Ala Met Asp Gly Val | 260 | 265 | 270 | | |
| Pro Ser Leu Glu Leu Gly Leu Pro Arg Lys Gln Ser Glu Met Gln Met | 275 | 280 | 285 | | |
| Lys Ala Gly Val Thr Cys Glu Val Cys Met Asn Val Val Gln Lys Leu | 290 | 295 | 300 | | |
| Asp His Trp Leu Met Ser Asn Ser Ser Glu Leu Met Ile Thr His Ala | 305 | 310 | 315 | | 320 |
| Leu Glu Arg Val Cys Ser Val Met Pro Ala Ser Ile Thr Lys Glu Cys | 325 | 330 | 335 | | |
| Ile Ile Leu Val Asp Thr Tyr Ser Pro Ser Leu Val Gln Leu Val Ala | 340 | 345 | 350 | | |
| Lys Ile Thr Pro Glu Lys Val Cys Lys Phe Ile Arg Leu Cys Gly Asn | 355 | 360 | 365 | | |
| Arg Arg Arg Ala Arg Ala Val His Asp Ala Tyr Ala Ile Val Pro Ser | 370 | 375 | 380 | | |
| Pro Glu Trp Asp Ala Glu Asn Gln Gly Ser Phe Cys Asn Gly Cys Lys | 385 | 390 | 395 | | 400 |
| Arg Leu Leu Thr Val Ser Ser His Asn Leu Glu Ser Lys Ser Thr Lys | 405 | 410 | 415 | | |
| Arg Asp Ile Leu Val Ala Phe Lys Gly Gly Cys Ser Ile Leu Pro Leu | 420 | 425 | 430 | | |
| Pro Tyr Met Ile Gln Cys Lys His Phe Val Thr Gln Tyr Glu Pro Val | 435 | 440 | 445 | | |
| Leu Ile Glu Ser Leu Lys Asp Met Met Asp Pro Val Ala Val Cys Lys | 450 | 455 | 460 | | |
| Lys Val Gly Ala Cys His Gly Pro Arg Thr Pro Leu Leu Gly Thr Asp | 465 | 470 | 475 | | 480 |
| Gln Cys Ala Leu Gly Pro Ser Phe Trp Cys Arg Ser Gln Glu Ala Ala | 485 | 490 | 495 | | |
| Ser Cys Ala Thr Leu Cys Asn Thr Ala Arg Ser Met Tyr Gly Lys Arg | 500 | 505 | 510 | | |
| Cys Thr Ser Thr Leu Gly Asn Thr Arg Asp Arg Gly Cys Gln Arg Pro | 515 | 520 | 525 | | |
| Arg Ala Cys | 530 | | | | |

71

<211> 186

<212> PRT

<213> Homo sapiens

<400> 92

Met Leu Pro Leu Val Ser Gly Cys Pro Gly Phe Ser Leu Gly Cys Phe
 1 5 10 15

Phe Phe Leu Thr Ala Leu His Ile Tyr Ala Asn Tyr Arg Ala Val Arg
 20 25 30

Ala Leu Val Met Glu Thr Leu Asn Glu Gly Arg Leu Arg Leu Val Leu
 35 40 45

Lys His Tyr Leu Gln Arg Gly Glu Val Leu Asp Pro Thr Ala Ala Asn
 50 55 60

Arg Met Glu Pro Leu Trp Thr Gly Phe Trp Pro Ala Pro Ser Leu Ser
 65 70 75 80

Leu Gly Val Pro Leu His Arg Leu Val Ser Ser Val Phe Glu Leu Gln
 85 90 95

Gln Leu Val Glu Gly His Gln Glu Ser Tyr Leu Leu Cys Trp Asp Gln
 100 105 110

Ser Gln Asn Gln Val Gln Val Val Leu Asn Gln Lys Ala Gly Pro Lys
 115 120 125

Thr Ile Leu Arg Ala Ala Thr His Gly Leu Met Leu Gly Ala Leu Gln
 130 135 140

Gly Asp Gly Pro Leu Pro Ala Glu Leu Glu Glu Leu Arg Asn Arg Val
 145 150 155 160

Arg Ala Gly Pro Arg Lys Arg Ala Gly Ser Ser Ser Arg Arg His Thr
 165 170 175

Lys Cys Trp Thr Cys Cys Ser Gln Ser Ser
 180 185

<210> 93

<211> 96

<212> PRT

<213> Homo sapiens

<400> 93

Met Ala Arg Ser Leu Leu Ile Ile Leu Gly Ala Asp Phe Thr Phe Pro
 1 5 10 15

Thr Ser Phe Asn Cys Phe Gln Lys Met Asn Leu Ala Lys Lys Ser Arg
 20 25 30

Gly Ser Phe Thr His Leu Leu Thr His Ser Trp Cys Leu Ser Leu Phe
 35 40 45

Leu Lys Glu Ala Asp Gln Gly Leu Arg Glu Asn Asn Phe Asp Phe Ser

72

50 55 60
 His Val Cys Pro Ser Lys Pro Pro Leu Trp Thr Asp Ser Pro Ser Val
 65 70 75 80
 Pro Gly Arg Asn Trp Asp Asn Pro Arg Thr Phe Leu Val Pro Ser Arg
 85 90 95

<210> 94
 <211> 77
 <212> PRT
 <213> Homo sapiens

<400> 94
 Met Leu Leu Phe Phe Phe Phe Leu Leu Phe Phe Phe Phe Phe Phe Trp
 1 5 10 15
 Leu Val Leu Phe Gly Ile Phe Phe Phe Ser Phe Leu Lys Lys Met Phe
 20 25 30
 Ser Gly Asn Met Asn Lys His Thr Ala Asn Tyr Ser Gly Ala Gly Lys
 35 40 45
 Ala Gln Glu Leu Ala Thr Ser Gln Leu His Ser Trp Asp Gly Lys Pro
 50 55 60
 Cys Cys Glu Leu Leu Arg Leu Phe Thr Tyr Phe Thr Tyr
 65 70 75

<210> 95
 <211> 40
 <212> PRT
 <213> Homo sapiens

<400> 95
 Met Arg Arg Gln Thr Phe Met Ser Ile Leu Val Phe Gln Cys Ser Pro
 1 5 10 15
 Ile Ser Phe Gly Leu Cys Ile Asn Lys Glu Arg Thr Val Val Ser Ser
 20 25 30
 Val Ile Thr Asp Asn Leu Cys Leu
 35 40

<210> 96
 <211> 58
 <212> PRT
 <213> Homo sapiens

<400> 96
 Met Lys Leu Thr Phe Ser Phe Pro Trp Phe Thr Leu Thr Ala Leu Gln
 1 5 10 15

Leu Trp Ser Ala Thr Glu Cys Gln Ala Val Val Asp Thr Met Ile Ala
 20 25 30

Val Trp Ser Glu Gly Lys Gly Thr Gly Val Ser Trp Glu Pro Trp Leu
 35 40 45

Leu Gly Lys Leu Gln Ser Ser Ser Phe Leu
 50 55

<210> 97
 <211> 127
 <212> PRT
 <213> Homo sapiens

<400> 97
 Met Pro Arg Ala Pro Trp Arg Ile Pro Leu Cys Ala Leu Pro Thr Leu
 1 5 10 15

Cys Leu Gly Ser Pro Leu Pro Ser Gln Pro Thr His Pro Ile Phe Tyr
 20 25 30

Asp His Arg Ala Pro Thr Trp Lys Met Ala His Pro Gly Gly Pro Arg
 35 40 45

Ser Ser His Ser Pro Arg Gly Pro Gly Gly His Pro Ala Leu Arg Gln
 50 55 60

Arg Leu Pro Cys Arg Arg Gly Glu Pro Glu Thr Ala Leu Cys Ser Ser
 65 70 75 80

Ala Pro Gly Ala Gly Phe Ala Glu Pro Pro Cys Lys Ala Ser Pro Gly
 85 90 95

Trp Gly Pro Pro Ser Arg Gly Pro Gln Gly Asp Arg Ser Gln Gly Glu
 100 105 110

Trp Leu Pro Ala Leu Gly Thr Pro Cys Gly Gly Pro Asp Asp Ser
 115 120 125

<210> 98
 <211> 86
 <212> PRT
 <213> Homo sapiens

<400> 98
 Met Ser Leu Cys Arg Ile Leu Gly Tyr Ser Phe Ser Ser Arg Leu Ser
 1 5 10 15

Ser Leu Ile Leu Pro Leu Ala Val Phe His Tyr Cys Leu Ser Cys Pro
 20 25 30

Leu His Phe Lys Leu Ser Phe Lys Tyr Leu Pro Phe Pro Ser Phe Pro
 35 40 45

Phe Ser Ser Leu Pro Cys Pro Ala Leu Pro Cys Pro Ala Leu Pro Ser

74

50 55 60
 Pro Pro Leu Pro Cys Pro Pro Leu Pro Ser Pro Pro Leu Pro Leu Pro
 65 70 75 80
 Ser Leu Ser Phe Phe Arg
 85

 <210> 99
 <211> 23
 <212> PRT
 <213> Homo sapiens

 <400> 99
 Met Phe Leu Lys Ala Gln Trp Leu Tyr Ser Leu Leu Leu Asn Cys Leu
 1 5 10 15
 Leu Pro Glu Gly Thr Ser Ser
 20

 <210> 100
 <211> 822
 <212> PRT
 <213> Homo sapiens

 <400> 100
 Met Ala Ala Ala Val Val Val Ala Glu Gly Asp Ser Asp Ser Arg Pro
 1 5 10 15
 Gly Gln Glu Leu Leu Val Ala Trp Asn Thr Val Ser Thr Gly Leu Val
 20 25 30
 Pro Pro Ala Ala Leu Gly Leu Val Ser Ser Arg Thr Ser Gly Ala Val
 35 40 45
 Pro Pro Lys Glu Glu Glu Leu Arg Ala Ala Val Glu Val Leu Arg Gly
 50 55 60
 His Gly Leu His Ser Val Leu Glu Glu Trp Phe Val Glu Val Leu Gln
 65 70 75 80
 Asn Asp Leu Gln Ala Asn Ile Ser Pro Glu Phe Trp Asn Ala Ile Ser
 85 90 95
 Gln Cys Glu Asn Ser Ala Asp Glu Pro Gln Cys Leu Leu Leu Leu Leu
 100 105 110
 Asp Ala Phe Gly Leu Leu Glu Ser Arg Leu Asp Pro Tyr Leu Arg Ser
 115 120 125
 Leu Glu Leu Leu Glu Lys Trp Thr Arg Leu Gly Leu Leu Met Gly Thr
 130 135 140
 Gly Ala Gln Gly Leu Arg Glu Glu Val His Thr Met Leu Arg Gly Val
 145 150 155 160

Leu Phe Phe Ser Thr Pro Arg Thr Phe Gln Glu Met Ile Gln Arg Leu
 165 170 175
 Tyr Gly Cys Phe Leu Arg Val Tyr Met Gln Ser Lys Arg Lys Gly Glu
 180 185 190
 Gly Gly Thr Asp Pro Glu Leu Glu Gly Glu Leu Asp Ser Arg Tyr Ala
 195 200 205
 Arg Arg Arg Tyr Tyr Arg Leu Leu Gln Ser Pro Leu Cys Ala Gly Cys
 210 215 220
 Ser Ser Asp Lys Gln Gln Cys Trp Cys Arg Gln Ala Leu Glu Gln Phe
 225 230 235 240
 His Gln Leu Ser Gln Val Leu His Arg Leu Ser Leu Leu Glu Arg Val
 245 250 255
 Ser Ala Glu Ala Val Thr Thr Thr Leu His Gln Val Thr Arg Glu Arg
 260 265 270
 Met Glu Asp Arg Cys Arg Gly Glu Tyr Glu Arg Ser Phe Leu Arg Glu
 275 280 285
 Phe His Lys Trp Ile Glu Arg Val Val Gly Trp Leu Gly Lys Val Phe
 290 295 300
 Leu Gln Asp Gly Pro Ala Arg Pro Ala Ser Pro Glu Ala Gly Asn Thr
 305 310 315 320
 Leu Arg Arg Trp Arg Cys His Val Gln Arg Phe Phe Tyr Arg Ile Tyr
 325 330 335
 Ala Ser Leu Arg Ile Glu Glu Leu Phe Ser Ile Val Arg Asp Phe Pro
 340 345 350
 Asp Ser Arg Pro Ala Ile Glu Asp Leu Lys Tyr Cys Leu Glu Arg Thr
 355 360 365
 Asp Gln Arg Gln Gln Leu Leu Val Ser Leu Lys Ala Ala Leu Glu Thr
 370 375 380
 Arg Leu Leu His Pro Gly Val Asn Thr Cys Asp Ile Ile Thr Leu Tyr
 385 390 395 400
 Ile Ser Ala Ile Lys Ala Leu Arg Val Leu Asp Pro Ser Met Val Ile
 405 410 415
 Leu Glu Val Ala Cys Glu Pro Ile Arg Arg Tyr Leu Arg Thr Arg Glu
 420 425 430
 Asp Thr Val Arg Gln Ile Val Ala Gly Leu Thr Gly Asp Ser Asp Gly
 435 440 445
 Thr Gly Asp Leu Ala Val Glu Leu Ser Lys Thr Asp Pro Ala Ser Leu
 450 455 460
 Glu Thr Gly Gln Asp Ser Glu Asp Asp Ser Gly Glu Pro Glu Asp Trp

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 465 | | 470 | | 475 | | 480 | | | | | | | | | |
| Val | Pro | Asp | Pro | Val | Asp | Ala | Asp | Pro | Gly | Lys | Ser | Ser | Ser | Lys | Arg |
| | | | | 485 | | | | | 490 | | | | | 495 | |
| Arg | Ser | Ser | Asp | Ile | Ile | Ser | Leu | Leu | Val | Ser | Ile | Tyr | Gly | Ser | Lys |
| | | | 500 | | | | | 505 | | | | | 510 | | |
| Asp | Leu | Phe | Ile | Asn | Glu | Tyr | Arg | Ser | Leu | Leu | Ala | Asp | Arg | Leu | Leu |
| | | 515 | | | | | 520 | | | | | 525 | | | |
| His | Gln | Phe | Ser | Phe | Ser | Pro | Glu | Arg | Glu | Ile | Arg | Asn | Val | Glu | Leu |
| | 530 | | | | | 535 | | | | | 540 | | | | |
| Leu | Lys | Leu | Arg | Phe | Gly | Glu | Ala | Pro | Met | His | Phe | Cys | Glu | Val | Met |
| 545 | | | | | 550 | | | | | 555 | | | | | 560 |
| Leu | Lys | Asp | Met | Ala | Asp | Ser | Arg | Arg | Ile | Asn | Ala | Asn | Ile | Arg | Glu |
| | | | | 565 | | | | | 570 | | | | | 575 | |
| Glu | Asp | Glu | Lys | Arg | Pro | Ala | Glu | Glu | Gln | Pro | Pro | Phe | Gly | Val | Tyr |
| | | | 580 | | | | | 585 | | | | | 590 | | |
| Ala | Val | Ile | Leu | Ser | Ser | Glu | Phe | Trp | Pro | Pro | Phe | Lys | Asp | Glu | Lys |
| | | 595 | | | | | 600 | | | | | 605 | | | |
| Leu | Glu | Val | Pro | Glu | Asp | Ile | Arg | Ala | Ala | Leu | Glu | Ala | Tyr | Cys | Lys |
| | 610 | | | | | 615 | | | | | 620 | | | | |
| Lys | Tyr | Glu | Gln | Leu | Lys | Ala | Met | Arg | Thr | Leu | Ser | Trp | Lys | His | Thr |
| 625 | | | | 630 | | | | | | 635 | | | | | 640 |
| Leu | Gly | Leu | Val | Thr | Met | Asp | Val | Glu | Leu | Ala | Asp | Arg | Thr | Leu | Ser |
| | | | | 645 | | | | | 650 | | | | | 655 | |
| Val | Ala | Val | Thr | Pro | Val | Gln | Ala | Val | Ile | Leu | Leu | Tyr | Phe | Gln | Asp |
| | | | 660 | | | | | 665 | | | | | 670 | | |
| Gln | Ala | Ser | Trp | Thr | Leu | Glu | Glu | Leu | Ser | Lys | Ala | Val | Lys | Met | Pro |
| | | 675 | | | | | 680 | | | | | 685 | | | |
| Val | Ala | Leu | Leu | Arg | Arg | Arg | Met | Ser | Val | Trp | Leu | Gln | Gln | Gly | Val |
| | | 690 | | | | 695 | | | | | 700 | | | | |
| Leu | Arg | Glu | Glu | Pro | Pro | Gly | Thr | Phe | Ser | Val | Ile | Glu | Glu | Glu | Arg |
| 705 | | | | 710 | | | | | | 715 | | | | | 720 |
| Pro | Gln | Asp | Arg | Asp | Asn | Met | Val | Leu | Ile | Asp | Ser | Asp | Asp | Glu | Ser |
| | | | | 725 | | | | | 730 | | | | | 735 | |
| Asp | Ser | Gly | Met | Ala | Ser | Gln | Ala | Asp | Gln | Lys | Glu | Glu | Glu | Leu | Leu |
| | | | 740 | | | | | 745 | | | | | 750 | | |
| Leu | Phe | Trp | Thr | Tyr | Ile | Gln | Ala | Met | Leu | Thr | Asn | Leu | Glu | Ser | Leu |
| | | 755 | | | | 760 | | | | | | 765 | | | |
| Ser | Leu | Asp | Arg | Ile | Tyr | Asn | Met | Leu | Arg | Met | Phe | Val | Val | Thr | Gly |
| | | 770 | | | | 775 | | | | | 780 | | | | |

Pro Ala Leu Ala Glu Ile Asp Leu Gln Glu Leu Gln Gly Tyr Leu Gln
785 790 795 800

Lys Lys Val Arg Asp Gln Gln Leu Val Tyr Ser Ala Gly Val Tyr Arg
805 810 815

Leu Pro Lys Asn Cys Ser
820

<210> 101
<211> 302
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (262)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (279)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (294)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (295)
<223> Xaa equals any of the naturally occurring L-amino acids

<400> 101
Met Leu Leu Leu Trp Lys Asn Phe Met Tyr Arg Arg Arg Gln Pro Val
1 5 10 15

Gln Leu Leu Val Glu Leu Leu Trp Pro Leu Phe Leu Phe Phe Ile Leu
20 25 30

Val Ala Val Arg His Ser His Pro Pro Leu Glu His His Glu Cys His
35 40 45

Phe Pro Asn Lys Pro Leu Pro Ser Ala Gly Thr Val Pro Trp Leu Gln
50 55 60

Gly Leu Ile Cys Asn Val Asn Asn Thr Cys Phe Pro Gln Leu Thr Pro
65 70 75 80

Gly Glu Glu Pro Gly Arg Leu Ser Asn Phe Asn Asp Ser Leu Val Ser
85 90 95

Arg Leu Leu Ala Asp Ala Arg Thr Val Leu Gly Gly Ala Ser Ala His
100 105 110

78

Arg Thr Leu Ala Gly Leu Gly Lys Leu Ile Ala Thr Leu Arg Ala Ala
 115 120 125
 Arg Ser Thr Ala Gln Pro Gln Pro Thr Lys Gln Ser Pro Leu Glu Pro
 130 135 140
 Pro Met Leu Asp Val Ala Glu Leu Leu Thr Ser Leu Leu Arg Thr Glu
 145 150 155 160
 Ser Leu Gly Leu Ala Leu Gly Gln Ala Gln Glu Pro Leu His Ser Leu
 165 170 175
 Leu Glu Ala Ala Glu Asp Leu Ala Gln Glu Leu Leu Ala Leu Arg Ser
 180 185 190
 Leu Val Glu Leu Arg Ala Leu Leu Gln Arg Pro Arg Gly Thr Ser Gly
 195 200 205
 Pro Leu Glu Leu Leu Ser Glu Ala Leu Cys Ser Val Arg Gly Pro Ser
 210 215 220
 Ser Thr Val Gly Pro Ser Leu Asn Trp Tyr Glu Ala Ser Asp Leu Met
 225 230 235 240
 Glu Leu Val Gly Gln Glu Pro Glu Ser Ala Cys Arg Gln Gln Leu Ser
 245 250 255
 Pro Leu Leu Gly Ala Xaa Trp Ser Leu Asp Ser Thr Arg Cys Pro Leu
 260 265 270
 Val Trp Asn Ala Glu Ala Xaa Ser Ser Glu Val Leu Leu Thr Asp His
 275 280 285
 Phe Thr Glu Val Met Xaa Xaa Glu Arg Leu Gln Ser Tyr Leu
 290 295 300

<210> 102

<211> 39

<212> PRT

<213> Homo sapiens

<400> 102

Met Ala Gly Gln Phe Arg Ser Tyr Val Trp Asp Pro Leu Leu Ile Leu
 1 5 10 15

Ser Gln Ser Ser Ser Cys Arg Pro Cys Ile Thr Ala Arg Trp Ala Cys
 20 25 30

Gly Trp Arg Trp Trp Thr Gly
 35

<210> 103

<211> 802

<212> PRT

<213> Homo sapiens

<400> 103

```

Met Leu Gly Ala Arg Ala Trp Leu Gly Arg Val Leu Leu Leu Pro Arg
 1              5              10              15

Ala Gly Ala Gly Leu Ala Ala Ser Arg Arg Cys Pro Gly Val Trp Pro
      20              25              30

Arg Thr Trp Pro His Arg Ser Pro Ser Arg Gly Ser Ser Ser Arg Asp
      35              40              45

Lys Asp Arg Ser Ala Thr Val Ser Ser Ser Val Pro Met Pro Ala Gly
      50              55              60

Gly Lys Gly Ser His Pro Ser Ser Thr Pro Gln Arg Val Pro Asn Arg
      65              70              75              80

Leu Ile His Glu Lys Ser Pro Tyr Leu Leu Gln His Ala Tyr Asn Pro
      85              90              95

Val Asp Trp Tyr Pro Trp Gly Gln Glu Ala Phe Asp Lys Ala Arg Lys
      100             105             110

Glu Asn Lys Pro Ile Phe Leu Ser Val Gly Tyr Ser Thr Cys His Trp
      115             120             125

Cys His Met Met Glu Glu Glu Ser Phe Gln Asn Glu Glu Ile Gly Arg
      130             135             140

Leu Leu Ser Glu Asp Phe Val Ser Val Lys Val Asp Arg Glu Glu Arg
      145             150             155             160

Pro Asp Val Asp Lys Val Tyr Met Thr Phe Val Gln Ala Thr Ser Ser
      165             170             175

Gly Gly Gly Trp Pro Met Asn Val Trp Leu Thr Pro Asn Leu Gln Pro
      180             185             190

Phe Val Gly Gly Thr Tyr Phe Pro Pro Glu Asp Gly Leu Thr Arg Val
      195             200             205

Gly Phe Arg Thr Val Leu Leu Arg Ile Arg Glu Gln Trp Lys Gln Asn
      210             215             220

Lys Asn Thr Leu Leu Glu Asn Ser Gln Arg Val Thr Thr Ala Leu Leu
      225             230             235             240

Ala Arg Ser Glu Ile Ser Val Gly Asp Arg Gln Leu Pro Pro Ser Ala
      245             250             255

Ala Thr Val Asn Asn Arg Cys Phe Gln Gln Leu Asp Glu Gly Tyr Asp
      260             265             270

Glu Glu Tyr Gly Gly Phe Ala Glu Ala Pro Lys Phe Pro Thr Pro Val
      275             280             285

Ile Leu Ser Phe Leu Phe Ser Tyr Trp Leu Ser His Arg Leu Thr Gln
      290             295             300

```

80

Asp Gly Ser Arg Ala Gln Gln Met Ala Leu His Thr Leu Lys Met Met
 305 310 315 320
 Ala Asn Gly Gly Ile Arg Asp His Val Gly Gln Gly Phe His Arg Tyr
 325 330 335
 Ser Thr Asp Arg Gln Trp His Val Pro His Phe Glu Lys Met Leu Tyr
 340 345 350
 Asp Gln Ala Gln Leu Ala Val Ala Tyr Ser Gln Ala Phe Gln Leu Ser
 355 360 365
 Gly Asp Glu Phe Tyr Ser Asp Val Ala Lys Gly Ile Leu Gln Tyr Val
 370 375 380
 Ala Arg Ser Leu Ser His Arg Ser Gly Gly Phe Tyr Ser Ala Glu Asp
 385 390 395 400
 Ala Asp Ser Pro Pro Glu Arg Gly Gln Arg Pro Lys Glu Gly Ala Tyr
 405 410 415
 Tyr Val Trp Thr Val Lys Glu Val Gln Gln Leu Leu Pro Glu Pro Val
 420 425 430
 Leu Gly Ala Thr Glu Pro Leu Thr Ser Gly Gln Leu Leu Met Lys His
 435 440 445
 Tyr Gly Leu Thr Glu Ala Gly Asn Ile Ser Pro Ser Gln Asp Pro Lys
 450 455 460
 Gly Glu Leu Gln Gly Gln Asn Val Leu Thr Val Arg Tyr Ser Leu Glu
 465 470 475 480
 Leu Thr Ala Ala Arg Phe Gly Leu Asp Val Glu Ala Val Arg Thr Leu
 485 490 495
 Leu Asn Ser Gly Leu Glu Lys Leu Phe Gln Ala Arg Lys His Arg Pro
 500 505 510
 Lys Pro His Leu Asp Ser Lys Met Leu Ala Ala Trp Asn Gly Leu Met
 515 520 525
 Val Ser Gly Tyr Ala Val Thr Gly Ala Val Leu Gly Gln Asp Arg Leu
 530 535 540
 Ile Asn Tyr Ala Thr Asn Gly Ala Lys Phe Leu Lys Arg His Met Phe
 545 550 555 560
 Asp Val Ala Ser Gly Arg Leu Met Arg Thr Cys Tyr Thr Gly Pro Gly
 565 570 575
 Gly Thr Val Glu His Ser Asn Pro Pro Cys Trp Gly Phe Leu Glu Asp
 580 585 590
 Tyr Ala Phe Val Val Arg Gly Leu Leu Asp Leu Tyr Glu Ala Ser Gln
 595 600 605
 Glu Ser Ala Trp Leu Glu Trp Ala Leu Arg Leu Gln Asp Thr Gln Asp

| 610 | 615 | 620 |
|--|-----|-----|
| Arg Leu Phe Trp Asp Ser Gln Gly Gly Gly Tyr Phe Cys Ser Glu Ala 625 630 635 640 | | |
| Glu Leu Gly Ala Gly Leu Pro Leu Arg Leu Lys Asp Asp Gln Asp Gly 645 650 655 | | |
| Ala Glu Pro Ser Ala Asn Ser Val Ser Ala His Asn Leu Leu Arg Leu 660 665 670 | | |
| His Gly Phe Thr Gly His Lys Asp Trp Met Asp Lys Cys Val Cys Leu 675 680 685 | | |
| Leu Thr Ala Phe Ser Glu Arg Met Arg Arg Val Pro Val Ala Leu Pro 690 695 700 | | |
| Glu Met Val Arg Ala Leu Ser Ala Gln Gln Gln Thr Leu Lys Gln Ile 705 710 715 720 | | |
| Val Ile Cys Gly Asp Arg Gln Ala Lys Asp Thr Lys Ala Leu Val Gln 725 730 735 | | |
| Cys Val His Ser Val Tyr Ile Pro Asn Lys Val Leu Ile Leu Ala Asp 740 745 750 | | |
| Gly Asp Pro Ser Ser Phe Leu Ser Arg Gln Leu Pro Phe Leu Ser Thr 755 760 765 | | |
| Leu Arg Arg Leu Glu Asp Gln Ala Thr Ala Tyr Val Cys Glu Asn Gln 770 775 780 | | |
| Ala Cys Ser Val Pro Ile Thr Asp Pro Cys Glu Leu Arg Lys Leu Leu 785 790 795 800 | | |
| His Pro | | |

<210> 104

<211> 89

<212> PRT

<213> Homo sapiens

<400> 104

| |
|---|
| Met Lys Pro Ala Thr Ala Ser Ala Leu Leu Leu Leu Leu Gly Leu 1 5 10 15 |
| Ala Trp Thr Gln Gly Ser His Gly Trp Gly Ala Asp Ala Ser Ser Leu 20 25 30 |
| Gln Lys Arg Ala Gly Arg Ala Asp Gln Val Ser Leu Cys Pro Gln Val 35 40 45 |
| Thr Leu Gln Gly Pro Trp Ser Pro Leu Ala Leu Leu Pro Gly Leu Gly 50 55 60 |
| Asn Leu Lys Phe Ser Phe Thr Pro Pro Phe Asn Gly Phe Leu Ser Arg |

82

65 70 75 80

Val Gln Asp Gly Arg Arg Trp Gln Leu
85

<210> 105
<211> 79
<212> PRT
<213> Homo sapiens

<400> 105
Met Arg Lys Trp Gly Leu Met Lys Leu Ile Ala Ser Met Met Gln Pro
1 5 10 15
Val Leu Leu Glu Leu Leu Ser Val Trp Arg Lys Glu Gly Arg Asp Ser
20 25 30
Arg Asn Ile His Asp Ser His Ser Met Tyr Val Leu Arg Lys Arg Leu
35 40 45
Ser Gly Ser Trp Leu Gln Ala Gly Leu Tyr Ser Thr Val Ile Ser Ala
50 55 60
Ala Leu Ile Leu Glu Ser Pro Arg Ala Cys Leu Pro Ser Lys Gly
65 70 75

<210> 106
<211> 98
<212> PRT
<213> Homo sapiens

<400> 106
Met His Cys Cys Gln Leu Pro Trp Arg Cys Ala Gln Ala Pro Gln Glu
1 5 10 15
Ala Phe Leu Leu Cys Leu Leu Phe Leu Ile Leu Val Leu Val Leu Leu
20 25 30
Gly Cys Ser Arg Gly Leu Pro Gly His Thr Pro Trp Arg Leu His Pro
35 40 45
Ala Ala Ala Ala Leu Leu Ala Pro Leu Leu His Asp Ala Leu Gly Ala
50 55 60
Cys Gly Phe Gln Gly Pro Glu Tyr Leu Leu Pro Cys Leu Leu Pro Leu
65 70 75 80
Pro Lys Pro Gly Gln Leu Gln Gly Pro Trp Gly Pro Leu Trp Ala Leu
85 90 95
Leu Pro

<210> 107
<211> 1887

<212> PRT

<213> Homo sapiens

<400> 107

```

Met Ala Ala Arg Gly Arg Gly Leu Leu Leu Leu Thr Leu Ser Val Leu
  1           5           10           15

Leu Ala Ala Gly Pro Ser Ala Ala Ala Lys Leu Asn Ile Pro Lys
          20           25           30

Val Leu Leu Pro Phe Thr Arg Ala Thr Arg Val Asn Phe Thr Leu Glu
      35           40           45

Ala Ser Glu Gly Cys Tyr Arg Trp Leu Ser Thr Arg Pro Glu Val Ala
      50           55           60

Ser Ile Glu Pro Leu Gly Leu Asp Glu Gln Gln Cys Ser Gln Lys Ala
      65           70           75           80

Val Val Gln Ala Arg Leu Thr Gln Pro Ala Arg Leu Thr Ser Ile Ile
          85           90           95

Phe Ala Glu Asp Ile Thr Thr Gly Gln Val Leu Arg Cys Asp Ala Ile
      100           105           110

Val Asp Leu Ile His Asp Ile Gln Ile Val Ser Thr Thr Arg Glu Leu
      115           120           125

Tyr Leu Glu Asp Ser Pro Leu Glu Leu Lys Ile Gln Ala Leu Asp Ser
      130           135           140

Glu Gly Asn Thr Phe Ser Thr Leu Ala Gly Leu Val Phe Glu Trp Thr
      145           150           155           160

Ile Val Lys Asp Ser Glu Ala Asp Arg Phe Ser Asp Ser His Asn Ala
          165           170           175

Leu Arg Ile Leu Thr Phe Leu Glu Ser Thr Tyr Ile Pro Pro Ser Tyr
          180           185           190

Ile Ser Glu Met Glu Lys Ala Ala Lys Gln Gly Asp Thr Ile Leu Val
          195           200           205

Ser Gly Met Lys Thr Gly Ser Ser Lys Leu Lys Ala Arg Ile Gln Glu
      210           215           220

Ala Val Tyr Lys Asn Val Arg Pro Ala Glu Val Arg Leu Leu Ile Leu
      225           230           235           240

Glu Asn Ile Leu Leu Asn Pro Ala Tyr Asp Val Tyr Leu Met Val Gly
          245           250           255

Thr Ser Ile His Tyr Lys Val Gln Lys Ile Arg Gln Gly Lys Ile Thr
          260           265           270

Glu Leu Ser Met Pro Ser Asp Gln Tyr Glu Leu Gln Leu Gln Asn Ser
      275           280           285

```

Ile Pro Gly Pro Glu Gly Asp Pro Thr Arg Pro Val Ala Val Leu Ala
 290 295 300
 Gln Asp Thr Ser Met Val Thr Ala Leu Gln Leu Gly Gln Ser Ser Leu
 305 310 315 320
 Val Leu Gly His Arg Ser Ile Arg Met Gln Gly Ala Ser Arg Leu Pro
 325 330 335
 Asn Ser Thr Ile Tyr Val Val Glu Pro Gly Tyr Leu Gly Phe Thr Val
 340 345 350
 His Pro Gly Asp Arg Trp Val Leu Glu Thr Gly Arg Leu Tyr Glu Ile
 355 360 365
 Thr Ile Glu Val Phe Asp Lys Phe Ser Asn Lys Val Tyr Val Ser Asp
 370 375 380
 Asn Ile Arg Ile Glu Thr Val Leu Pro Ala Glu Phe Phe Glu Val Leu
 385 390 395 400
 Ser Ser Ser Gln Asn Gly Ser Tyr His Arg Ile Arg Ala Leu Lys Arg
 405 410 415
 Gly Gln Thr Ala Ile Asp Ala Ala Leu Thr Ser Val Val Asp Gln Asp
 420 425 430
 Gly Gly Val His Ile Leu Gln Val Pro Val Trp Asn Gln Gln Glu Val
 435 440 445
 Glu Ile His Ile Pro Ile Thr Leu Tyr Pro Ser Ile Leu Thr Phe Pro
 450 455 460
 Trp Gln Pro Lys Thr Gly Ala Tyr Gln Tyr Thr Ile Arg Ala His Gly
 465 470 475 480
 Gly Ser Gly Asn Phe Ser Trp Ser Ser Ser Ser His Leu Val Ala Thr
 485 490 495
 Val Thr Val Lys Gly Val Met Thr Thr Gly Ser Asp Ile Gly Phe Ser
 500 505 510
 Val Ile Gln Ala His Asp Val Gln Asn Pro Leu His Phe Gly Glu Met
 515 520 525
 Lys Val Tyr Val Ile Glu Pro His Ser Met Glu Phe Ala Pro Cys Gln
 530 535 540
 Val Glu Ala Arg Val Gly Gln Ala Leu Glu Leu Pro Leu Arg Ile Ser
 545 550 555 560
 Gly Leu Met Pro Gly Gly Ala Ser Glu Val Val Thr Leu Ser Asp Cys
 565 570 575
 Ser His Phe Asp Leu Ala Val Glu Val Glu Asn Gln Gly Val Phe Gln
 580 585 590
 Pro Leu Pro Gly Arg Leu Pro Pro Gly Ser Glu His Cys Ser Gly Val

| 595 | 600 | 605 |
|--|-----|-----|
| Arg Val Lys Ala Glu Ala Gln Gly Ser Thr Thr Leu Leu Val Ser Tyr 610 615 620 | | |
| Arg His Gly His Val His Leu Ser Ala Lys Ile Thr Ile Ala Ala Tyr 625 630 635 640 | | |
| Leu Pro Leu Lys Ala Val Asp Pro Ser Ser Val Ala Leu Val Thr Leu 645 650 655 | | |
| Gly Ser Ser Lys Glu Met Leu Phe Glu Gly Gly Pro Arg Pro Trp Ile 660 665 670 | | |
| Leu Glu Pro Ser Lys Phe Phe Gln Asn Val Thr Ala Glu Asp Thr Asp 675 680 685 | | |
| Ser Ile Gly Leu Ala Leu Phe Ala Pro His Ser Ser Arg Asn Tyr Gln 690 695 700 | | |
| Gln His Trp Ile Leu Val Thr Cys Gln Ala Leu Gly Glu Gln Val Ile 705 710 715 720 | | |
| Ala Leu Ser Val Gly Asn Lys Pro Ser Leu Thr Asn Pro Phe Pro Ala 725 730 735 | | |
| Val Glu Pro Ala Val Val Lys Phe Val Cys Ala Pro Pro Ser Arg Leu 740 745 750 | | |
| Thr Leu Val Pro Val Tyr Thr Ser Pro Gln Leu Asp Met Ser Cys Pro 755 760 765 | | |
| Leu Leu Gln Gln Asn Lys Gln Val Val Pro Val Ser Ser His Arg Asn 770 775 780 | | |
| Pro Leu Leu Asp Leu Ala Ala Tyr Asp Gln Glu Gly Arg Arg Phe Asp 785 790 795 800 | | |
| Asn Phe Ser Ser Leu Ser Ile Gln Trp Glu Ser Thr Arg Pro Val Leu 805 810 815 | | |
| Ala Ser Ile Glu Pro Glu Leu Pro Met Gln Leu Val Ser Gln Asp Asp 820 825 830 | | |
| Glu Ser Gly Gln Lys Lys Leu His Gly Leu Gln Ala Ile Leu Val His 835 840 845 | | |
| Glu Ala Ser Gly Thr Thr Ala Ile Thr Ala Thr Ala Thr Gly Tyr Gln 850 855 860 | | |
| Glu Ser His Leu Ser Ser Ala Arg Thr Lys Gln Pro His Asp Pro Leu 865 870 875 880 | | |
| Val Pro Leu Ser Ala Ser Ile Glu Leu Ile Leu Val Glu Asp Val Arg 885 890 895 | | |
| Val Ser Pro Glu Glu Val Thr Ile Tyr Asn His Pro Gly Ile Gln Ala 900 905 910 | | |

Glu Leu Arg Ile Arg Glu Gly Ser Gly Tyr Phe Phe Leu Asn Thr Ser
 915 920 925
 Thr Ala Asp Val Val Lys Val Ala Tyr Gln Glu Ala Arg Gly Val Ala
 930 935 940
 Met Val His Pro Leu Leu Pro Gly Ser Ser Thr Ile Met Ile His Asp
 945 950 955 960
 Leu Cys Leu Val Phe Pro Ala Pro Ala Lys Ala Val Val Tyr Val Ser
 965 970 975
 Asp Ile Gln Glu Leu Tyr Ile Arg Val Val Asp Lys Val Glu Ile Gly
 980 985 990
 Lys Thr Val Lys Ala Tyr Val Arg Val Leu Asp Leu His Lys Lys Pro
 995 1000 1005
 Phe Leu Ala Lys Tyr Phe Pro Phe Met Asp Leu Lys Leu Arg Ala Ala
 1010 1015 1020
 Ser Pro Ile Ile Thr Leu Val Ala Leu Asp Glu Ala Leu Asp Asn Tyr
 1030 1035 1040
 Thr Ile Thr Phe Leu Ile Arg Gly Val Ala Ile Gly Gln Thr Ser Leu
 1045 1050 1055
 Thr Ala Ser Val Thr Asn Lys Ala Gly Gln Arg Ile Asn Ser Ala Pro
 1060 1065 1070
 Gln Gln Ile Glu Val Phe Pro Pro Phe Arg Leu Met Pro Arg Lys Val
 1075 1080 1085
 Thr Leu Leu Ile Gly Ala Thr Met Gln Val Thr Ser Glu Gly Gly Pro
 1090 1095 1100
 Gln Pro Gln Ser Asn Ile Leu Phe Ser Ile Ser Asn Glu Ser Val Ala
 1110 1115 1120
 Leu Val Ser Ala Ala Gly Leu Val Gln Gly Leu Ala Ile Gly Asn Gly
 1125 1130 1135
 Thr Val Ser Gly Leu Val Gln Ala Val Asp Ala Glu Thr Gly Lys Val
 1140 1145 1150
 Val Ile Ile Ser Gln Asp Leu Val Gln Val Glu Val Leu Leu Leu Arg
 1155 1160 1165
 Ala Val Arg Ile Arg Ala Pro Ile Met Arg Met Arg Thr Gly Thr Gln
 1170 1175 1180
 Met Pro Ile Tyr Val Thr Gly Ile Thr Asn His Gln Asn Pro Phe Ser
 1190 1195 1200
 Phe Gly Asn Ala Val Pro Gly Leu Thr Phe His Trp Ser Val Thr Lys
 1205 1210 1215

Arg Asp Val Leu Asp Leu Arg Gly Arg His His Glu Ala Ser Ile Arg
 1220 1225 1230
 Leu Pro Ser Gln Tyr Asn Phe Ala Met Asn Val Leu Gly Arg Val Lys
 1235 1240 1245
 Gly Arg Thr Gly Leu Arg Val Val Val Lys Ala Val Asp Pro Thr Ser
 1250 1255 1260
 Gly Gln Leu Tyr Gly Leu Ala Arg Glu Leu Ser Asp Glu Ile Gln Val
 1270 1275 1280
 Gln Val Phe Glu Lys Leu Gln Leu Leu Asn Pro Glu Ile Glu Ala Glu
 1285 1290 1295
 Gln Ile Leu Met Ser Pro Asn Ser Tyr Ile Lys Leu Gln Thr Asn Arg
 1300 1305 1310
 Asp Gly Ala Ala Ser Leu Ser Tyr Arg Val Leu Asp Gly Pro Glu Lys
 1315 1320 1325
 Val Pro Val Val His Val Asp Glu Lys Gly Phe Leu Ala Ser Gly Ser
 1330 1335 1340
 Met Ile Gly Thr Ser Thr Ile Gly Val Ile Ala Gln Glu Pro Phe Gly
 1350 1355 1360
 Ala Asn Gln Thr Ile Ile Val Ala Val Lys Val Ser Pro Val Ser Tyr
 1365 1370 1375
 Leu Arg Val Ser Met Ser Pro Val Leu His Thr Gln Asn Lys Glu Ala
 1380 1385 1390
 Leu Val Ala Val Pro Leu Gly Met Thr Val Thr Phe Thr Val His Phe
 1395 1400 1405
 His Asp Asn Ser Gly Asp Val Phe His Ala His Ser Ser Val Leu Asn
 1410 1415 1420
 Phe Ala Thr Asn Arg Asp Asp Phe Val Gln Ile Gly Lys Gly Pro Thr
 1430 1435 1440
 Asn Asn Thr Cys Val Val Arg Thr Val Ser Val Gly Leu Thr Leu Leu
 1445 1450 1455
 Arg Val Trp Asp Ala Glu His Pro Gly Leu Ser Asp Phe Met Pro Leu
 1460 1465 1470
 Pro Val Leu Gln Ala Ile Ser Pro Glu Leu Ser Gly Ala Met Val Val
 1475 1480 1485
 Gly Asp Val Leu Cys Leu Ala Thr Val Leu Thr Ser Leu Glu Gly Leu
 1490 1495 1500
 Ser Gly Thr Trp Ser Ser Ser Ala Asn Ser Ile Leu His Ile Asp Pro
 1510 1515 1520
 Lys Thr Gly Val Ala Val Ala Arg Ala Val Gly Ser Val Thr Val Tyr

| 1525 | 1530 | 1535 |
|---|------|------|
| Tyr Glu Val Ala Gly His Leu Arg Thr Tyr Lys Glu Val Val Val Ser 1540 | 1545 | 1550 |
| Val Pro Gln Arg Ile Met Ala Arg His Leu His Pro Ile Gln Thr Ser 1555 | 1560 | 1565 |
| Phe Gln Glu Ala Thr Ala Ser Lys Val Ile Val Ala Val Gly Asp Arg 1570 | 1575 | 1580 |
| Ser Ser Asn Leu Arg Gly Glu Cys Thr Pro Thr Gln Arg Glu Val Ile 1590 | 1595 | 1600 |
| Gln Ala Leu His Pro Glu Thr Leu Ile Ser Cys Gln Ser Gln Phe Lys 1605 | 1610 | 1615 |
| Pro Ala Val Phe Asp Phe Pro Ser Gln Asp Val Phe Thr Val Glu Pro 1620 | 1625 | 1630 |
| Gln Phe Asp Thr Ala Leu Gly Gln Tyr Phe Cys Ser Ile Thr Met His 1635 | 1640 | 1645 |
| Arg Leu Thr Asp Lys Gln Arg Lys His Leu Ser Met Lys Lys Thr Ala 1650 | 1655 | 1660 |
| Leu Val Val Ser Ala Ser Leu Ser Ser Ser His Phe Ser Thr Glu Gln 1670 | 1675 | 1680 |
| Val Gly Ala Glu Val Pro Phe Ser Pro Gly Leu Phe Ala Asp Gln Ala 1685 | 1690 | 1695 |
| Glu Ile Leu Leu Ser Asn His Tyr Thr Ser Ser Glu Ile Arg Val Phe 1700 | 1705 | 1710 |
| Gly Ala Pro Glu Val Leu Glu Asn Leu Glu Val Lys Ser Gly Ser Pro 1715 | 1720 | 1725 |
| Ala Val Leu Ala Phe Ala Lys Glu Lys Ser Phe Gly Trp Pro Ser Phe 1730 | 1735 | 1740 |
| Ile Thr Tyr Thr Val Gly Val Leu Asp Pro Ala Ala Gly Ser Gln Gly 1750 | 1755 | 1760 |
| Pro Leu Ser Thr Thr Leu Thr Phe Ser Ser Pro Val Thr Asn Gln Ala 1765 | 1770 | 1775 |
| Ile Ala Ile Pro Val Thr Val Ala Phe Val Val Asp Arg Arg Gly Pro 1780 | 1785 | 1790 |
| Gly Pro Tyr Gly Ala Ser Leu Phe Gln His Phe Leu Asp Ser Tyr Gln 1795 | 1800 | 1805 |
| Val Met Phe Phe Thr Leu Phe Ala Leu Leu Ala Gly Thr Ala Val Met 1810 | 1815 | 1820 |
| Ile Ile Ala Tyr His Thr Val Cys Thr Pro Arg Asp Leu Ala Val Pro 1830 | 1835 | 1840 |

Ala Ala Leu Thr Pro Arg Ala Ser Pro Gly His Ser Pro His Tyr Phe
 1845 1850 1855

Ala Ala Ser Ser Pro Thr Ser Pro Asn Ala Leu Pro Pro Ala Arg Lys
 1860 1865 1870

Ala Ser Pro Pro Ser Gly Leu Trp Ser Pro Ala Tyr Ala Ser His
 1875 1880 1885

<210> 108

<211> 29

<212> PRT

<213> Homo sapiens

<400> 108

Met Tyr Val Trp Val Ser Gly Ala Leu Val Leu Val Leu Ser Pro His
 1 5 10 15

Pro Ala Ser Arg Thr Leu Cys Leu Met Ala Gln Ala Val
 20 25

<210> 109

<211> 378

<212> PRT

<213> Homo sapiens

<400> 109

Ala Arg Glu Lys Pro Tyr Leu Val Glu Glu Ala Val Ser Tyr Asn Glu
 1 5 10 15

Leu Asp Tyr Val Ser Val Gly Leu Asp Gln Gln Thr Val Lys Leu Val
 20 25 30

Cys Thr Asn Arg Arg Lys Gln Phe Leu Leu Asp Thr Ala Asp Val Ala
 35 40 45

Leu Ala Glu Phe Phe Leu Ala Ser Leu Lys Ser Ala Met Ile Lys Gly
 50 55 60

Cys Arg Glu Pro Pro Tyr Pro Ser Ile Leu Thr Asp Ala Thr Met Glu
 65 70 75 80

Lys Leu Ala Leu Ala Lys Phe Val Ala Gln Glu Ser Lys Cys Glu Ala
 85 90 95

Ser Ala Val Thr Val Arg Phe Tyr Gly Leu Val His Trp Glu Asp Pro
 100 105 110

Thr Asp Glu Ser Leu Gly Pro Thr Pro Cys His Cys Ser Pro Pro Glu
 115 120 125

Gly Thr Ile Thr Lys Glu Gly Met Leu His Tyr Lys Ala Gly Thr Ser
 130 135 140

Tyr Leu Gly Lys Glu His Trp Lys Thr Cys Phe Val Val Leu Ser Asn

90

145 150 155 160
 Gly Ile Leu Tyr Gln Tyr Pro Asp Arg Thr Asp Val Ile Pro Leu Leu
 165 170 175
 Ser Val Asn Met Gly Gly Glu Gln Cys Gly Gly Cys Arg Arg Ala Asn
 180 185 190
 Thr Thr Asp Arg Pro His Ala Phe Gln Val Ile Leu Ser Asp Arg Pro
 195 200 205
 Cys Leu Glu Leu Ser Ala Glu Ser Glu Ala Glu Met Ala Glu Trp Met
 210 215 220
 Gln His Leu Cys Gln Ala Val Ser Lys Gly Val Ile Pro Gln Gly Val
 225 230 235 240
 Ala Pro Ser Pro Cys Ile Pro Cys Cys Leu Val Leu Thr Asp Asp Arg
 245 250 255
 Leu Phe Thr Cys His Glu Asp Cys Gln Thr Ser Phe Phe Arg Ser Leu
 260 265 270
 Gly Thr Ala Lys Leu Gly Asp Ile Ser Ala Val Ser Thr Glu Pro Gly
 275 280 285
 Lys Glu Tyr Cys Val Leu Glu Phe Ser Gln Asp Ser Gln Gln Leu Leu
 290 295 300
 Pro Pro Trp Val Ile Tyr Leu Ser Cys Thr Ser Glu Leu Asp Arg Leu
 305 310 315 320
 Leu Ser Ala Leu Asn Ser Gly Trp Lys Thr Ile Tyr Gln Val Asp Leu
 325 330 335
 Pro His Thr Ala Ile Gln Glu Ala Ser Asn Lys Lys Lys Phe Glu Asp
 340 345 350
 Ala Leu Ser Leu Ile His Ser Ala Trp Gln Arg Ser Asp Ser Leu Cys
 355 360 365
 Arg Gly Arg Ala Ser Arg Asp Pro Trp Cys
 370 375

<210> 110

<211> 66

<212> PRT

<213> Homo sapiens

<400> 110

Met Ser Trp Val Ile Val Val Ile Ile Trp Gly Tyr Leu Leu Glu Gly
 1 5 10 15
 His Gly Val Pro Phe Cys Lys Ser Tyr Gly Pro Ser Pro Trp Lys Leu
 20 25 30
 His Thr His His Ala Ala Tyr Asn Ser Gly Ser Ser Gln Val Tyr Arg

91

35 40 45
 Ile Leu Glu Thr Leu Met Ser Gly Ser Thr His Cys Ser Phe Ser Gly
 50 55 60
 Thr Phe
 65

<210> 111
 <211> 48
 <212> PRT
 <213> Homo sapiens

<400> 111
 Met Lys Pro Pro Pro Leu Phe Phe Phe Leu Lys Ile Val Leu Ala Ile
 1 5 10 15
 Trp Gly Pro Leu Trp Phe His Met Asn Phe Arg Phe Val Phe Ser Ile
 20 25 30
 Ser Met Lys Asn Ala Ile Gly Ile Leu Ile Gly Ile Ala Leu Asn Leu
 35 40 45

<210> 112
 <211> 778
 <212> PRT
 <213> Homo sapiens

<400> 112
 Met Leu Ser Ser Leu Gly Cys Leu Leu Leu Cys Gly Ser Ile Thr Leu
 1 5 10 15
 Ala Leu Gly Asn Ala Gln Lys Leu Pro Lys Gly Lys Arg Pro Asn Leu
 20 25 30
 Lys Val His Ile Asn Thr Thr Ser Asp Ser Ile Leu Leu Lys Phe Leu
 35 40 45
 Arg Pro Ser Pro Asn Val Lys Leu Glu Gly Leu Leu Leu Gly Tyr Gly
 50 55 60
 Ser Asn Val Ser Pro Asn Gln Tyr Phe Pro Leu Pro Ala Glu Gly Lys
 65 70 75 80
 Phe Thr Glu Ala Ile Val Asp Ala Glu Pro Lys Tyr Leu Ile Val Val
 85 90 95
 Arg Pro Ala Pro Pro Pro Ser Gln Lys Lys Ser Cys Ser Gly Lys Thr
 100 105 110
 Arg Ser Arg Lys Pro Leu Gln Leu Val Val Gly Thr Leu Thr Pro Ser
 115 120 125

Ser Val Phe Leu Ser Trp Gly Phe Leu Ile Asn Pro His His Asp Trp
 130 135 140
 Thr Leu Pro Ser His Cys Pro Asn Asp Arg Phe Tyr Thr Ile Arg Tyr
 145 150 155 160
 Arg Glu Lys Asp Lys Glu Lys Lys Trp Ile Phe Gln Ile Cys Pro Ala
 165 170 175
 Thr Glu Thr Ile Val Glu Asn Leu Lys Pro Asn Thr Val Tyr Glu Phe
 180 185 190
 Gly Val Lys Asp Asn Val Glu Gly Gly Ile Trp Ser Lys Ile Phe Asn
 195 200 205
 His Lys Thr Val Val Gly Ser Lys Lys Val Asn Gly Lys Ile Gln Ser
 210 215 220
 Thr Tyr Asp Gln Asp His Thr Val Pro Ala Tyr Val Pro Arg Lys Leu
 225 230 235 240
 Ile Pro Ile Thr Ile Ile Lys Gln Val Ile Gln Asn Val Thr His Lys
 245 250 255
 Asp Ser Ala Lys Ser Pro Glu Lys Ala Pro Leu Gly Gly Val Ile Leu
 260 265 270
 Val His Leu Ile Ile Pro Gly Leu Asn Glu Thr Thr Val Lys Leu Pro
 275 280 285
 Ala Ser Leu Met Phe Glu Ile Ser Asp Ala Leu Lys Thr Gln Leu Ala
 290 295 300
 Lys Asn Glu Thr Leu Ala Leu Pro Ala Glu Ser Lys Thr Pro Glu Val
 305 310 315 320
 Glu Lys Ile Ser Ala Arg Pro Thr Thr Val Thr Pro Glu Thr Val Pro
 325 330 335
 Arg Ser Thr Lys Pro Thr Thr Ser Ser Ala Leu Asp Val Ser Glu Thr
 340 345 350
 Thr Leu Val Leu Ser Lys Arg Thr Pro Glu Thr Leu Gln Thr Ile Leu
 355 360 365
 Ile Pro Gln Phe Glu Leu Pro Leu Ser Thr Leu Ala Pro Lys Ser Leu
 370 375 380
 Pro Glu Phe Pro Glu Ala Lys Thr Pro Phe Pro Phe Glu Lys Pro Arg
 385 390 395 400
 Gly Thr Leu Ala Ser Ser Glu Lys Pro Trp Ile Val Pro Thr Ala Lys
 405 410 415
 Ile Ser Glu Asp Ser Lys Val Leu Gln Pro Gln Thr Ala Thr Tyr Asp
 420 425 430
 Val Phe Ser Ser Pro Thr Thr Ser Asp Glu Pro Glu Ile Ser Asp Ser

| | | |
|---|-----|-----|
| 435 | 440 | 445 |
| Tyr Thr Ala Thr Ser Asp Arg Ile Leu Asp Ser Ile Pro Pro Lys Thr | | |
| 450 | 455 | 460 |
| Ser Arg Thr Leu Glu Gln Pro Arg Ala Thr Leu Ala Pro Ser Glu Thr | | |
| 465 | 470 | 475 |
| Pro Phe Val Pro Gln Lys Leu Glu Ile Phe Thr Ser Pro Glu Met Gln | | |
| | 485 | 490 |
| Pro Thr Thr Pro Ala Pro Gln Gln Thr Thr Ser Ile Pro Ser Thr Pro | | |
| | 500 | 510 |
| Lys Arg Arg Pro Arg Pro Lys Pro Pro Arg Thr Lys Pro Glu Arg Thr | | |
| | 515 | 525 |
| Thr Ser Ala Gly Thr Ile Thr Pro Lys Ile Ser Lys Ser Pro Glu Pro | | |
| | 530 | 540 |
| Thr Trp Thr Thr Pro Ala Pro Gly Lys Thr Gln Phe Ile Ser Leu Lys | | |
| | 545 | 555 |
| Pro Lys Ile Pro Leu Ser Pro Glu Val Thr His Thr Lys Pro Ala Pro | | |
| | 565 | 575 |
| Lys Gln Thr Pro Arg Ala Pro Pro Lys Pro Lys Thr Ser Pro Arg Pro | | |
| | 580 | 590 |
| Arg Ile Pro Gln Thr Gln Pro Val Pro Lys Val Pro Gln Arg Val Thr | | |
| | 595 | 605 |
| Ala Lys Pro Lys Thr Ser Pro Ser Pro Glu Val Ser Tyr Thr Thr Pro | | |
| | 610 | 620 |
| Ala Pro Lys Asp Val Leu Leu Pro His Lys Pro Tyr Pro Glu Val Ser | | |
| | 625 | 635 |
| Gln Ser Glu Pro Ala Pro Leu Glu Thr Arg Gly Ile Pro Phe Ile Pro | | |
| | 645 | 655 |
| Met Ile Ser Pro Ser Pro Ser Gln Glu Glu Leu Gln Thr Thr Leu Glu | | |
| | 660 | 670 |
| Glu Thr Asp Gln Ser Thr Gln Glu Pro Phe Thr Thr Lys Ile Pro Arg | | |
| | 675 | 685 |
| Thr Thr Glu Leu Ala Lys Thr Thr Gln Ala Pro His Arg Phe Tyr Thr | | |
| | 690 | 700 |
| Thr Val Arg Pro Arg Thr Ser Asp Lys Pro His Ile Arg Pro Val Leu | | |
| | 705 | 715 |
| Asn Arg Thr Thr Thr Arg Pro Thr Arg Pro Lys Pro Ser Gly Met Pro | | |
| | 725 | 735 |
| Ser Gly Asn Gly Val Gly Thr Gly Val Lys Gln Ala Pro Arg Pro Ser | | |
| | 740 | 750 |

Gly Ala Asp Arg Asn Val Ser Val Asp Ser Thr His Pro Thr Lys Lys
755 760 765

Pro Gly Thr Arg Arg Pro Pro Leu Pro Pro
770 775

```
<210> 113
<211> 55
<212> PRT
<213> Homo sapiens
```

<400> 113
Met Cys Val Gly Leu Phe Leu Ser Ser Val Phe Phe His Ile Cys Val
1 5 10 15

His Pro Phe Ala Asn Ala Thr Leu Ser Cys Leu Leu Glu Ile Gly Lys
20 25 30

Leu Cys Glu Ser Phe Asn Phe Val Leu Phe Gln Ile Val Leu Ala Ile
35 40 45

Leu Val Pro Leu Thr Phe Ile
50 55

```
<210> 114
<211> 33
<212> PRT
<213> Homo sapiens
```

```
<400> 114
Met Ile Ser Cys Leu Cys Asn Phe Ile Ala His Cys Val Ala Leu Val
  1             5             10            15
```

Met Arg Thr Cys Met Leu Val Val Ser Ser Asn Phe Ala Pro Ser Phe
20 25 30

Leu

```
<210> 115
<211> 157
<212> PRT
<213> Homo sapiens
```

```
<400> 115
Met Ser Pro Trp Leu Leu Leu Leu Val Val Gly Ser Trp Leu Leu
  1             5             10            15
```

Ala Arg Ile Leu Ala Trp Thr Tyr Ala Phe Tyr Asn Asn Cys Arg Arg
20 25 30

Leu Gln Cys Phe Pro Gln Pro Pro Lys Arg Asn Trp Phe Trp Gly His
35 40 45

95

Leu Gly Leu Ile Thr Pro Thr Glu Glu Gly Leu Lys Asp Ser Thr Gln
 50 55 60
 Met Ser Ala Thr Tyr Ser Gln Gly Phe Thr Val Trp Leu Gly Pro Ile
 65 70 75 80
 Ile Pro Phe Ile Val Leu Cys His Pro Asp Thr Ile Arg Ser Ile Thr
 85 90 95
 Asn Ala Ser Ala Ala Ile Ala Pro Lys Asp Asn Leu Phe Ile Arg Phe
 100 105 110
 Leu Lys Pro Trp Leu Gly Glu Tyr Leu Gln Val Lys Gly Val Gly Asp
 115 120 125
 Asn Leu Ala Gly Arg Val Gly Glu Val Leu Leu Leu Pro Ile Val Leu
 130 135 140
 Gly Cys Pro Thr Arg Arg Arg Asp Thr Ala Glu Trp Arg
 145 150 155

<210> 116
 <211> 102
 <212> PRT
 <213> Homo sapiens

<400> 116
 Met Asn Tyr Leu His Cys Asn Val Leu Leu Thr Leu Phe Cys Leu Leu
 1 5 10 15
 Phe Leu Leu His Ser Cys Ile Lys Ile Ile Lys His His Ser Gln Ala
 20 25 30
 Lys Arg Thr Arg Phe Pro Ser His Ile Ser His Lys Gly Glu Ala Asn
 35 40 45
 Thr His Gln Gly Gly Asn Tyr Thr Glu Leu Gly Trp Gly Leu Asp Ile
 50 55 60
 Tyr Phe Thr Ser Glu Leu Phe Ile Ser Ala Val Asn Leu Gly Glu Gly
 65 70 75 80
 Leu Gly Glu Val Leu Ser Gly Glu Gln Arg Gly Pro Gly Gly Lys Leu
 85 90 95
 Met Lys Thr Ser Asp Asp
 100

<210> 117
 <211> 72
 <212> PRT
 <213> Homo sapiens

<400> 117
 Met Ala Gly Asn Ile Gln Ala Val Glu Thr Gly Tyr Val Leu Ile Cys
 1 5 10 15

96

Leu Ile Val Pro Leu Leu Leu Cys Gly Leu Arg Glu Gly Gln Glu Val
 20 25 30

Pro Phe Asp Val Asn Lys Ala Lys Tyr Leu Pro Thr Phe Leu Lys Lys
 35 40 45

Lys Lys Lys Lys Lys Lys Lys Lys Lys Lys Lys Lys Lys Lys Lys Lys
 50 55 60

Lys Lys Lys Lys Lys Lys Lys Lys
 65 70

<210> 118

<211> 46

<212> PRT

<213> Homo sapiens

<400> 118

Met Arg Ile Pro Val Phe Pro Lys Gln Leu Met Phe Thr Gly Leu Val
 1 5 10 15

Phe Leu Leu Leu Leu Ser Lys Asp Glu Gly Ile His Asn Arg Leu Ser
 20 25 30

Leu Glu Asn Thr Asn Asp Gly Gln Leu Phe Gly Val Ile Lys
 35 40 45

<210> 119

<211> 19

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (7)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (11)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 119

Met Pro Phe Thr Leu Gly Xaa Thr Arg Arg Xaa Arg Gly Leu Ala Lys
 1 5 10 15

Lys Pro Lys

<210> 120

<211> 245

<212> PRT

<213> Homo sapiens

97

<400> 120

Met Ala Asp Val Ser Ala Lys Asp Ser Ser Gln Glu Thr Leu Val Asn
 1 5 10 15

Leu Ala Gly Leu Leu Val Ser Leu Leu Met Leu Pro Leu Val Ser Gly
 20 25 30

Cys Pro Gly Phe Ser Leu Gly Cys Phe Phe Phe Leu Thr Ala Leu His
 35 40 45

Ile Tyr Ala Asn Tyr Arg Ala Val Arg Ala Leu Val Met Glu Thr Leu
 50 55 60

Asn Glu Gly Arg Leu Arg Leu Val Leu Lys His Tyr Leu Gln Arg Gly
 65 70 75 80

Glu Val Leu Asp Pro Thr Ala Ala Asn Arg Met Glu Pro Leu Trp Thr
 85 90 95

Gly Phe Trp Pro Ala Pro Ser Leu Ser Leu Gly Val Pro Leu His Arg
 100 105 110

Leu Val Ser Ser Val Phe Glu Leu Gln Gln Leu Val Glu Gly His Gln
 115 120 125

Glu Ser Tyr Leu Leu Cys Trp Asp Gln Ser Gln Asn Gln Val Gln Val
 130 135 140

Val Leu Asn Gln Lys Ala Gly Pro Lys Thr Ile Leu Arg Ala Ala Thr
 145 150 155 160

His Gly Leu Met Leu Gly Ala Leu Gln Gly Asp Gly Pro Leu Pro Ala
 165 170 175

Glu Leu Glu Glu Leu Arg Asn Arg Val Arg Ala Gly Pro Lys Lys Glu
 180 185 190

Ser Trp Val Val Val Lys Glu Thr His Glu Val Leu Asp Met Leu Phe
 195 200 205

Pro Lys Phe Leu Lys Gly Leu Gln Asp Ala Gly Trp Lys Thr Glu Lys
 210 215 220

His Gln Leu Glu Val Asp Glu Trp Arg Ala Thr Trp Leu Leu Ser Pro
 225 230 235 240

Glu Lys Lys Val Leu
 245

<210> 121

<211> 75

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (62)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (63)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 121

Leu Pro Val Gln Asn Gly Cys Pro Glu Ser Ala Met Glu Met Asn Gly
1 5 10 15

Arg Ala Pro Cys Trp Glu Val Gly Leu Glu Glu Leu Ser Ser Arg Lys
20 25 30

Leu Thr Ala Gly Pro Gln Phe Pro Ser Glu Pro Gln Ala Pro Ala Pro
35 40 45

Ser Leu Phe Arg Gln Cys Leu Leu Trp Phe Cys Gly Met Xaa Xaa Gly
50 55 60

Gly Val Gly Ser Pro Pro Pro Leu Thr Gln Glu
65 70 75

<210> 122

<211> 96

<212> PRT

<213> Homo sapiens

<400> 122

Met Ala Arg Ser Leu Leu Ile Ile Leu Gly Ala Asp Phe Thr Phe Pro
1 5 10 15

Thr Ser Phe Asn Cys Phe Gln Lys Met Asn Leu Ala Lys Lys Ser Arg
20 25 30

Gly Ser Phe Thr His Leu Leu Thr His Ser Trp Cys Leu Ser Leu Phe
35 40 45

Leu Lys Glu Ala Asp Gln Gly Leu Arg Glu Asn Asn Phe Asp Phe Ser
50 55 60

His Val Cys Pro Ser Lys Pro Pro Leu Trp Thr Asp Ser Pro Ser Val
65 70 75 80

Pro Gly Arg Asn Trp Asp Asn Pro Arg Thr Phe Leu Val Pro Ser Arg
85 90 95

<210> 123

<211> 9

<212> PRT

<213> Homo sapiens

<400> 123

99

Leu Gly Ala Phe Ser Trp Ser Pro Lys
 1 5

<210> 124

<211> 44

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (3)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 124

Leu Lys Xaa Ile Thr Ile Cys Cys Leu Gln Lys Thr His Leu His Ser
 1 5 10 15

Lys Gly Thr Glu Arg Met Lys Val Lys Gly Trp Glu Arg Val Tyr Trp
 20 25 30

Gly Asn Ile Thr Glu Gly Asn Met Met Asn Leu Tyr
 35 40

<210> 125

<211> 77

<212> PRT

<213> Homo sapiens

<400> 125

Met Leu Leu Phe Phe Phe Phe Leu Leu Phe Phe Phe Phe Phe Trp
 1 5 10 15

Leu Val Leu Phe Gly Ile Phe Phe Phe Ser Phe Leu Lys Lys Met Phe
 20 25 30

Ser Gly Asn Met Asn Lys His Thr Ala Asn Tyr Ser Gly Ala Gly Lys
 35 40 45

Ala Gln Glu Leu Ala Thr Ser Gln Leu His Ser Trp Asp Gly Lys Pro
 50 55 60

Cys Cys Glu Leu Leu Arg Leu Phe Thr Tyr Phe Thr Tyr
 65 70 75

<210> 126

<211> 26

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (3)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

100

<221> SITE

<222> (5)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 126

Met Leu Xaa Phe Xaa Phe Phe Leu Leu Phe Phe Phe Phe Phe Trp Trp
 1 5 10 15

Cys Cys Leu Ala Phe Phe Ser Phe Pro Phe
 20 25

<210> 127

<211> 40

<212> PRT

<213> Homo sapiens

<400> 127

Met Arg Arg Gln Thr Phe Met Ser Ile Leu Val Phe Gln Cys Ser Pro
 1 5 10 15

Ile Ser Phe Gly Leu Cys Ile Asn Lys Glu Arg Thr Val Val Ser Ser
 20 25 30

Val Ile Thr Asp Asn Leu Cys Leu
 35 40

<210> 128

<211> 58

<212> PRT

<213> Homo sapiens

<400> 128

Met Lys Leu Thr Phe Ser Phe Pro Trp Phe Thr Leu Thr Ala Leu Gln
 1 5 10 15

Leu Trp Ser Ala Thr Glu Cys Gln Ala Val Val Asp Thr Met Ile Ala
 20 25 30

Val Trp Ser Glu Gly Lys Gly Thr Gly Val Ser Trp Glu Pro Trp Leu
 35 40 45

Leu Gly Lys Leu Gln Ser Ser Ser Phe Leu
 50 55

<210> 129

<211> 6

<212> PRT

<213> Homo sapiens

<400> 129

Thr Leu Thr Ala Lys Thr
 1 5

<210> 130

101

<211> 34

<212> PRT

<213> Homo sapiens

<400> 130

Leu Cys Val Ser His Pro Gly Ile Thr Cys Thr Pro Leu Trp Leu Cys
 1 5 10 15

Val Ile Ser Gln Asn Met Glu Leu Ile Leu Met Phe Arg Arg Pro Lys
 20 25 30

Leu Thr

<210> 131

<211> 90

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (22)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (25)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (26)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 131

Gln Asn Thr Ile Glu Cys Gly Ser Ser Thr Ala Gly Val Cys Cys Ser
 1 5 10 15

Gln Leu Trp Arg Leu Xaa Val Gln Xaa Xaa Gly Thr Gly Arg Leu His
 20 25 30

Val Trp Trp Gly Pro Ala Ser Trp Ser Ile Ala Ser Thr Phe Ser Leu
 35 40 45

His Pro Tyr Val Val Glu Glu Ala Gly Glu Leu Ser Gly Val Ser Phe
 50 55 60

Val Thr Pro Phe Leu Arg Leu Val His Ser His Asp Leu Ile Thr Ser
 65 70 75 80

Gln Arg Pro Cys Leu Leu Thr Pro Leu Pro
 85 90

<210> 132

<211> 90

<212> PRT

102

<213> Homo sapiens

<220>

<221> SITE

<222> (31)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (57)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 132

Met Pro Arg Ala Pro Trp Arg Ile Pro Leu Cys Ala Leu Pro Thr Leu
 1 5 10 15

Cys Leu Gly Ser Pro Leu Pro Ser Gln Pro Thr His Pro Ile Xaa Tyr
 20 25 30

Asp His Arg Ala Pro Thr Trp Lys Met Ala His Pro Gly Gly Pro Arg
 35 40 45

Ser Ser His Ser Pro Arg Thr Trp Xaa Thr Pro Ser Ser Gln Thr Lys
 50 55 60

Ala Ala Leu Pro Ala Gly Gly Ala Arg Asn Ser Pro Leu Gln Leu Cys
 65 70 75 80

Thr Arg Ser Arg Phe Cys Gly Thr Pro Met
 85 90

<210> 133

<211> 67

<212> PRT

<213> Homo sapiens

<400> 133

Met Ser Leu Cys Arg Ile Leu Gly Tyr Ser Phe Ser Ser Arg Leu Ser
 1 5 10 15

Ser Leu Ile Leu Pro Leu Ala Val Phe His Tyr Cys Leu Ser Cys Pro
 20 25 30

Leu His Phe Lys Leu Ser Phe Lys Tyr Leu Pro Phe Pro Ser Phe Pro
 35 40 45

Phe Ser Ser Leu Pro Cys Pro Ala Leu Pro Cys Pro Ala Leu Pro Ser
 50 55 60

Pro Pro Leu
 65

<210> 134

<211> 23

<212> PRT

<213> Homo sapiens

103

<400> 134

Met Phe Leu Lys Ala Gln Trp Leu Tyr Ser Leu Leu Leu Asn Cys Leu
 1 5 10 15

Leu Pro Glu Gly Thr Ser Ser
 20

<210> 135

<211> 7

<212> PRT

<213> Homo sapiens

<400> 135

Glu Ala Pro Trp Gln Phe Ser
 1 5

<210> 136

<211> 73

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (60)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 136

Ser Leu Leu Leu Ser Cys Cys Pro Leu Gly Asn Arg Ala Tyr Gly Ala
 1 5 10 15

Thr Gly Ala Glu Val Ala Ser Arg Ala Ser Leu Glu Gly Ser Glu His
 20 25 30

Ser Met Gln Arg Ser His Arg Glu Ala Gly Asn Gln Gly Pro Gly Arg
 35 40 45

Ala Ala Ser Cys Ala Ser Pro Ala Phe Val Met Xaa Phe Ser Phe Phe
 50 55 60

Thr His Cys Gln Ile Cys Phe Leu Pro
 65 70

<210> 137

<211> 64

<212> PRT

<213> Homo sapiens

<400> 137

Arg Thr Leu Arg Met Ser Pro Ser Ala Phe Cys Tyr Ser Leu Thr Leu
 1 5 10 15

Leu Ala Cys Trp Arg Ala Ala Trp Ile Pro Thr Cys Val Pro Arg Ala
 20 25 30

Ala Gly Glu Met Asp Ser Pro Gly Leu Ala Asp Gly His Trp Cys Ser
35 40 45

Gly Ala Ala Arg Arg Ser Pro His Tyr Val Ala Arg Ser Leu Val Leu
50 55 60

```

<400> 138
Leu Pro Trp Leu Pro Phe Phe Phe Ser Cys Leu Val Ser Thr Leu Pro
  1             5             10             15

Ser Met Ser Val Ser Ala Phe Ser Leu Val Val Arg Gly Arg Arg Ala
  20             25             30

Phe Thr Ser Val Arg
  35

```

```
<400> 139
Pro Leu Cys Leu Ala Leu Glu Leu Gly Trp Val Cys Leu Ser Ser Thr
  1             5             10            15
```

```
<220>
<221> SITE
<222> (43)
<223> Xaa equals any of the naturally occurring L-amino acids
```

```

<400> 140
Met Ala Gly Gln Phe Arg Ser Tyr Val Trp Asp Pro Leu Leu Ile Leu
  1             5             10             15
Ser Gln Ile Val Leu Met Gln Thr Val Tyr Tyr Gly Ser Leu Gly Leu
      20             25             30
Trp Leu Ala Leu Val Asp Gly Leu Val Arg Xaa Ala Pro Arg Trp Thr
    35             40             45

```

105

Arg Cys Ser Thr Pro Arg Ser Trp Ala Phe Pro Pro Leu Gln Ala Gly
 50 55 60

Ser Pro
 65

<210> 141
 <211> 124
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (28)
 <223> Xaa equals any of the naturally occurring L-amino acids

<400> 141
 Thr Gln Ile Pro Thr His Ile Ser Arg Tyr Thr Pro Leu His Ser Ser
 1 5 10 15

Leu Gly Asn Arg Ala Arg Leu Arg Leu Lys Lys Xaa Lys Ile Lys Tyr
 20 25 30

Ala Tyr Leu Cys Pro Pro Ser Leu Lys Gln Leu Leu Asn Tyr Ala Val
 35 40 45

Ile Asn Gly Leu Ser Ser Ala Asn Tyr Phe Cys Leu Tyr Thr Lys Val
 50 55 60

Pro Gln Ala Met Leu Leu Leu Ala Ser Gly Leu Ser Ser Ala Phe Pro
 65 70 75 80

Tyr Asp Ser Leu Gly Phe Thr Leu Ser Met Leu Leu Phe Phe Glu Arg
 85 90 95

Asn Lys Ser Arg Val Glu Val Leu Ala Lys Glu Pro Ser Ala Pro Ser
 100 105 110

Ser Tyr Trp Asp Ser Glu Asn Arg Gly Cys Gln Leu
 115 120

<210> 142
 <211> 214
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (199)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (206)
 <223> Xaa equals any of the naturally occurring L-amino acids

106

<220>

<221> SITE

<222> (214)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 142

Met Leu Gly Ala Arg Ala Trp Leu Gly Arg Val Leu Leu Leu Pro Arg
 1 5 10 15

Ala Gly Ala Gly Leu Ala Ala Ser Arg Arg Cys Pro Gly Val Trp Pro
 20 25 30

Arg Thr Trp Pro His Arg Ser Pro Ser Arg Gly Ser Ser Ser Arg Asp
 35 40 45

Lys Asp Arg Ser Ala Thr Val Ser Ser Ser Val Pro Met Pro Ala Gly
 50 55 60

Gly Lys Gly Ser His Pro Ser Ser Thr Pro Gln Arg Val Pro Asn Arg
 65 70 75 80

Leu Ile His Glu Lys Ser Pro Tyr Leu Leu Gln His Ala Tyr Asn Pro
 85 90 95

Val Asp Trp Tyr Pro Trp Gly Gln Glu Ala Phe Asp Lys Ala Arg Lys
 100 105 110

Glu Asn Lys Pro Ile Phe Leu Ser Val Gly Tyr Ser Thr Cys His Trp
 115 120 125

Cys His Met Met Glu Glu Glu Ser Phe Gln Asn Glu Glu Ile Gly Arg
 130 135 140

Leu Leu Ser Glu Asp Phe Val Ser Val Lys Val Asp Arg Glu Glu Arg
 145 150 155 160

Pro Asp Val Asp Lys Val Tyr Met Thr Phe Val Gln Ala Thr Ser Ser
 165 170 175

Gly Gly Gly Trp Pro Met Asn Val Trp Leu Thr Pro Asn Leu Gln Pro
 180 185 190

Phe Val Gly Gly Thr Ile Xaa Leu Leu Lys Asp Gly Leu Xaa Arg Val
 195 200 205

Gly Ser Ala Gln Cys Xaa
 210

<210> 143

<211> 43

<212> PRT

<213> Homo sapiens

<400> 143

Met Leu Gly Ala Arg Ala Trp Leu Gly Arg Val Leu Leu Leu Pro Arg
 1 5 10 15

107

Ala Gly Ala Gly Leu Ala Ala Ser Arg Arg Ser Ala Cys Ser Pro Thr
 20 25 30

Ser Arg Leu Asn Ser Leu Arg Ser Leu Ile Pro
 35 40

<210> 144

<211> 99

<212> PRT

<213> Homo sapiens

<400> 144

Ser Lys Ser Asn Pro Lys Pro Arg Cys Gln Lys Gly Thr Pro Trp Val
 1 5 10 15

Ile Arg Pro His Phe His Ser Asp Gly Val Ala Ser Ser Lys Thr Gly
 20 25 30

Leu Thr Val Phe Gln Met Ser Gly Leu Gln Ala Pro Ile Pro Ser Arg
 35 40 45

Cys Ser Ala Ala Ala Leu Ile Leu Arg Gly Gly Leu Pro Cys Thr Pro
 50 55 60

Leu Glu Ala Phe His Trp Gly Asn Cys Leu Pro Gly Ser Ala Leu Arg
 65 70 75 80

Ile Arg Ile Ala Lys Ala Gly Gln Ser Leu Pro Gln Gly Cys Ser Thr
 85 90 95

Gly Gln Ala

<210> 145

<211> 60

<212> PRT

<213> Homo sapiens

<400> 145

Met Arg Lys Trp Gly Leu Met Lys Leu Ile Ala Ser Met Met Gln Pro
 1 5 10 15

Val Leu Leu Glu Leu Leu Ser Val Trp Arg Lys Glu Gly Arg Asp Ser
 20 25 30

Arg Asn Ile His Asp Ser His Ser Met Tyr Val Leu Arg Lys Arg Leu
 35 40 45

Ser Gly Ser Trp Leu Gln Gln Val Cys Thr Leu Leu
 50 55 60

<210> 146

<211> 101

<212> PRT

108

<213> Homo sapiens

<400> 146

Val Asp Pro Arg Val Arg Thr Ser Ser Arg Ser Arg Ala Ala Ala Leu
 1 5 10 15
 Phe Glu Cys Phe Leu Met Val Phe Leu Leu Lys Cys Gln Val Asn Asn
 20 25 30
 Phe Asn Pro Ile Gln Gln Tyr Ser Leu Phe Pro Leu Lys Ser Ser Gly
 35 40 45
 Thr Cys Ser Ile Ser Leu Phe Cys Met Arg Gly Leu Tyr Phe Cys Leu
 50 55 60
 Gly Val Val Ile Cys Thr His Ala Ile Leu Leu Lys Pro Ser Cys Leu
 65 70 75 80
 Val Leu Phe Leu Glu Ser Phe Phe Phe Pro Val Leu Met Tyr Ala Gly
 85 90 95
 Phe Gly Asn Ser Ser
 100

<210> 147

<211> 98

<212> PRT

<213> Homo sapiens

<400> 147

Met His Cys Cys Gln Leu Pro Trp Arg Cys Ala Gln Ala Pro Gln Glu
 1 5 10 15
 Ala Phe Leu Leu Cys Leu Leu Phe Leu Ile Leu Val Leu Val Leu Leu
 20 25 30
 Gly Cys Ser Arg Gly Leu Pro Gly His Thr Pro Trp Arg Leu His Pro
 35 40 45
 Ala Ala Ala Ala Leu Leu Ala Pro Leu Leu His Asp Ala Leu Gly Ala
 50 55 60
 Cys Gly Phe Gln Gly Pro Glu Tyr Leu Leu Pro Cys Leu Leu Pro Leu
 65 70 75 80
 Pro Lys Pro Gly Gln Leu Gln Gly Pro Trp Gly Pro Leu Trp Ala Leu
 85 90 95
 Leu Pro

<210> 148

<211> 22

<212> PRT

<213> Homo sapiens

109

<400> 148

Leu Pro Arg Pro Cys Ala Pro Ser Pro Val Trp Arg Gln Val Gly Arg
1 5 10 15

Glu Glu Ala Ser Leu Leu
20

<210> 149

<211> 25

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (9)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 149

Cys Ala Val Arg Phe Arg Glu Gln Xaa Ala Pro Glu Arg Val Phe Leu
1 5 10 15

Pro Thr Arg Gly Arg Lys Ser Glu Pro
20 25

<210> 150

<211> 886

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (26)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (216)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (234)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (275)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (871)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 150

Met Ala Ala Arg Gly Arg Gly Leu Leu Leu Leu Thr Leu Ser Val Leu

110

| 1 | 5 | 10 | 15 |
|---|-----|-----|-----|
| Leu Ala Ala Gly Pro Ser Ala Ala Ala Xaa Lys Leu Asn Ile Pro Lys | 20 | 25 | 30 |
| Val Leu Leu Pro Phe Thr Arg Ala Thr Arg Val Asn Phe Thr Leu Glu | 35 | 40 | 45 |
| Ala Ser Glu Gly Cys Tyr Arg Trp Leu Ser Thr Arg Pro Glu Val Ala | 50 | 55 | 60 |
| Ser Ile Glu Pro Leu Gly Leu Asp Glu Gln Gln Cys Ser Gln Lys Ala | 65 | 70 | 80 |
| Val Val Gln Ala Arg Leu Thr Gln Pro Ala Arg Leu Thr Ser Ile Ile | 85 | 90 | 95 |
| Phe Ala Glu Asp Ile Thr Thr Gly Gln Val Leu Arg Cys Asp Ala Ile | 100 | 105 | 110 |
| Val Asp Leu Ile His Asp Ile Gln Ile Val Ser Thr Thr Arg Glu Leu | 115 | 120 | 125 |
| Tyr Leu Glu Asp Ser Pro Leu Glu Leu Lys Ile Gln Ala Leu Asp Ser | 130 | 135 | 140 |
| Glu Gly Asn Thr Phe Ser Thr Leu Ala Gly Leu Val Phe Glu Trp Thr | 145 | 150 | 155 |
| Ile Val Lys Asp Ser Glu Ala Asp Arg Phe Ser Asp Ser His Asn Ala | 165 | 170 | 175 |
| Leu Arg Ile Leu Thr Phe Leu Glu Ser Thr Tyr Ile Pro Pro Ser Tyr | 180 | 185 | 190 |
| Ile Ser Glu Met Glu Lys Ala Ala Lys Gln Gly Asp Thr Ile Leu Val | 195 | 200 | 205 |
| Ser Gly Met Lys Thr Gly Ser Xaa Lys Leu Lys Ala Arg Ile Gln Glu | 210 | 215 | 220 |
| Ala Val Tyr Lys Asn Val Arg Pro Ala Xaa Val Arg Leu Leu Ile Leu | 225 | 230 | 235 |
| Glu Asn Ile Leu Leu Asn Pro Ala Tyr Asp Val Tyr Leu Met Val Gly | 245 | 250 | 255 |
| Thr Ser Ile His Tyr Lys Val Gln Lys Ile Arg Gln Gly Lys Ile Thr | 260 | 265 | 270 |
| Glu Leu Xaa Met Pro Ser Asp Gln Tyr Glu Leu Gln Leu Gln Asn Ser | 275 | 280 | 285 |
| Ile Pro Gly Pro Glu Gly Asp Pro Thr Arg Pro Val Ala Val Leu Ala | 290 | 295 | 300 |
| Gln Asp Thr Ser Met Val Thr Ala Leu Gln Leu Gly Gln Ser Ser Leu | 305 | 310 | 315 |
| | | | 320 |

Val Leu Gly His Arg Ser Ile Arg Met Gln Gly Ala Ser Arg Leu Pro
 325 330 335
 Asn Ser Thr Ile Tyr Val Val Glu Pro Gly Tyr Leu Gly Phe Thr Val
 340 345 350
 His Pro Gly Asp Arg Trp Val Leu Glu Thr Gly Arg Leu Tyr Glu Ile
 355 360 365
 Thr Ile Glu Val Phe Asp Lys Phe Ser Asn Lys Val Tyr Val Ser Asp
 370 375 380
 Asn Ile Arg Ile Glu Thr Val Leu Pro Ala Glu Phe Phe Glu Val Leu
 385 390 395 400
 Ser Ser Ser Gln Asn Gly Ser Tyr His Arg Ile Arg Ala Leu Lys Arg
 405 410 415
 Gly Gln Thr Ala Ile Asp Ala Ala Leu Thr Ser Val Val Asp Gln Asp
 420 425 430
 Gly Gly Val His Ile Leu Gln Val Pro Val Trp Asn Gln Gln Glu Val
 435 440 445
 Glu Ile His Ile Pro Ile Thr Leu Tyr Pro Ser Ile Leu Thr Phe Pro
 450 455 460
 Trp Gln Pro Lys Thr Gly Ala Tyr Gln Tyr Thr Ile Arg Ala His Gly
 465 470 475 480
 Gly Ser Gly Asn Phe Ser Trp Ser Ser Ser Ser His Leu Val Ala Thr
 485 490 495
 Val Thr Val Lys Gly Val Met Thr Thr Gly Ser Asp Ile Gly Phe Ser
 500 505 510
 Val Ile Gln Ala His Asp Val Gln Asn Pro Leu His Phe Gly Glu Met
 515 520 525
 Lys Val Tyr Val Ile Glu Pro His Ser Met Glu Phe Ala Pro Cys Gln
 530 535 540
 Val Glu Ala Arg Val Gly Gln Ala Leu Glu Leu Pro Leu Arg Ile Ser
 545 550 555 560
 Gly Leu Met Pro Gly Gly Ala Ser Glu Val Val Thr Leu Ser Asp Cys
 565 570 575
 Ser His Phe Asp Leu Ala Val Glu Val Glu Asn Gln Gly Val Phe Gln
 580 585 590
 Pro Leu Pro Gly Arg Leu Pro Pro Gly Ser Glu His Cys Ser Gly Val
 595 600 605
 Arg Val Lys Ala Glu Ala Gln Gly Ser Thr Thr Leu Leu Val Ser Tyr
 610 615 620

112

Arg His Gly His Val His Leu Ser Ala Lys Ile Thr Ile Ala Ala Tyr
 625 630 635 640
 Leu Pro Leu Lys Ala Val Asp Pro Ser Ser Val Ala Leu Val Thr Leu
 645 650 655
 Gly Ser Ser Lys Glu Met Leu Phe Glu Gly Gly Pro Arg Pro Trp Ile
 660 665 670
 Leu Glu Pro Ser Lys Phe Phe Gln Asn Val Thr Ala Glu Asp Thr Asp
 675 680 685
 Ser Ile Gly Leu Ala Leu Phe Ala Pro His Ser Ser Arg Asn Tyr Gln
 690 695 700
 Gln His Trp Ile Leu Val Thr Cys Gln Ala Leu Gly Glu Gln Val Ile
 705 710 715 720
 Ala Leu Ser Val Gly Asn Lys Pro Ser Leu Thr Asn Pro Phe Pro Ala
 725 730 735
 Val Glu Pro Ala Val Val Lys Phe Val Cys Ala Pro Pro Ser Arg Leu
 740 745 750
 Thr Leu Val Pro Val Tyr Thr Ser Pro Gln Leu Asp Met Ser Cys Pro
 755 760 765
 Leu Leu Gln Gln Asn Lys Gln Val Val Pro Val Ser Ser His Arg Asn
 770 775 780
 Pro Leu Leu Asp Leu Ala Ala Tyr Asp Gln Glu Gly Arg Arg Phe Asp
 785 790 795 800
 Asn Phe Ser Ser Leu Ser Ile Gln Trp Glu Ser Thr Arg Pro Val Leu
 805 810 815
 Ala Ser Ile Glu Pro Glu Leu Pro Met Gln Leu Val Ser Gln Asp Asp
 820 825 830
 Glu Ser Gly Gln Lys Lys Leu His Gly Leu Gln Ala Ile Leu Val His
 835 840 845
 Glu Ala Ser Gly Thr Thr Ala Ser Leu Pro Leu Pro Leu Ala Thr Arg
 850 855 860
 Ser Pro Thr Ser Ala Leu Xaa Glu Gln Ser Ser Arg Met Thr Leu Trp
 865 870 875 880
 Cys Leu Cys Arg Pro Pro
 885

<210> 151

<211> 498

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (11)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (20)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (398)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 151

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Glu | Ala | Leu | Gly | Gly | Arg | Cys | Leu | Trp | Glu | Xaa | Pro | Val | Thr | Phe | Thr |
| 1 | | | | 5 | | | | | 10 | | | | | 15 | |

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Val | His | Phe | Xaa | Asp | Asn | Ser | Gly | Asp | Val | Phe | His | Ala | His | Ser | Ser |
| | | | 20 | | | | | 25 | | | | | | 30 | |

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Val | Leu | Asn | Phe | Ala | Thr | Asn | Arg | Asp | Asp | Phe | Val | Gln | Ile | Gly | Lys |
| | | 35 | | | | | 40 | | | | | 45 | | | |

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Gly | Pro | Thr | Asn | Asn | Thr | Cys | Val | Val | Arg | Thr | Val | Ser | Val | Gly | Leu |
| | 50 | | | | | 55 | | | | | 60 | | | | |

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Thr | Leu | Leu | Arg | Val | Trp | Asp | Ala | Glu | His | Pro | Gly | Leu | Ser | Asp | Phe |
| 65 | | | | | 70 | | | | | 75 | | | | | 80 |

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Met | Pro | Leu | Pro | Val | Leu | Gln | Ala | Ile | Ser | Pro | Glu | Leu | Ser | Gly | Ala |
| | | | | 85 | | | | | 90 | | | | | 95 | |

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Met | Val | Val | Gly | Asp | Val | Leu | Cys | Leu | Ala | Thr | Val | Leu | Thr | Ser | Leu |
| | | | 100 | | | | | 105 | | | | | 110 | | |

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Glu | Gly | Leu | Ser | Gly | Thr | Trp | Ser | Ser | Ser | Ala | Asn | Ser | Ile | Leu | His |
| | | 115 | | | | | 120 | | | | | 125 | | | |

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Ile | Asp | Pro | Lys | Thr | Gly | Val | Ala | Val | Ala | Arg | Ala | Val | Gly | Ser | Val |
| | 130 | | | | | 135 | | | | | 140 | | | | |

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Thr | Val | Tyr | Tyr | Glu | Val | Ala | Gly | His | Leu | Arg | Thr | Tyr | Lys | Glu | Val |
| 145 | | | | | 150 | | | | | 155 | | | | | 160 |

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Val | Val | Ser | Val | Pro | Gln | Arg | Ile | Met | Ala | Arg | His | Leu | His | Pro | Ile |
| | | | | 165 | | | | | 170 | | | | | 175 | |

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Gln | Thr | Ser | Phe | Gln | Glu | Ala | Thr | Ala | Ser | Lys | Val | Ile | Val | Ala | Val |
| | | | 180 | | | | | 185 | | | | | 190 | | |

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Gly | Asp | Arg | Ser | Ser | Asn | Leu | Arg | Gly | Glu | Cys | Thr | Pro | Thr | Gln | Arg |
| | | 195 | | | | 200 | | | | | | 205 | | | |

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Glu | Val | Ile | Gln | Ala | Leu | His | Pro | Glu | Thr | Leu | Ile | Ser | Cys | Gln | Ser |
| | 210 | | | | | 215 | | | | | 220 | | | | |

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Gln | Phe | Lys | Pro | Ala | Val | Phe | Asp | Phe | Pro | Ser | Gln | Asp | Val | Phe | Thr |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|

114

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 225 | | 230 | | 235 | | 240 | | | | | | | | | |
| Val | Glu | Pro | Gln | Phe | Asp | Thr | Ala | Leu | Gly | Gln | Tyr | Phe | Cys | Ser | Ile |
| | | | | 245 | | | | | 250 | | | | | 255 | |
| Thr | Met | His | Arg | Leu | Thr | Asp | Lys | Gln | Arg | Lys | His | Leu | Ser | Met | Lys |
| | | | 260 | | | | | 265 | | | | | 270 | | |
| Lys | Thr | Ala | Leu | Val | Val | Ser | Ala | Ser | Leu | Ser | Ser | Ser | His | Phe | Ser |
| | | 275 | | | | | 280 | | | | | 285 | | | |
| Thr | Glu | Gln | Val | Gly | Ala | Glu | Val | Pro | Phe | Ser | Pro | Gly | Leu | Phe | Ala |
| | 290 | | | | | 295 | | | | | 300 | | | | |
| Asp | Gln | Ala | Glu | Ile | Leu | Leu | Ser | Asn | His | Tyr | Thr | Ser | Ser | Glu | Ile |
| 305 | | | | | 310 | | | | | 315 | | | | 320 | |
| Arg | Val | Phe | Gly | Ala | Pro | Glu | Val | Leu | Glu | Asn | Leu | Glu | Val | Lys | Ser |
| | | | 325 | | | | | | 330 | | | | | 335 | |
| Gly | Ser | Pro | Ala | Val | Leu | Ala | Phe | Ala | Lys | Glu | Lys | Ser | Phe | Gly | Trp |
| | | | 340 | | | | | 345 | | | | | 350 | | |
| Pro | Ser | Phe | Ile | Thr | Tyr | Thr | Val | Gly | Val | Leu | Asp | Pro | Ala | Ala | Gly |
| | | 355 | | | | | 360 | | | | | 365 | | | |
| Ser | Gln | Gly | Pro | Leu | Ser | Thr | Thr | Leu | Thr | Phe | Ser | Ser | Pro | Val | Thr |
| | 370 | | | | | 375 | | | | | 380 | | | | |
| Asn | Gln | Ala | Ile | Ala | Ile | Pro | Val | Thr | Val | Ala | Phe | Val | Xaa | Asp | Arg |
| 385 | | | | | 390 | | | | | 395 | | | | 400 | |
| Arg | Gly | Pro | Gly | Pro | Tyr | Gly | Ala | Ser | Leu | Phe | Gln | His | Phe | Leu | Asp |
| | | | 405 | | | | | | 410 | | | | | 415 | |
| Ser | Tyr | Gln | Val | Met | Phe | Phe | Thr | Leu | Phe | Ala | Leu | Leu | Ala | Gly | Thr |
| | | 420 | | | | | | 425 | | | | | 430 | | |
| Ala | Val | Met | Ile | Ile | Ala | Tyr | His | Thr | Val | Cys | Thr | Pro | Arg | Asp | Leu |
| | | 435 | | | | | 440 | | | | | 445 | | | |
| Ala | Val | Pro | Ala | Ala | Leu | Thr | Pro | Arg | Ala | Ser | Pro | Gly | His | Ser | Pro |
| | 450 | | | | | 455 | | | | | 460 | | | | |
| His | Tyr | Phe | Ala | Ala | Ser | Ser | Pro | Thr | Ser | Pro | Asn | Ala | Leu | Pro | Pro |
| 465 | | | | | 470 | | | | | 475 | | | | 480 | |
| Ala | Arg | Lys | Ala | Ser | Pro | Pro | Ser | Gly | Leu | Trp | Ser | Pro | Ala | Tyr | Ala |
| | | | 485 | | | | | | 490 | | | | | 495 | |

Ser His

<210> 152

<211> 29

<212> PRT

<213> Homo sapiens

115

<220>

<221> SITE

<222> (26)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (29)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 152

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Met | Tyr | Val | Trp | Val | Ser | Gly | Ala | Leu | Val | Leu | Val | Leu | Ser | Pro | His |
| 1 | | | | 5 | | | | | 10 | | | | | 15 | |

| | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Pro | Ala | Ser | Arg | Thr | Leu | Cys | Leu | Met | Xaa | Gln | Ala | Xaa |
| | | | 20 | | | | | 25 | | | | |

<210> 153

<211> 80

<212> PRT

<213> Homo sapiens

<400> 153

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Pro | His | Cys | Ala | Ser | Arg | Ala | Val | Pro | Tyr | Pro | Pro | Gly | Pro | Ala | Ala |
| 1 | | | | 5 | | | | | 10 | | | | | 15 | |

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Ala | Ala | Phe | Pro | Arg | Gln | Gly | Leu | Gln | Leu | Ala | Thr | Thr | Cys | Gly | His |
| | | | 20 | | | | | 25 | | | | | 30 | | |

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Ser | Ser | Asp | Pro | Ala | Cys | Phe | Gly | Gln | Cys | Pro | Cys | His | Leu | Cys | Ala |
| | | 35 | | | | | 40 | | | | | 45 | | | |

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Asn | His | Pro | Gly | Tyr | Leu | Trp | Ser | Tyr | Arg | Val | His | Leu | Ser | Pro | Gln |
| | 50 | | | | | 55 | | | | | 60 | | | | |

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Pro | His | Leu | His | Pro | Pro | Gln | His | Leu | Leu | Pro | Pro | His | Cys | Thr | Leu |
| | 65 | | | | | 70 | | | | 75 | | | | | 80 |

<210> 154

<211> 11

<212> PRT

<213> Homo sapiens

<400> 154

| | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Met | Trp | Arg | Trp | Leu | Ser | Ser | Phe | Trp | Leu | Leu |
| 1 | | | | 5 | | | | | 10 | |

<210> 155

<211> 11

<212> PRT

<213> Homo sapiens

116

<400> 155

Met Trp Arg Trp Leu Ser Ser Phe Trp Leu Leu
 1 5 10

<210> 156

<211> 64

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (7)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (27)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 156

Ser Trp Val Ile Val Val Xaa Ile Trp Gly Tyr Leu Leu Glu Gly His
 1 5 10 15

Gly Val Pro Phe Cys Lys Ser Tyr Gly Pro Xaa Pro Trp Lys Leu His
 20 25 30

Thr His His Ala Ala Tyr Asn Ser Gly Ser Ser Gln Val Tyr Arg Ile
 35 40 45

Leu Gly Asn Ser Pro Cys Pro Val Leu Ile His Cys Ser Phe Ser Gly
 50 55 60

<210> 157

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (9)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (14)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 157

Trp Lys Gly Leu Leu Glu Gly Ser Xaa Glu Ala Thr Met Xaa
 1 5 10

117

<210> 158

<211> 107

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (66)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 158

Pro Leu Gly Arg Glu Pro Leu Ala Gly Phe Leu Ser Phe Leu Ser Phe
 1 5 10 15

Ser Leu Leu Trp Cys Leu Glu Ala Phe Pro Arg Leu Gln Phe Leu Thr
 20 25 30

Thr Leu Thr Asp Phe Ala Ile Val Leu Ser Pro Pro Leu Ser Phe Pro
 35 40 45

Lys Leu Thr Leu Trp Arg Leu Ile Lys Arg Lys Asn His Arg Pro Gly
 50 55 60

Ala Xaa Leu Thr Pro Arg Arg Arg Ala Asn His Leu Arg Cys Gly Val
 65 70 75 80

Arg Asp Gln Pro Asp Gln Asn Arg Glu Thr Pro Ser Leu Leu Asn Asn
 85 90 95

Thr Lys Leu Ala Gly Arg Gly Gly Ala Arg Leu
 100 105

<210> 159

<211> 48

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (15)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (29)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 159

Met Lys Pro Pro Pro Leu Phe Phe Phe Leu Lys Ile Val Leu Xaa Ile
 1 5 10 15

Trp Gly Pro Leu Trp Phe His Met Asn Phe Arg Phe Xaa Phe Ser Ile
 20 25 30

Ser Met Lys Asn Ala Ile Gly Ile Leu Ile Gly Ile Ala Leu Asn Leu
 35 40 45

118

<210> 160

<211> 73

<212> PRT

<213> Homo sapiens

<400> 160

Thr Leu Phe Val Ser Tyr Gln Leu Ser Asn Pro Gln Tyr Ser Ser Phe
 1 5 10 15

Ile Ser Gln Asn Arg Lys Leu Lys Gln Arg Glu Glu Lys Leu His Glu
 20 25 30

Arg Phe Tyr Thr Ala Val Arg Ser Leu Asn Trp Ile Leu Asn Leu Ala
 35 40 45

Phe Trp Leu Glu Ser Pro Ser Phe Tyr Gln Leu Cys Ile Ala Val Arg
 50 55 60

Val Asp Ser Pro Trp Lys Gly Lys Ser
 65 70

<210> 161

<211> 161

<212> PRT

<213> Homo sapiens

<400> 161

Met Leu Ser Ser Leu Gly Cys Leu Leu Leu Cys Gly Ser Ile Thr Leu
 1 5 10 15

Ala Leu Gly Asn Ala Gln Lys Leu Pro Lys Gly Lys Arg Pro Asn Leu
 20 25 30

Lys Val His Ile Asn Thr Thr Ser Asp Ser Ile Leu Leu Lys Phe Leu
 35 40 45

Arg Pro Ser Pro Asn Val Lys Leu Glu Gly Leu Leu Leu Gly Tyr Gly
 50 55 60

Ser Asn Val Ser Pro Asn Gln Tyr Phe Pro Leu Pro Ala Glu Gly Lys
 65 70 75 80

Phe Thr Glu Ala Ile Val Asp Ala Glu Pro Lys Tyr Leu Ile Val Val
 85 90 95

Arg Pro Ala Pro Pro Pro Ser Gln Lys Lys Ser Cys Ser Gly Lys Thr
 100 105 110

Arg Ser Arg Lys Pro Leu Gln Leu Val Val Gly Thr Leu Thr Pro Ser
 115 120 125

Ser Val Phe Leu Ser Trp Gly Phe Leu Ile Asn Pro His His Asp Trp
 130 135 140

119

Thr Leu Pro Ser His Cys Pro Asn Asp Arg Phe Tyr Thr Ile Arg Tyr
 145 150 155 160

Arg

<210> 162

<211> 181

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (80)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (151)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (162)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 162

Pro Lys Thr Ser Pro Ser Pro Glu Val Ser Tyr Thr Thr Pro Ala Pro
 1 5 10 15

Lys Asp Val Leu Leu Pro His Lys Pro Tyr Pro Glu Val Ser Gln Ser
 20 25 30

Glu Pro Ala Pro Leu Glu Thr Arg Gly Ile Pro Phe Ile Pro Met Ile
 35 40 45

Ser Pro Ser Pro Ser Gln Glu Glu Leu Gln Thr Thr Leu Glu Glu Thr
 50 55 60

Asp Gln Ser Thr Gln Glu Pro Phe Thr Thr Lys Ile Pro Arg Thr Xaa
 65 70 75 80

Glu Leu Ala Lys Thr Thr Gln Ala Pro His Arg Phe Tyr Thr Thr Val
 85 90 95

Arg Pro Arg Thr Ser Asp Lys Pro His Ile Arg Pro Val Leu Asn Arg
 100 105 110

Thr Thr Thr Arg Pro Thr Arg Pro Lys Pro Ser Gly Met Pro Ser Gly
 115 120 125

Asn Gly Val Gly Thr Gly Val Lys Gln Ala Pro Arg Pro Ser Gly Ala
 130 135 140

Asp Arg Asn Val Ser Val Xaa Ser Thr His Pro Thr Lys Lys Pro Gly
 145 150 155 160

120

Thr Xaa Arg Pro Pro Leu Pro Pro Ser Arg Arg Gly Arg Glu Phe Pro
 165 170 175

Gly Arg Arg Ala His
 180

<210> 163
 <211> 33
 <212> PRT
 <213> Homo sapiens

<400> 163
 Met Ile Ser Cys Leu Cys Asn Phe Ile Ala His Cys Val Ala Leu Val
 1 5 10 15

Met Arg Thr Cys Met Leu Val Val Ser Ser Asn Phe Ala Pro Ser Phe
 20 25 30

Leu

<210> 164
 <211> 85
 <212> PRT
 <213> Homo sapiens

<400> 164
 Ile Lys Thr Val Phe Leu Gly Gln Arg Tyr Thr Asp Pro Asn Phe Ile
 1 5 10 15

Ala Val Val Phe Ile His Leu Pro Ile Asp Ile Leu Lys Ala Pro Ala
 20 25 30

Arg Pro Gly Thr Val Ala His Ala Cys Asn Leu Ser Thr Leu Val Gly
 35 40 45

Arg Gly Gly Arg Ile Thr Arg Ser Arg Asp Gln Asp His Pro Gly Gln
 50 55 60

Arg Gly Glu Thr Leu Ser Leu Leu Lys Ile Gln Lys Leu Ala Gly His
 65 70 75 80

Gly Gly Ala Arg Leu
 85

<210> 165
 <211> 126
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (119)
 <223> Xaa equals any of the naturally occurring L-amino acids

121

<400> 165

Met Ser Pro Trp Leu Leu Leu Leu Val Val Gly Ser Trp Leu Leu
 1 5 10 15

Ala Arg Ile Leu Ala Trp Thr Tyr Ala Phe Tyr Asn Asn Cys Arg Arg
 20 25 30

Leu Gln Cys Phe Pro Gln Pro Pro Lys Arg Asn Trp Phe Trp Gly His
 35 40 45

Leu Gly Leu Ile Thr Pro Thr Glu Glu Gly Leu Lys Asp Ser Thr Gln
 50 55 60

Met Ser Ala Thr Tyr Ser Gln Gly Phe Thr Val Trp Leu Gly Pro Ile
 65 70 75 80

Ile Pro Phe Ile Val Leu Cys His Pro Asp Thr Ile Arg Ser Ile Thr
 85 90 95

Asn Ala Ser Ala Ala Ile Ala Pro Lys Asp Asn Leu Phe Ile Arg Phe
 100 105 110

Leu Lys Pro Trp Leu Gly Xaa Arg Asp Thr Ala Glu Trp Arg
 115 120 125

<210> 166

<211> 13

<212> PRT

<213> Homo sapiens

<400> 166

Leu Val Ile Gly Gly Trp Gly Gln Arg Arg Leu Tyr Arg
 1 5 10

<210> 167

<211> 33

<212> PRT

<213> Homo sapiens

<400> 167

Met Asn His Leu Ser Ile Ser Ile Ala Leu Phe Leu Leu Cys Cys Val
 1 5 10 15

His Leu Ser Leu Gly Leu Ser Val Phe Pro Phe Gln Glu Asp Arg Ser
 20 25 30

Val

<210> 168

<211> 132

<212> PRT

<213> Homo sapiens

122

<220>

<221> SITE

<222> (70)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 168

Ser Phe Arg Thr Ala Pro Arg Gly Pro His Val Lys Glu Ser His Ala
 1 5 10 15

Ser Gly Leu Leu Ser Asn Gln Ile Asn Leu Gln Ser Phe Asp Phe Lys
 20 25 30

Arg Met Leu Leu Cys Arg Leu Asn Ile Thr Gly Leu Cys Trp Gly Pro
 35 40 45

Lys Arg Thr Arg Cys Ala Leu Gly Gly Gln Thr Gly Leu Gln His His
 50 55 60

Pro Ser Asn Glu Lys Xaa Arg His Ser Gly Lys Glu Asp Leu Phe Leu
 65 70 75 80

Ser Ile Cys Leu Gly Trp Gly Thr Thr Val Asn Met Ala Cys Asn Asn
 85 90 95

Gln Arg Gly Arg Gly Tyr Gln Thr Gln Arg Asn Ser Ser Pro Val Tyr
 100 105 110

Gln Glu Glu Leu Leu Phe Phe Cys Thr Ser Leu Phe Ser Arg Leu Phe
 115 120 125

Ser Leu Lys Gly
 130

<210> 169

<211> 73

<212> PRT

<213> Homo sapiens

<400> 169

Met Ala Gly Asn Ile Gln Ala Val Glu Thr Gly Tyr Val Leu Ile Cys
 1 5 10 15

Leu Ile Val Pro Leu Leu Leu Cys Gly Leu Arg Glu Gly Gln Glu Val
 20 25 30

Pro Phe Asp Val Asn Lys Ala Lys Tyr Leu Pro Thr Phe Leu Lys Lys
 35 40 45

Lys Lys Lys Lys Lys Lys Lys Lys Lys Lys Lys Lys Lys Lys Lys Lys
 50 55 60

Lys Lys Lys Lys Lys Lys Lys Lys Ile
 65 70

<210> 170

<211> 53

123

<212> PRT

<213> Homo sapiens

<400> 170

Met Arg Ile Pro Val Phe Pro Lys Gln Leu Met Phe Thr Gly Leu Val
 1 5 10 15

Phe Leu Leu Leu Leu Ser Lys Asp Glu Gly Ile His Asn Arg Leu Ser
 20 25 30

Leu Glu Asn Thr Asn Asp Gly Gln Leu Phe Gly Val Ile Asn Glu Leu
 35 40 45

Ala Thr Thr Leu Met
 50

<210> 171

<211> 344

<212> PRT

<213> Homo sapiens

<400> 171

Asp Gly Pro Arg Ser Lys Pro Gln Pro Lys Asp Asn Gly Asp Val Cys
 1 5 10 15

Gln Asp Cys Ile Gln Met Val Thr Asp Ile Gln Thr Ala Val Arg Thr
 20 25 30

Asn Ser Thr Phe Val Gln Ala Leu Val Glu His Val Lys Glu Glu Cys
 35 40 45

Asp Arg Leu Gly Pro Gly Met Ala Asp Ile Cys Lys Asn Tyr Ile Ser
 50 55 60

Gln Tyr Ser Glu Ile Ala Ile Gln Met Met Met His Met Gln Asp Gln
 65 70 75 80

Gln Pro Lys Glu Ile Cys Ala Leu Val Gly Phe Cys Asp Glu Val Lys
 85 90 95

Glu Met Pro Met Gln Thr Leu Val Pro Ala Lys Val Ala Ser Lys Asn
 100 105 110

Val Ile Pro Ala Leu Glu Leu Val Glu Pro Ile Lys Lys His Glu Val
 115 120 125

Pro Ala Lys Ser Asp Val Tyr Cys Glu Val Cys Glu Phe Leu Val Lys
 130 135 140

Glu Val Thr Lys Leu Ile Asp Asn Asn Lys Thr Glu Lys Glu Ile Leu
 145 150 155 160

Asp Ala Phe Asp Lys Met Cys Ser Lys Leu Pro Lys Ser Leu Ser Glu
 165 170 175

Glu Cys Gln Glu Val Val Asp Thr Tyr Gly Ser Ser Ile Leu Ser Ile
 180 185 190

124

Leu Leu Glu Glu Val Ser Pro Glu Leu Val Cys Ser Met Leu His Leu
 195 200 205

Cys Ser Gly Thr Arg Leu Pro Ala Leu Thr Val His Val Thr Gln Pro
 210 215 220

Lys Asp Gly Gly Phe Cys Glu Val Cys Lys Lys Leu Val Gly Tyr Leu
 225 230 235 240

Asp Arg Asn Leu Glu Lys Asn Ser Thr Lys Gln Glu Ile Leu Ala Ala
 245 250 255

Leu Glu Lys Gly Cys Ser Phe Leu Pro Asp Pro Tyr Gln Lys Gln Cys
 260 265 270

Asp Gln Phe Val Ala Glu Tyr Glu Pro Val Leu Ile Glu Ile Leu Val
 275 280 285

Glu Val Met Asp Pro Ser Phe Val Cys Leu Lys Ile Gly Ala Cys Pro
 290 295 300

Ser Ala His Lys Pro Leu Leu Gly Thr Glu Lys Cys Ile Trp Gly Pro
 305 310 315 320

Ser Tyr Trp Cys Gln Asn Thr Glu Thr Ala Ala Gln Cys Asn Ala Val
 325 330 335

Glu His Cys Lys Arg His Val Trp
 340

<210> 172

<211> 345

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (268)

<223> Xaa equals stop translation

<220>

<221> SITE

<222> (317)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (323)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 172

Asn Gly Pro Leu Thr Phe His Pro Arg Gln Ala Pro Glu Gly Ala Leu
 1 5 10 15

Cys Gln Asp Cys Val Arg Gln Val Ser Arg Leu Gln Glu Ala Val Arg
 20 25 30

125

Ser Asn Leu Thr Leu Ala Asp Leu Asn Ile Gln Glu Gln Cys Glu Ser
 35 40 45
 Leu Gly Pro Gly Leu Ala Val Leu Cys Lys Asn Tyr Leu Phe Gln Phe
 50 55 60
 Phe Val Pro Ala Asp Gln Ala Leu Arg Leu Leu Pro Pro Gln Glu Leu
 65 70 75 80
 Cys Arg Lys Gly Gly Phe Cys Glu Glu Leu Gly Ala Pro Ala Arg Leu
 85 90 95
 Thr Gln Val Val Ala Met Asp Gly Val Pro Ser Leu Glu Leu Gly Leu
 100 105 110
 Pro Arg Lys Gln Ser Glu Met Gln Met Lys Ala Gly Val Thr Cys Glu
 115 120 125
 Val Cys Met Asn Val Val Gln Lys Leu Asp His Trp Leu Met Ser Asn
 130 135 140
 Ser Ser Glu Leu Met Ile Thr His Ala Leu Glu Arg Val Cys Ser Val
 145 150 155 160
 Met Pro Ala Ser Ile Thr Lys Glu Cys Ile Ile Leu Val Asp Thr Tyr
 165 170 175
 Ser Pro Ser Leu Val Gln Leu Val Ala Lys Ile Thr Pro Glu Lys Val
 180 185 190
 Cys Lys Phe Ile Arg Leu Cys Gly Asn Arg Arg Arg Ala Arg Ala Val
 195 200 205
 His Asp Ala Tyr Ala Ile Val Pro Ser Pro Glu Trp Asp Ala Glu Asn
 210 215 220
 Gln Gly Ser Phe Cys Asn Gly Cys Lys Arg Leu Leu Thr Val Ser Ser
 225 230 235 240
 His Asn Leu Glu Ser Lys Ser Thr Lys Arg Asp Ile Leu Val Ala Phe
 245 250 255
 Lys Gly Gly Cys Ser Ile Leu Pro Leu Pro Ile Xaa Ser Lys Cys Lys
 260 265 270
 His Phe Arg His Pro Ile Arg Ser Pro Cys Ser Leu Arg Val Ser Lys
 275 280 285
 Asp Met Met Gly Pro Arg Gly Cys Val Gln Glu Gly Gly Gly Pro Gly
 290 295 300
 Thr Ala Pro Arg Pro His Leu Leu Gly Thr Asp Gln Xaa Gly Leu Ala
 305 310 315 320
 Gln Ala Xaa Gly Ala Arg Thr Lys Lys Ala Ala Ser Phe Gly Asn Ala
 325 330 335

126

Leu Gln His Trp Gln Lys His Gly Trp
 340 345

<210> 173

<211> 822

<212> PRT

<213> Homo sapiens

<400> 173

Met Ala Ala Ala Val Val Val Ala Glu Gly Asp Ser Asp Ser Arg Pro
 1 5 10 15

Gly Gln Glu Leu Leu Val Ala Trp Asn Thr Val Ser Thr Gly Leu Val
 20 25 30

Pro Pro Ala Ala Leu Gly Leu Val Ser Ser Arg Thr Ser Gly Ala Val
 35 40 45

Pro Pro Lys Glu Glu Glu Leu Arg Ala Ala Val Glu Val Leu Arg Gly
 50 55 60

His Gly Leu His Ser Val Leu Glu Glu Trp Phe Val Glu Val Leu Gln
 65 70 75 80

Asn Asp Leu Gln Ala Asn Ile Ser Pro Glu Phe Trp Asn Ala Ile Ser
 85 90 95

Gln Cys Glu Asn Ser Ala Asp Glu Pro Gln Cys Leu Leu Leu Leu Leu
 100 105 110

Asp Ala Phe Gly Leu Leu Glu Ser Arg Leu Asp Pro Tyr Leu Arg Ser
 115 120 125

Leu Glu Leu Leu Glu Lys Trp Thr Arg Leu Gly Leu Leu Met Gly Thr
 130 135 140

Gly Ala Gln Gly Leu Arg Glu Glu Val His Thr Met Leu Arg Gly Val
 145 150 155 160

Leu Phe Phe Ser Thr Pro Arg Thr Phe Gln Glu Met Ile Gln Arg Leu
 165 170 175

Tyr Gly Cys Phe Leu Arg Val Tyr Met Gln Ser Lys Arg Lys Gly Glu
 180 185 190

Gly Gly Thr Asp Pro Glu Leu Glu Gly Glu Leu Asp Ser Arg Tyr Ala
 195 200 205

Arg Arg Arg Tyr Tyr Arg Leu Leu Gln Ser Pro Leu Cys Ala Gly Cys
 210 215 220

Ser Ser Asp Lys Gln Gln Cys Trp Cys Arg Gln Ala Leu Glu Gln Phe
 225 230 235 240

His Gln Leu Ser Gln Val Leu His Arg Leu Ser Leu Leu Glu Arg Val
 245 250 255

127

Ser Ala Glu Ala Val Thr Thr Thr Leu His Gln Val Thr Arg Glu Arg
 260 265 270
 Met Glu Asp Arg Cys Arg Gly Glu Tyr Glu Arg Ser Phe Leu Arg Glu
 275 280 285
 Phe His Lys Trp Ile Glu Arg Val Val Gly Trp Leu Gly Lys Val Phe
 290 295 300
 Leu Gln Asp Gly Pro Ala Arg Pro Ala Ser Pro Glu Ala Gly Asn Thr
 305 310 315 320
 Leu Arg Arg Trp Arg Cys His Val Gln Arg Phe Phe Tyr Arg Ile Tyr
 325 330 335
 Ala Ser Leu Arg Ile Glu Glu Leu Phe Ser Ile Val Arg Asp Phe Pro
 340 345 350
 Asp Ser Arg Pro Ala Ile Glu Asp Leu Lys Tyr Cys Leu Glu Arg Thr
 355 360 365
 Asp Gln Arg Gln Gln Leu Leu Val Ser Leu Lys Ala Ala Leu Glu Thr
 370 375 380
 Arg Leu Leu His Pro Gly Val Asn Thr Cys Asp Ile Ile Thr Leu Tyr
 385 390 395 400
 Ile Ser Ala Ile Lys Ala Leu Arg Val Leu Asp Pro Ser Met Val Ile
 405 410 415
 Leu Glu Val Ala Cys Glu Pro Ile Arg Arg Tyr Leu Arg Thr Arg Glu
 420 425 430
 Asp Thr Val Arg Gln Ile Val Ala Gly Leu Thr Gly Asp Ser Asp Gly
 435 440 445
 Thr Gly Asp Leu Ala Val Glu Leu Ser Lys Thr Asp Pro Ala Ser Leu
 450 455 460
 Glu Thr Gly Gln Asp Ser Glu Asp Asp Ser Gly Glu Pro Glu Asp Trp
 465 470 475 480
 Val Pro Asp Pro Val Asp Ala Asp Pro Gly Lys Ser Ser Ser Lys Arg
 485 490 495
 Arg Ser Ser Asp Ile Ile Ser Leu Leu Val Ser Ile Tyr Gly Ser Lys
 500 505 510
 Asp Leu Phe Ile Asn Glu Tyr Arg Ser Leu Leu Ala Asp Arg Leu Leu
 515 520 525
 His Gln Phe Ser Phe Ser Pro Glu Arg Glu Ile Arg Asn Val Glu Leu
 530 535 540
 Leu Lys Leu Arg Phe Gly Glu Ala Pro Met His Phe Cys Glu Val Met
 545 550 555 560
 Leu Lys Asp Met Ala Asp Ser Arg Arg Ile Asn Ala Asn Ile Arg Glu

128

| 565 | | | | | | | | | | 570 | | | | | 575 | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|--|--|--|--|
| Glu | Asp | Glu | Lys | Arg | Pro | Ala | Glu | Glu | Gln | Pro | Pro | Phe | Gly | Val | Tyr | | | | |
| | | | 580 | | | | | 585 | | | | | 590 | | | | | | |
| Ala | Val | Ile | Leu | Ser | Ser | Glu | Phe | Trp | Pro | Pro | Phe | Lys | Asp | Glu | Lys | | | | |
| | | 595 | | | | | 600 | | | | | 605 | | | | | | | |
| Leu | Glu | Val | Pro | Glu | Asp | Ile | Arg | Ala | Ala | Leu | Glu | Ala | Tyr | Cys | Lys | | | | |
| | 610 | | | | | 615 | | | | | 620 | | | | | | | | |
| Lys | Tyr | Glu | Gln | Leu | Lys | Ala | Met | Arg | Thr | Leu | Ser | Trp | Lys | His | Thr | | | | |
| 625 | | | | | 630 | | | | | 635 | | | | | 640 | | | | |
| Leu | Gly | Leu | Val | Thr | Met | Asp | Val | Glu | Leu | Ala | Asp | Arg | Thr | Leu | Ser | | | | |
| | | | | 645 | | | | | 650 | | | | | | 655 | | | | |
| Val | Ala | Val | Thr | Pro | Val | Gln | Ala | Val | Ile | Leu | Leu | Tyr | Phe | Gln | Asp | | | | |
| | | | 660 | | | | | 665 | | | | | 670 | | | | | | |
| Gln | Ala | Ser | Trp | Thr | Leu | Glu | Glu | Leu | Ser | Lys | Ala | Val | Lys | Met | Pro | | | | |
| | | 675 | | | | | 680 | | | | | 685 | | | | | | | |
| Val | Ala | Leu | Leu | Arg | Arg | Arg | Met | Ser | Val | Trp | Leu | Gln | Gln | Gly | Val | | | | |
| | | 690 | | | | 695 | | | | | 700 | | | | | | | | |
| Leu | Arg | Glu | Glu | Pro | Pro | Gly | Thr | Phe | Ser | Val | Ile | Glu | Glu | Glu | Arg | | | | |
| 705 | | | | | 710 | | | | | 715 | | | | | 720 | | | | |
| Pro | Gln | Asp | Arg | Asp | Asn | Met | Val | Leu | Ile | Asp | Ser | Asp | Asp | Glu | Ser | | | | |
| | | | | 725 | | | | | 730 | | | | | 735 | | | | | |
| Asp | Ser | Gly | Met | Ala | Ser | Gln | Ala | Asp | Gln | Lys | Glu | Glu | Glu | Leu | Leu | | | | |
| | | | 740 | | | | | 745 | | | | | 750 | | | | | | |
| Leu | Phe | Trp | Thr | Tyr | Ile | Gln | Ala | Met | Leu | Thr | Asn | Leu | Glu | Ser | Leu | | | | |
| | | 755 | | | | 760 | | | | | | 765 | | | | | | | |
| Ser | Leu | Asp | Arg | Ile | Tyr | Asn | Met | Leu | Arg | Met | Phe | Val | Val | Thr | Gly | | | | |
| | | 770 | | | | 775 | | | | | 780 | | | | | | | | |
| Pro | Ala | Leu | Ala | Glu | Ile | Asp | Leu | Gln | Glu | Leu | Gln | Gly | Tyr | Leu | Gln | | | | |
| 785 | | | | | 790 | | | | | 795 | | | | | 800 | | | | |
| Lys | Lys | Val | Arg | Asp | Gln | Gln | Leu | Val | Tyr | Ser | Ala | Gly | Val | Tyr | Arg | | | | |
| | | | 805 | | | | | | 810 | | | | | 815 | | | | | |
| Leu | Pro | Lys | Asn | Cys | Ser | | | | | | | | | | | | | | |
| | | | 820 | | | | | | | | | | | | | | | | |

<210> 174

<211> 822

<212> PRT

<213> Homo sapiens

<400> 174

Met Ala Ala Ala Val Val Val Ala Glu Gly Asp Ser Asp Ser Arg Pro

129

| | | | |
|---|-----|-----|-----|
| 1 | 5 | 10 | 15 |
| Gly Gln Glu Leu Leu Val Ala Trp Asn Thr Val Ser Thr Gly Leu Val | 20 | 25 | 30 |
| Pro Pro Ala Ala Leu Gly Leu Val Ser Ser Arg Thr Ser Gly Ala Val | 35 | 40 | 45 |
| Pro Pro Lys Glu Glu Glu Leu Arg Ala Ala Val Glu Val Leu Arg Gly | 50 | 55 | 60 |
| His Gly Leu His Ser Val Leu Glu Glu Trp Phe Val Glu Val Leu Gln | 65 | 70 | 75 |
| Asn Asp Leu Gln Ala Asn Ile Ser Pro Glu Phe Trp Asn Ala Ile Ser | 85 | 90 | 95 |
| Gln Cys Glu Asn Ser Ala Asp Glu Pro Gln Cys Leu Leu Leu Leu Leu | 100 | 105 | 110 |
| Asp Ala Phe Gly Leu Leu Glu Ser Arg Leu Asp Pro Tyr Leu Arg Ser | 115 | 120 | 125 |
| Leu Glu Leu Leu Glu Lys Trp Thr Arg Leu Gly Leu Leu Met Gly Thr | 130 | 135 | 140 |
| Gly Ala Gln Gly Leu Arg Glu Glu Val His Thr Met Leu Arg Gly Val | 145 | 150 | 155 |
| Leu Phe Phe Ser Thr Pro Arg Thr Phe Gln Glu Met Ile Gln Arg Leu | 165 | 170 | 175 |
| Tyr Gly Cys Phe Leu Arg Val Tyr Met Gln Ser Lys Arg Lys Gly Glu | 180 | 185 | 190 |
| Gly Gly Thr Asp Pro Glu Leu Glu Gly Glu Leu Asp Ser Arg Tyr Ala | 195 | 200 | 205 |
| Arg Arg Arg Tyr Tyr Arg Leu Leu Gln Ser Pro Leu Cys Ala Gly Cys | 210 | 215 | 220 |
| Ser Ser Asp Lys Gln Gln Cys Trp Cys Arg Gln Ala Leu Glu Gln Phe | 225 | 230 | 235 |
| His Gln Leu Ser Gln Val Leu His Arg Leu Ser Leu Leu Glu Arg Val | 245 | 250 | 255 |
| Ser Ala Glu Ala Val Thr Thr Thr Leu His Gln Val Thr Arg Glu Arg | 260 | 265 | 270 |
| Met Glu Asp Arg Cys Arg Gly Glu Tyr Glu Arg Ser Phe Leu Arg Glu | 275 | 280 | 285 |
| Phe His Lys Trp Ile Glu Arg Val Val Gly Trp Leu Gly Lys Val Phe | 290 | 295 | 300 |
| Leu Gln Asp Gly Pro Ala Arg Pro Ala Ser Pro Glu Ala Gly Asn Thr | 305 | 310 | 315 |
| | | | 320 |

130

Leu Arg Arg Trp Arg Cys His Val Gln Arg Phe Phe Tyr Arg Ile Tyr
 325 330 335
 Ala Ser Leu Arg Ile Glu Glu Leu Phe Ser Ile Val Arg Asp Phe Pro
 340 345 350
 Asp Ser Arg Pro Ala Ile Glu Asp Leu Lys Tyr Cys Leu Glu Arg Thr
 355 360 365
 Asp Gln Arg Gln Gln Leu Leu Val Ser Leu Lys Ala Ala Leu Glu Thr
 370 375 380
 Arg Leu Leu His Pro Gly Val Asn Thr Cys Asp Ile Ile Thr Leu Tyr
 385 390 395 400
 Ile Ser Ala Ile Lys Ala Leu Arg Val Leu Asp Pro Ser Met Val Ile
 405 410 415
 Leu Glu Val Ala Cys Glu Pro Ile Arg Arg Tyr Leu Arg Thr Arg Glu
 420 425 430
 Asp Thr Val Arg Gln Ile Val Ala Gly Leu Thr Gly Asp Ser Asp Gly
 435 440 445
 Thr Gly Asp Leu Ala Val Glu Leu Ser Lys Thr Asp Pro Ala Ser Leu
 450 455 460
 Glu Thr Gly Gln Asp Ser Glu Asp Asp Ser Gly Glu Pro Glu Asp Trp
 465 470 475 480
 Val Pro Asp Pro Val Asp Ala Asp Pro Gly Lys Ser Ser Ser Lys Arg
 485 490 495
 Arg Ser Ser Asp Ile Ile Ser Leu Leu Val Ser Ile Tyr Gly Ser Lys
 500 505 510
 Asp Leu Phe Ile Asn Glu Tyr Arg Ser Leu Leu Ala Asp Arg Leu Leu
 515 520 525
 His Gln Phe Ser Phe Ser Pro Glu Arg Glu Ile Arg Asn Val Glu Leu
 530 535 540
 Leu Lys Leu Arg Phe Gly Glu Ala Pro Met His Phe Cys Glu Val Met
 545 550 555 560
 Leu Lys Asp Met Ala Asp Ser Arg Arg Ile Asn Ala Asn Ile Arg Glu
 565 570 575
 Glu Asp Glu Lys Arg Pro Ala Glu Glu Gln Pro Pro Phe Gly Val Tyr
 580 585 590
 Ala Val Ile Leu Ser Ser Glu Phe Trp Pro Pro Phe Lys Asp Glu Lys
 595 600 605
 Leu Glu Val Pro Glu Asp Ile Arg Ala Ala Leu Glu Ala Tyr Cys Lys
 610 615 620

131

Lys Tyr Glu Gln Leu Lys Ala Met Arg Thr Leu Ser Trp Lys His Thr
625 630 635 640

Leu Gly Leu Val Thr Met Asp Val Glu Leu Ala Asp Arg Thr Leu Ser
645 650 655

Val Ala Val Thr Pro Val Gln Ala Val Ile Leu Leu Tyr Phe Gln Asp
660 665 670

Gln Ala Ser Trp Thr Leu Glu Glu Leu Ser Lys Ala Val Lys Met Pro
675 680 685

Val Ala Leu Leu Arg Arg Arg Met Ser Val Trp Leu Gln Gln Gly Val
690 695 700

Leu Arg Glu Glu Pro Pro Gly Thr Phe Ser Val Ile Glu Glu Glu Arg
705 710 715 720

Pro Gln Asp Arg Asp Asn Met Val Leu Ile Asp Ser Asp Asp Glu Ser
725 730 735

Asp Ser Gly Met Ala Ser Gln Ala Asp Gln Lys Glu Glu Glu Leu Leu
740 745 750

Leu Phe Trp Thr Tyr Ile Gln Ala Met Leu Thr Asn Leu Glu Ser Leu
755 760 765

Ser Leu Asp Arg Ile Tyr Asn Met Leu Arg Met Phe Val Val Thr Gly
770 775 780

Pro Ala Leu Ala Glu Ile Asp Leu Gln Glu Leu Gln Gly Tyr Leu Gln
785 790 795 800

Lys Lys Val Arg Asp Gln Gln Leu Val Tyr Ser Ala Gly Val Tyr Arg
805 810 815

Leu Pro Lys Asn Cys Ser
820

<210> 175

<211> 145

<212> PRT

<213> Homo sapiens

<400> 175

Met Ala Phe Trp Thr Gln Leu Met Leu Leu Leu Trp Lys Asn Phe Met
1 5 10 15

Tyr Arg Arg Arg Gln Pro Val Gln Leu Val Glu Leu Leu Trp Pro
20 25 30

Leu Phe Leu Phe Phe Ile Leu Val Ala Val Arg His Ser His Pro Pro
35 40 45

Leu Glu His His Glu Cys His Phe Pro Asn Lys Pro Leu Pro Ser Ala
50 55 60

132

Gly Thr Val Pro Trp Leu Gln Gly Leu Ile Cys Asn Val Asn Asn Thr
 65 70 75 80
 Cys Phe Pro Gln Leu Thr Pro Gly Glu Glu Pro Gly Arg Leu Ser Asn
 85 90 95
 Phe Asn Asp Ser Leu Val Ser Arg Leu Leu Ala Asp Ala Arg Thr Val
 100 105 110
 Leu Gly Gly Ala Ser Ala His Arg Thr Leu Ala Gly Leu Gly Lys Leu
 115 120 125
 Ile Ala Thr Leu Arg Ala Ala Arg Ser Thr Ala Gln Pro Gln Pro Thr
 130 135 140
 Lys
 145

<210> 176
 <211> 50
 <212> PRT
 <213> Homo sapiens

<400> 176
 Arg Pro Leu Met Gln Asn Gly Gly Pro Glu Thr Phe Thr Lys Leu Met
 1 5 10 15
 Gly Ile Leu Ser Asp Leu Leu Cys Gly Tyr Pro Glu Gly Gly Gly Ser
 20 25 30
 Arg Val Leu Ser Phe Asn Trp Tyr Glu Asp Asn Asn Tyr Lys Ala Phe
 35 40 45
 Leu Gly
 50

<210> 177
 <211> 48
 <212> PRT
 <213> Homo sapiens

<400> 177
 Arg Ala Leu Leu Gln Arg Pro Arg Gly Thr Ser Gly Pro Leu Glu Leu
 1 5 10 15
 Leu Ser Glu Ala Leu Cys Ser Val Arg Gly Pro Ser Ser Thr Val Gly
 20 25 30
 Pro Ser Leu Asn Trp Tyr Glu Ala Ser Asp Leu Met Glu Leu Val Gly
 35 40 45

<210> 178

133

<211> 145

<212> PRT

<213> Homo sapiens

<400> 178

Met Gly Phe Val Arg Gln Ile Gln Leu Leu Leu Trp Lys Asn Trp Thr
 1 5 10 15

Leu Arg Lys Arg Gln Lys Ile Arg Phe Val Val Glu Leu Val Trp Pro
 20 25 30

Leu Ser Leu Phe Leu Val Leu Ile Trp Leu Arg Asn Ala Asn Pro Leu
 35 40 45

Tyr Ser His His Glu Cys His Phe Pro Asn Lys Ala Met Pro Ser Ala
 50 55 60

Gly Met Leu Pro Trp Leu Gln Gly Ile Phe Cys Asn Val Asn Asn Pro
 65 70 75 80

Cys Phe Gln Ser Pro Thr Pro Gly Glu Ser Pro Gly Ile Val Ser Asn
 85 90 95

Tyr Asn Asn Ser Ile Leu Ala Arg Val Tyr Arg Asp Phe Gln Glu Leu
 100 105 110

Leu Met Asn Ala Pro Glu Ser Gln His Leu Gly Arg Ile Trp Thr Glu
 115 120 125

Leu His Ile Leu Ser Gln Phe Met Asp Thr Leu Arg Thr His Pro Glu
 130 135 140

Arg
 145

<210> 179

<211> 833

<212> PRT

<213> Homo sapiens

<400> 179

Ala Val Ala Ala Lys Leu Asn Ile Pro Lys Val Leu Leu Pro Phe Thr
 1 5 10 15

Arg Ala Thr Arg Val Asn Phe Thr Leu Glu Ala Ser Glu Gly Cys Tyr
 20 25 30

Arg Trp Ser Ser Thr Arg Pro Glu Val Ala Ser Ile Glu Pro Leu Gly
 35 40 45

Ser Ser Glu Gln Gln Cys Ser Gln Lys Ala Val Val Gln Ala Arg Leu
 50 55 60

Thr Gln Pro Ala Arg Leu Thr Ser Ile Ile Phe Ala Glu Asp Ile Thr
 65 70 75 80

Thr Gly Gln Val Leu Arg Cys Asp Ala Ile Val Asp Leu Ile His Gly

134

| 85 | | | | | 90 | | | | | 95 | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Ile | Gln | Ile | Val | Ser | Thr | Thr | Arg | Glu | Leu | Tyr | Leu | Glu | Asp | Ser | Pro |
| | | | 100 | | | | | 105 | | | | | 110 | | |
| Leu | Glu | Leu | Lys | Ile | Gln | Ala | Leu | Asp | Ser | Glu | Gly | Asn | Thr | Phe | Ser |
| | | | 115 | | | | | 120 | | | | | 125 | | |
| Thr | Leu | Ala | Gly | Leu | Val | Phe | Asp | Trp | Thr | Ile | Val | Lys | Asp | Thr | Glu |
| | | | 130 | | | | | 135 | | | | | 140 | | |
| Ala | Asn | Gly | Phe | Ser | Asp | Ser | His | Asn | Ala | Leu | Arg | Ile | Leu | Thr | Phe |
| | | | | | | | | | | | | | | | 160 |
| Leu | Glu | Ser | Thr | Tyr | Ile | Pro | Pro | Ser | Tyr | Ile | Ser | Glu | Met | Glu | Lys |
| | | | | | | | | | | | | | | | 175 |
| Ala | Ala | Lys | Gln | Gly | Asp | Thr | Ile | Leu | Val | Ser | Gly | Met | Lys | Thr | Gly |
| | | | | | | | | | | | | | | | 190 |
| Ser | Ser | Lys | Leu | Lys | Ala | Arg | Ile | Gln | Glu | Ala | Val | Tyr | Lys | Asn | Val |
| | | | | | | | | | | | | | | | 205 |
| Arg | Pro | Ala | Glu | Val | Arg | Leu | Leu | Ile | Leu | Glu | Asn | Ile | Leu | Leu | Asn |
| | | | | | | | | | | | | | | | 220 |
| Pro | Ala | Tyr | Asp | Val | Tyr | Leu | Leu | Val | Gly | Thr | Ser | Ile | His | Tyr | Lys |
| | | | | | | | | | | | | | | | 240 |
| Val | Gln | Lys | Ile | Arg | Gln | Gly | Lys | Ile | Thr | Glu | Leu | Ser | Met | Pro | Ser |
| | | | | | | | | | | | | | | | 255 |
| Asp | Gln | Tyr | Glu | Leu | Gln | Leu | Gln | Asn | Ser | Ile | Pro | Asp | Pro | Gln | Gly |
| | | | | | | | | | | | | | | | 270 |
| Asp | Pro | Ala | Arg | Pro | Val | Ala | Val | Leu | Thr | Gln | Asp | Thr | Ser | Arg | Val |
| | | | | | | | | | | | | | | | 285 |
| Thr | Ala | Met | Gln | Met | Gly | Gln | Ser | Asn | Leu | Val | Leu | Gly | His | Arg | Ser |
| | | | | | | | | | | | | | | | 300 |
| Ile | Arg | Met | Gln | Gly | Ala | Ser | Arg | Leu | Pro | Asn | Ser | Thr | Ile | Tyr | Val |
| | | | | | | | | | | | | | | | 320 |
| Val | Glu | Ala | Gly | Tyr | Leu | Gly | Phe | Thr | Val | His | Pro | Gly | Asp | Arg | Trp |
| | | | | | | | | | | | | | | | 335 |
| Val | Leu | Glu | Thr | Gly | His | Leu | Tyr | Ala | Val | Thr | Ile | Glu | Val | Phe | Asp |
| | | | | | | | | | | | | | | | 350 |
| Arg | Ser | Ser | Asn | Lys | Val | Tyr | Pro | Ser | Asp | Asn | Ile | Arg | Ile | Glu | Ala |
| | | | | | | | | | | | | | | | 365 |
| Val | Phe | Pro | Ala | Glu | Phe | Phe | Glu | Val | Leu | Ser | Ser | Ser | Gln | Asn | Gly |
| | | | | | | | | | | | | | | | 380 |
| Ser | Tyr | His | His | Val | Arg | Ala | Ile | Gln | Ser | Gly | Gln | Thr | Thr | Ile | Ser |
| | | | | | | | | | | | | | | | 400 |

135

Ala Ser Leu Thr Ser Val Val Asp Gln Asp Gly Gly Val His Val Leu
 405 410 415
 Gln Val Pro Val Trp Asn Gln Gln Glu Val Asp Ile His Ile Pro Ile
 420 425 430
 Thr Leu Tyr Pro Ser Ile Leu Thr Phe Pro Trp Gln Pro Lys Thr Gly
 435 440 445
 Ala Tyr Gln Tyr Thr Ile Lys Ala His Gly Gly Ser Gly Asn Phe Thr
 450 455 460
 Trp Ser Ser Ser Ser Tyr Met Val Ala Thr Val Thr Val Lys Gly Val
 465 470 475 480
 Met Thr Thr Gly Gly Asp Thr Gly Leu Ser Val Ile Arg Ala His Asp
 485 490 495
 Val Gln Asn Pro Leu His Phe Gly Glu Met Lys Val Tyr Val Ile Glu
 500 505 510
 Pro Ser Ser Met Glu Phe Ala Pro Cys Gln Val Glu Ala Arg Val Gly
 515 520 525
 His Thr Leu Glu Leu Pro Leu Thr Ile Ser Gly Leu Met Pro Gly Gly
 530 535 540
 Ser Ser Glu Val Val Thr Leu Ser Asp Cys Ser His Phe Asp Leu Val
 545 550 555 560
 Val Glu Val Glu Asn Gln Gly Val Phe Gln Pro Leu Pro Gly Arg Leu
 565 570 575
 Pro Pro Gly Pro Glu His Cys Ser Gly Val Lys Val Arg Ala Asp Ala
 580 585 590
 Gln Gly Ser Thr Thr Leu Leu Val Ser Tyr Thr His Gly His Val His
 595 600 605
 Leu Gly Ala Lys Ile Thr Leu Ala Ala Tyr Leu Pro Leu Lys Ala Val
 610 615 620
 Asp Pro Ser Ser Val Ala Val Val Thr Leu Gly Ser Ser Lys Glu Met
 625 630 635 640
 Leu Phe Glu Gly Gly Pro Arg Pro Trp Val Leu Glu Pro Ser Lys Phe
 645 650 655
 Phe Arg Asn Val Thr Ser Glu Asp Thr Gly Ser Ile Ser Leu Ser Leu
 660 665 670
 Leu Gly Pro Pro Ala Ser Arg Asn Tyr Gln Gln His Arg Val Leu Val
 675 680 685
 Thr Cys Gln Ala Leu Gly Glu Gln Val Ile Ala Leu Ser Val Gly Asn
 690 695 700

136

Arg Pro Ser Leu Ser Asn Pro Phe Pro Ala Val Glu Pro Thr Val Val
705 710 715 720

Lys Ser Val Cys Ala Pro Pro Ser Arg Leu Thr Leu Met Pro Val Tyr
725 730 735

Ala Leu Pro Gln Leu Asp Leu Ser Cys Pro Leu Leu Gln Gln Asn Lys
740 745 750

Gln Val Val Pro Val Ser Ser His Arg Asn Pro Leu Leu Asp Leu Gly
755 760 765

Ala Tyr Asp Gln Gln Gly Arg Arg Phe Asp Asn Phe Ser Ser Leu Ser
770 775 780

Ile Gln Trp Glu Ser Phe Arg Pro Leu Leu Ala Ser Ile Glu Val Asp
785 790 795 800

Gln Pro Met Gln Leu Val Ser Gln Asp Asp Gly Asn Gly Gln Lys Lys
805 810 815

Leu His Gly Leu Gln Thr Val Ser Val His Glu Ala Ser Gly Thr Thr
820 825 830

Ala

<210> 180

<211> 833

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (4)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (194)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (212)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (253)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 180

Ala Ala Ala Xaa Lys Leu Asn Ile Pro Lys Val Leu Leu Pro Phe Thr
1 5 10 15

Arg Ala Thr Arg Val Asn Phe Thr Leu Glu Ala Ser Glu Gly Cys Tyr

137

| 20 | | | | | 25 | | | | | 30 | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Arg | Trp | Leu | Ser | Thr | Arg | Pro | Glu | Val | Ala | Ser | Ile | Glu | Pro | Leu | Gly |
| | | 35 | | | | | 40 | | | | | 45 | | | |
| Leu | Asp | Glu | Gln | Gln | Cys | Ser | Gln | Lys | Ala | Val | Val | Gln | Ala | Arg | Leu |
| | 50 | | | | | 55 | | | | | 60 | | | | |
| Thr | Gln | Pro | Ala | Arg | Leu | Thr | Ser | Ile | Ile | Phe | Ala | Glu | Asp | Ile | Thr |
| | 65 | | | | | 70 | | | | | 75 | | | | 80 |
| Thr | Gly | Gln | Val | Leu | Arg | Cys | Asp | Ala | Ile | Val | Asp | Leu | Ile | His | Asp |
| | | | | 85 | | | | | 90 | | | | | 95 | |
| Ile | Gln | Ile | Val | Ser | Thr | Thr | Arg | Glu | Leu | Tyr | Leu | Glu | Asp | Ser | Pro |
| | | | 100 | | | | | 105 | | | | | 110 | | |
| Leu | Glu | Leu | Lys | Ile | Gln | Ala | Leu | Asp | Ser | Glu | Gly | Asn | Thr | Phe | Ser |
| | 115 | | | | | | 120 | | | | | 125 | | | |
| Thr | Leu | Ala | Gly | Leu | Val | Phe | Glu | Trp | Thr | Ile | Val | Lys | Asp | Ser | Glu |
| | 130 | | | | | 135 | | | | | 140 | | | | |
| Ala | Asp | Arg | Phe | Ser | Asp | Ser | His | Asn | Ala | Leu | Arg | Ile | Leu | Thr | Phe |
| | 145 | | | | | 150 | | | | | 155 | | | | 160 |
| Leu | Glu | Ser | Thr | Tyr | Ile | Pro | Pro | Ser | Tyr | Ile | Ser | Glu | Met | Glu | Lys |
| | | | 165 | | | | | 170 | | | | | 175 | | |
| Ala | Ala | Lys | Gln | Gly | Asp | Thr | Ile | Leu | Val | Ser | Gly | Met | Lys | Thr | Gly |
| | | | 180 | | | | | 185 | | | | | 190 | | |
| Ser | Xaa | Lys | Leu | Lys | Ala | Arg | Ile | Gln | Glu | Ala | Val | Tyr | Lys | Asn | Val |
| | | 195 | | | | | 200 | | | | | 205 | | | |
| Arg | Pro | Ala | Xaa | Val | Arg | Leu | Leu | Ile | Leu | Glu | Asn | Ile | Leu | Leu | Asn |
| | 210 | | | | | 215 | | | | | 220 | | | | |
| Pro | Ala | Tyr | Asp | Val | Tyr | Leu | Met | Val | Gly | Thr | Ser | Ile | His | Tyr | Lys |
| | 225 | | | | | 230 | | | | | 235 | | | | 240 |
| Val | Gln | Lys | Ile | Arg | Gln | Gly | Lys | Ile | Thr | Glu | Leu | Xaa | Met | Pro | Ser |
| | | | 245 | | | | | 250 | | | | | 255 | | |
| Asp | Gln | Tyr | Glu | Leu | Gln | Leu | Gln | Asn | Ser | Ile | Pro | Gly | Pro | Glu | Gly |
| | | 260 | | | | | 265 | | | | | 270 | | | |
| Asp | Pro | Thr | Arg | Pro | Val | Ala | Val | Leu | Ala | Gln | Asp | Thr | Ser | Met | Val |
| | | 275 | | | | | 280 | | | | | 285 | | | |
| Thr | Ala | Leu | Gln | Leu | Gly | Gln | Ser | Ser | Leu | Val | Leu | Gly | His | Arg | Ser |
| | 290 | | | | | 295 | | | | | 300 | | | | |
| Ile | Arg | Met | Gln | Gly | Ala | Ser | Arg | Leu | Pro | Asn | Ser | Thr | Ile | Tyr | Val |
| | 305 | | | | | 310 | | | | | 315 | | | | 320 |
| Val | Glu | Pro | Gly | Tyr | Leu | Gly | Phe | Thr | Val | His | Pro | Gly | Asp | Arg | Trp |
| | | | 325 | | | | | 330 | | | | | 335 | | |

Val Leu Glu Thr Gly Arg Leu Tyr Glu Ile Thr Ile Glu Val Phe Asp
 340 345 350
 Lys Phe Ser Asn Lys Val Tyr Val Ser Asp Asn Ile Arg Ile Glu Thr
 355 360 365
 Val Leu Pro Ala Glu Phe Phe Glu Val Leu Ser Ser Ser Gln Asn Gly
 370 375 380
 Ser Tyr His Arg Ile Arg Ala Leu Lys Arg Gly Gln Thr Ala Ile Asp
 385 390 395 400
 Ala Ala Leu Thr Ser Val Val Asp Gln Asp Gly Gly Val His Ile Leu
 405 410 415
 Gln Val Pro Val Trp Asn Gln Gln Glu Val Glu Ile His Ile Pro Ile
 420 425 430
 Thr Leu Tyr Pro Ser Ile Leu Thr Phe Pro Trp Gln Pro Lys Thr Gly
 435 440 445
 Ala Tyr Gln Tyr Thr Ile Arg Ala His Gly Gly Ser Gly Asn Phe Ser
 450 455 460
 Trp Ser Ser Ser Ser His Leu Val Ala Thr Val Thr Val Lys Gly Val
 465 470 475 480
 Met Thr Thr Gly Ser Asp Ile Gly Phe Ser Val Ile Gln Ala His Asp
 485 490 495
 Val Gln Asn Pro Leu His Phe Gly Glu Met Lys Val Tyr Val Ile Glu
 500 505 510
 Pro His Ser Met Glu Phe Ala Pro Cys Gln Val Glu Ala Arg Val Gly
 515 520 525
 Gln Ala Leu Glu Leu Pro Leu Arg Ile Ser Gly Leu Met Pro Gly Gly
 530 535 540
 Ala Ser Glu Val Val Thr Leu Ser Asp Cys Ser His Phe Asp Leu Ala
 545 550 555 560
 Val Glu Val Glu Asn Gln Gly Val Phe Gln Pro Leu Pro Gly Arg Leu
 565 570 575
 Pro Pro Gly Ser Glu His Cys Ser Gly Val Arg Val Lys Ala Glu Ala
 580 585 590
 Gln Gly Ser Thr Thr Leu Leu Val Ser Tyr Arg His Gly His Val His
 595 600 605
 Leu Ser Ala Lys Ile Thr Ile Ala Ala Tyr Leu Pro Leu Lys Ala Val
 610 615 620
 Asp Pro Ser Ser Val Ala Leu Val Thr Leu Gly Ser Ser Lys Glu Met
 625 630 635 640

139

Leu Phe Glu Gly Gly Pro Arg Pro Trp Ile Leu Glu Pro Ser Lys Phe
645 650 655

Phe Gln Asn Val Thr Ala Glu Asp Thr Asp Ser Ile Gly Leu Ala Leu
660 665 670

Phe Ala Pro His Ser Ser Arg Asn Tyr Gln Gln His Trp Ile Leu Val
675 685

Thr Cys Gln Ala Leu Gly Glu Gln Val Ile Ala Leu Ser Val Gly Asn
690 695 700

Lys Pro Ser Leu Thr Asn Pro Phe Pro Ala Val Glu Pro Ala Val Val
705 710 715 720

Lys Phe Val Cys Ala Pro Pro Ser Arg Leu Thr Leu Val Pro Val Tyr
725 730 735

Thr Ser Pro Gln Leu Asp Met Ser Cys Pro Leu Leu Gln Gln Asn Lys
740 745 750

Gln Val Val Pro Val Ser Ser His Arg Asn Pro Leu Leu Asp Leu Ala
755 760 765

Ala Tyr Asp Gln Glu Gly Arg Arg Phe Asp Asn Phe Ser Ser Leu Ser
770 775 780

Ile Gln Trp Glu Ser Thr Arg Pro Val Leu Ala Ser Ile Glu Pro Glu
785 790 795 800

Leu Pro Met Gln Leu Val Ser Gln Asp Asp Glu Ser Gly Gln Lys Lys
805 810 815

Leu His Gly Leu Gln Ala Ile Leu Val His Glu Ala Ser Gly Thr Thr
820 825 830

Ala

<210> 181

<211> 315

<212> PRT

<213> Homo sapiens

<400> 181

Ala Ile Ser Ala Thr Ala Thr Gly Tyr Gln Gln Ser His Leu Ser Ala
1 5 10 15

Ala Gly Val Lys Gln Leu Arg Asp Pro Leu Val Pro Val Ser Ala Ser
20 25 30

Ile Glu Leu Ile Leu Val Glu Asp Val Arg Val Ser Pro Glu Glu Val
35 40 45

Thr Ile Tyr Asn His Pro Gly Val Gln Val Glu Leu His Ile Thr Glu
50 55 60

140

Gly Ser Gly Tyr Phe Phe Leu Asn Thr Ser Thr Gln Asp Ile Ile Asn
 65 70 75 80
 Val Ala Tyr Gln Asp Thr Arg Gly Val Ala Met Val His Pro Leu Phe
 85 90 95
 Pro Gly Ser Ser Thr Val Met Val His Asp Leu Cys Leu Thr Phe Pro
 100 105 110
 Ala Pro Ala Lys Ala Thr Ile His Val Ser Asp Ile Gln Glu Leu Tyr
 115 120 125
 Val Arg Val Val Asp Lys Val Glu Ile Gly Lys Ala Val Lys Ala Tyr
 130 135 140
 Val Arg Val Leu Asp Phe Tyr Lys Lys Pro Phe Leu Ala Lys Tyr Phe
 145 150 155 160
 Thr Phe Met Asp Leu Lys Leu Arg Ala Ala Ser Gln Ile Ile Thr Leu
 165 170 175
 Val Thr Leu Asp Glu Ala Leu Asp Asn Tyr Thr Ala Thr Phe Leu Val
 180 185 190
 His Gly Val Ala Ile Gly Gln Thr Ser Leu Ser Ala Ser Val Thr Asp
 195 200 205
 Lys Ser Gly Gln Arg Val Ser Ser Thr Ala Gln Gln Ile Glu Val Phe
 210 215 220
 Pro Pro Phe Arg Leu Ile Pro Arg Lys Val Thr Leu Ile Ile Gly Ala
 225 230 235 240
 Met Ile Gln Ile Thr Ser Glu Gly Gly Pro Gln Pro Gln Ser Asn Ile
 245 250 255
 Leu Phe Ser Ile Asn Asn Glu Ser Val Ala Ala Val Ser Ser Ala Gly
 260 265 270
 Leu Val Arg Gly Leu Met Val Gly Asn Gly Ser Val Leu Gly Val Val
 275 280 285
 Gln Ala Val Asp Ala Glu Thr Gly Lys Val Ile Ile Val Ser Gln Asp
 290 295 300
 His Val Glu Val Glu Val Leu Gln Leu Gln Ala
 305 310 315

<210> 182

<211> 315

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (17)

<223> Xaa equals any of the naturally occurring L-amino acids

141

<220>

<221> SITE

<222> (217)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 182

Ser Ile Thr Ala Thr Ala Thr Gly Tyr Gln Glu Ser His Leu Ser Ser
 1 5 10 15

Xaa Arg Thr Lys Gln Pro His Asp Pro Leu Val Pro Leu Ser Ala Ser
 20 25 30

Ile Glu Leu Ile Leu Val Glu Asp Val Arg Val Ser Pro Glu Glu Val
 35 40 45

Thr Ile Tyr Asn His Pro Gly Ile Gln Ala Glu Leu Arg Ile Arg Glu
 50 55 60

Gly Ser Gly Tyr Phe Phe Leu Asn Thr Ser Thr Ala Asp Val Val Lys
 65 70 75 80

Val Ala Tyr Gln Glu Ala Arg Gly Val Ala Met Val His Pro Leu Leu
 85 90 95

Pro Gly Ser Ser Thr Ile Met Ile His Asp Leu Cys Leu Val Phe Pro
 100 105 110

Ala Pro Ala Lys Ala Val Val Tyr Val Ser Asp Ile Gln Glu Leu Tyr
 115 120 125

Ile Arg Val Val Asp Lys Val Glu Ile Gly Lys Thr Val Lys Ala Tyr
 130 135 140

Val Arg Val Leu Asp Leu His Lys Lys Pro Phe Leu Ala Lys Tyr Phe
 145 150 155 160

Pro Phe Met Asp Leu Lys Leu Arg Ala Ala Ser Pro Ile Ile Thr Leu
 165 170 175

Val Ala Leu Asp Glu Ala Leu Asp Asn Tyr Thr Ile Thr Phe Leu Ile
 180 185 190

Arg Gly Val Ala Ile Gly Gln Thr Ser Leu Thr Ala Ser Val Thr Asn
 195 200 205

Lys Ala Gly Gln Arg Ile Asn Ser Xaa Pro Gln Gln Ile Glu Val Phe
 210 215 220

Pro Pro Phe Arg Leu Met Pro Arg Lys Val Thr Leu Leu Ile Gly Ala
 225 230 235 240

Thr Met Gln Val Thr Ser Glu Gly Gly Pro Gln Pro Gln Ser Asn Ile
 245 250 255

Leu Phe Ser Ile Ser Asn Glu Ser Val Ala Leu Val Ser Ala Ala Gly
 260 265 270

142

Leu Val Gln Gly Leu Ala Ile Gly Asn Gly Thr Val Ser Gly Leu Val
 275 280 285

Gln Ala Val Asp Ala Glu Thr Gly Lys Val Val Ile Ile Ser Gln Asp
 290 295 300

Leu Val Gln Val Glu Val Leu Leu Leu Arg Ala
 305 310 315

<210> 183

<211> 47

<212> PRT

<213> Homo sapiens

<400> 183

Pro Leu Ala Leu Thr Pro His Ala Ser Pro Gln His Ser Pro His Tyr
 1 5 10 15

Leu Ala Ser Ser Pro Thr Ala Phe Asn Thr Leu Pro Ser Asp Arg Lys
 20 25 30

Ala Ser Pro Pro Ser Gly Leu Trp Ser Pro Ala Tyr Ala Ser His
 35 40 45

<210> 184

<211> 46

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (27)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (34)

<223> Xaa equals stop translation

<400> 184

Pro Phe Trp Phe Thr Arg His Gln Glu Pro Gln His His Cys His Cys
 1 5 10 15

His Trp Leu Pro Gly Val Pro Pro Gln Leu Xaa Gln Asn Lys Ala Ala
 20 25 30

Ala Xaa Pro Ser Gly Ala Ser Val Gly Leu His Arg Ala His
 35 40 45

<210> 185

<211> 94

<212> PRT

<213> Homo sapiens

<400> 185

143

Val Glu Ser Glu Leu Leu His Thr Tyr Ser Lys Val Asp Pro Phe Asp
 1 5 10 15
 Val Leu Ile Leu Cys Val Arg Val Ala Val Leu Thr Ala Val Thr Leu
 20 25 30
 Thr Val Pro Ile Val Leu Phe Pro Val Arg Arg Ala Ile Gln Gln Met
 35 40 45
 Leu Phe Pro Asn Gln Glu Phe Ser Trp Leu Arg His Val Leu Ile Ala
 50 55 60
 Val Gly Leu Leu Thr Cys Ile Asn Leu Leu Val Ile Phe Ala Pro Asn
 65 70 75 80
 Ile Leu Gly Ile Phe Gly Val Ile Gly Ala Thr Ser Ala Pro
 85 90

<210> 186

<211> 91

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (38)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (43)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (51)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (83)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 186

Val Lys Ala Glu Met Leu His Met Tyr Ser Gln Lys Asp Pro Leu Ile
 1 5 10 15
 Leu Cys Val Arg Leu Ala Val Leu Leu Ala Val Thr Leu Thr Val Pro
 20 25 30
 Val Val Leu Phe Pro Xaa Arg Arg Ala Leu Xaa Gln Leu Leu Phe Pro
 35 40 45
 Gly Lys Xaa Phe Ser Trp Pro Arg His Val Ala Ile Ser Ser Asp Leu
 50 55 60
 Leu Val Trp Ser Met Ser Leu Ser Ser Val Cys Gln His Pro Gly Tyr

144

65 70 75 80
 Leu Trp Xaa Leu Ile Gly Ser Thr Gln Ala Pro
 85 90

<210> 187
 <211> 374
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (320)
 <223> Xaa equals any of the naturally occurring L-amino acids

<400> 187
 Glu Lys Pro Tyr Leu Val Glu Glu Ala Val Ser Tyr Asn Glu Leu Asp
 1 5 10 15
 Tyr Val Ser Val Gly Leu Asp Gln Gln Thr Val Lys Leu Val Cys Thr
 20 25 30
 Asn Arg Arg Lys Gln Phe Leu Leu Asp Thr Ala Asp Val Ala Leu Ala
 35 40 45
 Glu Phe Phe Leu Ala Ser Leu Lys Ser Ala Met Ile Lys Gly Cys Arg
 50 55 60
 Glu Pro Pro Tyr Pro Ser Ile Leu Thr Asp Ala Thr Met Glu Lys Leu
 65 70 75 80
 Ala Leu Ala Lys Phe Val Ala Gln Glu Ser Lys Cys Glu Ala Ser Ala
 85 90 95
 Val Thr Val Arg Phe Tyr Gly Leu Val His Trp Glu Asp Pro Thr Asp
 100 105 110
 Glu Ser Leu Gly Pro Thr Pro Cys His Cys Ser Pro Pro Glu Gly Thr
 115 120 125
 Ile Thr Lys Glu Gly Met Leu His Tyr Lys Ala Gly Thr Ser Tyr Leu
 130 135 140
 Gly Lys Glu His Trp Lys Thr Cys Phe Val Val Leu Ser Asn Gly Ile
 145 150 155 160
 Leu Tyr Gln Tyr Pro Asp Arg Thr Asp Val Ile Pro Leu Leu Ser Val
 165 170 175
 Asn Met Gly Gly Glu Gln Cys Gly Gly Cys Arg Arg Ala Asn Thr Thr
 180 185 190
 Asp Arg Pro His Ala Phe Gln Val Ile Leu Ser Asp Arg Pro Cys Leu
 195 200 205
 Glu Leu Ser Ala Glu Ser Glu Ala Glu Met Ala Glu Trp Met Gln His
 210 215 220

145

Leu Cys Gln Ala Val Ser Lys Gly Val Ile Pro Gln Gly Val Ala Pro
 225 230 235 240
 Ser Pro Cys Ile Pro Cys Cys Leu Val Leu Thr Asp Asp Arg Leu Phe
 245 250 255
 Thr Cys His Glu Asp Cys Gln Thr Ser Phe Phe Arg Ser Leu Gly Thr
 260 265 270
 Ala Lys Leu Gly Asp Ile Ser Ala Val Ser Thr Glu Pro Gly Lys Glu
 275 280 285
 Tyr Cys Val Leu Glu Phe Ser Gln Asp Ser Gln Gln Leu Leu Pro Pro
 290 295 300
 Trp Val Ile Tyr Leu Ser Cys Thr Ser Glu Leu Asp Arg Leu Leu Xaa
 305 310 315 320
 Ala Leu Asn Ser Gly Trp Lys Thr Ile Tyr Gln Val Asp Leu Pro His
 325 330 335
 Thr Ala Ile Gln Glu Ala Phe Asn Lys Lys Lys Phe Glu Asp Ala Leu
 340 345 350
 Ser Leu Ile His Ser Ala Trp Gln Arg Ser Asp Ser Leu Cys Arg Gly
 355 360 365
 Arg Ala Ser Glu Thr Pro
 370

<210> 188

<211> 145

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (21)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (35)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (78)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (80)

<223> Xaa equals stop translation

<220>

146

<221> SITE
 <222> (85)
 <223> Xaa equals stop translation

<220>
 <221> SITE
 <222> (93)
 <223> Xaa equals stop translation

<400> 188
 Tyr Arg Phe Asn Ala Ile Pro Ile Lys Ile Pro Met Ala Phe Phe Met
 1 5 10 15
 Glu Ile Glu Lys Xaa Asn Leu Lys Phe Ile Trp Asn His Lys Gly Pro
 20 25 30
 Gln Ile Xaa Lys Thr Ile Leu Arg Lys Lys Asn Lys Gly Gly Gly Phe
 35 40 45
 Ile Phe Pro Asp Phe Lys Val Tyr Tyr Lys Ala Thr Val Ile Lys Thr
 50 55 60
 Val Arg Tyr Trp Gln Lys Asp Arg His Ile Thr Asn Gly Xaa Lys Xaa
 65 70 75 80
 Arg Ala Gln Lys Xaa Thr His Ala Asn Met Phe Asn Xaa Phe Ser Thr
 85 90 95
 Arg Val Pro Gly Thr His Asn Glu Ala Arg Ile Val Ser Ser Thr Asn
 100 105 110
 Thr Trp Glu Asn Trp Met Ser Thr Cys Lys Arg Met Lys Phe Asp Pro
 115 120 125
 Tyr Leu Thr Leu Tyr Thr Lys Ile Tyr Ser Lys Trp Thr Lys Asp Leu
 130 135 140
 Ser
 145

<210> 189
 <211> 311
 <212> PRT
 <213> Homo sapiens

<400> 189
 Gly Glu Gly Ile Leu Leu Ser Gly Gly Asp Lys Trp Ser Arg His Arg
 1 5 10 15
 Arg Met Leu Thr Pro Ala Phe His Phe Asn Ile Leu Lys Ser Tyr Ile
 20 25 30
 Thr Ile Phe Asn Lys Ser Ala Asn Ile Met Leu Asp Lys Trp Gln His
 35 40 45
 Leu Ala Ser Glu Gly Ser Ser Cys Leu Asp Met Phe Glu His Ile Ser
 50 55 60

147

Leu Met Thr Leu Asp Ser Leu Gln Lys Cys Ile Phe Ser Phe Asp Ser
 65 70 75 80
 His Cys Gln Glu Arg Pro Ser Glu Tyr Ile Ala Thr Ile Leu Glu Leu
 85 90 95
 Ser Ala Leu Val Glu Lys Arg Ser Gln His Ile Leu Gln His Met Asp
 100 105 110
 Phe Leu Tyr Tyr Leu Ser His Asp Gly Arg Arg Phe His Arg Ala Cys
 115 120 125
 Arg Leu Val His Asp Phe Thr Asp Ala Val Ile Arg Glu Arg Arg Arg
 130 135 140
 Thr Leu Pro Thr Gln Gly Ile Asp Asp Phe Phe Lys Asp Lys Ala Lys
 145 150 155 160
 Ser Lys Thr Leu Asp Phe Ile Asp Val Leu Leu Leu Ser Lys Asp Glu
 165 170 175
 Asp Gly Lys Ala Leu Ser Asp Glu Asp Ile Arg Ala Glu Ala Asp Thr
 180 185 190
 Phe Met Phe Gly Gly His Asp Thr Thr Ala Ser Gly Leu Ser Trp Val
 195 200 205
 Leu Tyr Asn Leu Ala Arg His Pro Glu Tyr Gln Glu Arg Cys Arg Gln
 210 215 220
 Glu Val Gln Glu Leu Leu Lys Asp Arg Asp Pro Lys Glu Ile Glu Trp
 225 230 235 240
 Asp Asp Leu Ala Gln Leu Pro Phe Leu Thr Met Cys Val Lys Glu Ser
 245 250 255
 Leu Arg Leu His Pro Pro Ala Pro Phe Ile Ser Arg Cys Cys Thr Gln
 260 265 270
 Asp Ile Val Leu Pro Asp Gly Arg Val Ile Pro Lys Gly Ile Thr Cys
 275 280 285
 Leu Ile Asp Ile Ile Gly Val His His Asn Pro Thr Val Trp Pro Asp
 290 295 300
 Pro Glu Val Tyr Asp Pro Phe
 305 310

<210> 190

<211> 312

<212> PRT

<213> Homo sapiens

<400> 190

Gly Glu Gly Ile Leu Leu Ser Gly Gly Asp Lys Trp Ser Arg His Arg
 1 5 10 15

148

Arg Met Leu Thr Pro Ala Phe His Phe Asn Ile Leu Lys Ser Tyr Ile
 20 25 30
 Thr Ile Phe Asn Lys Ser Ala Asn Ile Met Leu Asp Lys Trp Gln His
 35 40 45
 Leu Ala Ser Glu Gly Ser Ser Arg Leu Asp Met Phe Glu His Ile Ser
 50 55 60
 Leu Met Thr Leu Asp Ser Leu Gln Lys Cys Ile Phe Ser Phe Asp Ser
 65 70 75 80
 His Cys Gln Glu Arg Pro Ser Glu Tyr Ile Ala Thr Ile Leu Glu Leu
 85 90 95
 Ser Ala Leu Val Glu Lys Arg Ser Gln His Ile Leu Gln His Met Asp
 100 105 110
 Phe Leu Tyr Tyr Leu Ser His Asp Gly Arg Arg Phe His Arg Ala Cys
 115 120 125
 Arg Leu Val His Asp Phe Thr Asp Ala Val Ile Arg Glu Arg Arg Arg
 130 135 140
 Thr Leu Pro Thr Gln Gly Ile Asp Asp Phe Phe Lys Asp Lys Ala Lys
 145 150 155 160
 Ser Lys Thr Leu Asp Phe Ile Asp Val Leu Leu Leu Ser Lys Asp Glu
 165 170 175
 Asp Gly Lys Ala Leu Ser Asp Glu Asp Ile Arg Ala Glu Ala Asp Thr
 180 185 190
 Phe Met Phe Gly Gly His Asp Thr Thr Ala Ser Gly Leu Ser Trp Val
 195 200 205
 Leu Tyr Asn Leu Ala Arg His Pro Glu Tyr Gln Glu Arg Cys Arg Gln
 210 215 220
 Glu Val Gln Glu Leu Leu Lys Asp Arg Asp Pro Lys Glu Ile Glu Trp
 225 230 235 240
 Asp Asp Leu Ala Gln Leu Pro Phe Leu Thr Met Cys Val Lys Glu Ser
 245 250 255
 Leu Arg Leu His Pro Pro Ala Pro Phe Ile Ser Arg Cys Cys Thr Gln
 260 265 270
 Asp Ile Val Leu Pro Asp Gly Arg Val Ile Pro Lys Gly Ile Thr Cys
 275 280 285
 Leu Ile Asp Ile Ile Gly Val His His Asn Pro Thr Val Trp Pro Asp
 290 295 300
 Pro Glu Val Leu Pro Ser Pro Phe
 305 310

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/30628

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 536/23.1, 23.5, 24.31; 530/300, 350; 435/6, 69.1, 252.3, 320.1, 325

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|---|-----------------------|
| X, P | Database GenEmbl; Accession NO: Y56966; MICHALOVICH et al. "New polypeptides of prosaposin family, polynucleotides useful for identifying agonist, antagonists and inhibitors for treatment of neuronal degeneration, psychiatric disorders, stroke etc."; 03 February 2000, having 98.6% sequence identity with SEQ ID NO: 91; see entire document. | 1-7, 21 |



Further documents are listed in the continuation of Box C.



See patent family annex.

| | |
|---|--|
| * Special categories of cited documents: | *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention |
| *A* document defining the general state of the art which is not considered to be of particular relevance | *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone |
| *E* earlier document published on or after the international filing date | *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art |
| *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) | *Z* document member of the same patent family |
| *O* document referring to an oral disclosure, use, exhibition or other means | |
| *P* document published prior to the international filing date but later than the priority date claimed | |

Date of the actual completion of the international search

24 FEBRUARY 2001

Date of mailing of the international search report

26 APR 2001

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

RITA MITRA

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/30628

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|--|-----------------------|
| X, P | WO 00/05361 A1 (SMITHKLINE BEECHAM, P.L.C.) 03 February 2000, "Human SBPSAPL gene with homology to the prosaposin family of neurotrophic factors"; relates to SBPSAPL polynucleotides encoding the polypeptides; recombinants and methods for their production, methods for isolating polypeptides expressed by such recombinants; vector: pQE; host cell: bacteria and animal cells; see entire document. | 1-10, 14, 15, 21 |
| X, P | Database: N_Genseq_36; Accession NO: Z56885; MICHALOVICH et al. "This DNA encodes a human SBPSAPL polypeptide having homology to the human prosaposin family"; 03 February 2000; having 99.8% sequence identity with SEQ ID NO: 11; see entire document. | 1-7, 21 |
| X, P | Database: GenEmbl; Accession NO: AC004578; STONE et al. "Homo sapiens chromosome 4"; 16 November 1999; having 99.8% sequence identity to SEQ ID NO: 11; see entire document. | 1-7, 21 |
| X | Database: EST; Accession NO: AA429294; HILLIER et al. 16 October 1997, "Soares_NhHMPu_S1 Homo sapiens cDNA clone"; having 98.8% sequence identity to SEQ ID NO: 11; vector: pT7T3D-Pac; host cell: DH10B; see entire document. | 1-10, 21 |

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/30628

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-10, 14, 15 and 21 {all in part}

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/30628

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (7):

CO7H 21/04; CO7H 21/02; CO7K 5/00; CO7K 14/00; C12Q 1/68; C12N 1/21; C12N 15/63; C12N 15/85, 15/86

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

536/23.1, 23.5, 24.31; 530/300, 350; 435/6, 69.1, 252.3, 320.1, 325

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

EAST, STN (Database: GenEmbl, N_GenSeq_36, Issued_Patents_N, EST, A_GenSeq_36, PIR_65, SwissProt_39, Sptrembl_14

Search Terms: sulfatide activator protein, SAP-I, saposin, prosaposin, leukodystrophy, SBPSAPL polypeptide

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Groups 1-80, claim(s) 1-10, 14, 15 and 21, all in part, drawn to an isolated nucleic acid of SEQ ID NO X or a peptide of SEQ ID NO Y, wherein X and Y are values that correlate to those listed in Table 1, and correspond to one of the cDNA Clone IDs, respectively. For examples,

If group 1 is elected, this correlates to Gene no 1, cDNA clone ID HWEAC77 of Table 1, wherein X is 11 and Y is 91.

If group 2 is elected, this correlates to Gene No 2, cDNA clone ID HRAEH37, wherein X is 12 and Y is 92.

Groups 81-161, claim(s) 11, 12, 16 and 23, all in part, each group directed to a peptide of SEQ ID NO; Y, wherein Y correlates to one of those listed in Table 1, and corresponds to one of the cDNA Clone IDs, respectively. For examples,

If group 81 is elected, this correlates to Gene no 1, cDNA clone ID HWEAC77 of Table 1, wherein Y is 91.

If group 82 is elected, this correlates to Gene No 2, cDNA clone ID HRAEH37, wherein Y is 92.

Groups 161-240, claim 13, in part, drawn to an isolated antibody which binds to a protein with SEQ ID NO Y, wherein Y correlates to one of those listed in Table 1, and corresponds to one of the cDNA Clone IDs, respectively. For examples,

If group 161 is elected, this correlates to Gene no 1, cDNA clone ID HWEAC77, of Table 1, wherein Y is 91.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/30628

If group 162 is elected, this correlates to Gene No 2, cDNA clone ID HRAEH37, wherein Y is 92.

Groups 241-320, claim 17, in part, drawn to a method for preventing, treating or ameliorating an undefined medical condition by administering a polypeptide of SEQ ID NO Y, wherein Y correlates to one of those listed in Table 1, and corresponds to one of the cDNA Clone IDs, respectively. For examples,

If group 241 is elected, this correlates to Gene no 1, cDNA clone ID HWEAC77 of Table 1, wherein Y is 91.

If group 242 is elected, this correlates to Gene No 2, cDNA clone ID HRAEH37, wherein Y is 92.

Groups 321-400, claim 17, in part, drawn to a method for preventing, treating or ameliorating an undefined medical condition by administering a polynucleotide of SEQ ID NO X encoding a protein of SEQ ID NO Y, wherein X and Y correlate to one of those listed in Table 1, and correspond to one of the cDNA Clone IDs, respectively. For examples,

If group 321 is elected, this correlates to Gene no 1, cDNA clone ID HWEAC77 of Table 1, wherein X is 11 and Y is 91.

If group 322 is elected, this correlates to Gene No 2, cDNA clone ID HRAEH37, wherein X is 12 and Y is 92.

Groups 401-480, claims 18 and 19, in part, drawn to a method of diagnosis of an undefined pathological condition by determining the presence or absence of a mutation in a polynucleotide of SEQ ID NO X, wherein X correlates to one of those listed in Table 1, and corresponds to one of the cDNA Clone IDs, respectively. For examples,

If group 401 is elected, this correlates to Gene no 1, cDNA clone ID HWEAC77 of Table 1, wherein X is 11.

If group 402 is elected, this correlates to Gene No 2, cDNA clone ID HRAEH37, wherein X is 12.

Groups 481-560, claim 20, in part, drawn to a method of identifying a binding partner to a polypeptide defined by SEQ ID NO Y, wherein Y correlates to one of those listed in Table 1, and corresponds to one of the cDNA Clone IDs, respectively. For examples,

If group 481 is elected, this correlates to Gene no 1, cDNA clone ID HWEAC77 of Table 1, wherein Y is 91.

If group 638 is elected, this correlates to Gene No 2, cDNA clone ID HRAEH37, wherein Y is 92.

Groups 561-640, claim 22, in part, drawn to a method of identifying an activity in a biological assay by identification of the protein in the supernatant wherein the cell expresses a polypeptide encoded by SEQ ID NO X, wherein X correlates to one of those listed in Table 1, and corresponds to one of the cDNA Clone IDs, respectively. For examples,

If group 561 is elected, this correlates to Gene no 1, cDNA clone ID HWEAC77 of Table 1, wherein X is 11.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/30628

If group 562 is elected, this correlates to Gene No 2, cDNA clone ID HRAEH37, wherein X is 12.

The inventions listed as Groups 1-640 do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

The polynucleotides and polypeptides of each of the clones in Table 1 are unrelated, each to the other. The polynucleotide sequences encode structurally distinct polypeptides and do not share a special technical feature.

Furthermore, the technical feature that links the DNA, protein, antibody, methods of CDNA clone HWEAC77 (see Table 1) is not a contribution over the prior art. See the various documents cited in the search report. Thus the technical feature of the polynucleotide sequence is not special and the groups are not so linked under PCT Rule 13.1. Additionally the claimed methods produce different products and/or different results which are not coextensive and which do not share the same technical feature.

THIS PAGE BLANK (USPTO)